

# International Journal of Pharmacy and Pharmaceutical Sciences Vol 7, Issue 3, 2015

*by* Artikel 6 Cytotoxicity Studies Of Tetraprelyltoluquinone, A

---

**Submission date:** 15-Aug-2018 12:01PM (UTC+0800)

**Submission ID:** 990073762

**File name:** droquinone\_from\_Garcina\_cowa\_Roxb\_on\_H-460,\_MCF-7\_and\_DU-145.pdf (204.98K)

**Word count:** 2496

**Character count:** 12915

Original Article

CYTOTOXICITY STUDIES OF TETRAPRENYLTOLUQUINONE, A PRENYLATED HYDROQUINONE FROM *GARCINIA COWA* ROXB ON H-460, MCF-7 AND DU-145

FATMA SRI WAHYUNI<sup>1</sup>, JOHNSON STANSLAS<sup>3</sup>, NORDIN HJ. LAJIS<sup>2</sup>, DACHRIYANUS<sup>1</sup>

<sup>1</sup>Faculty of Pharmacy, Andalas University, Kampus Limau Manis, Padang, West Sumatra, Indonesia, 25163, <sup>2</sup>Laboratory of Natural Products, Institute of Bioscience, University Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia, <sup>3</sup>Faculty of Medicine and Health Sciences, University Putra Malaysia 43400 UPM, Serdang, Selangor, Malaysia.  
Email: dachriyanus@ffarmasi.unand.ac.id

Received: 23 Oct 2014 Revised and Accepted: 15 Nov 2014

ABSTRACT

**Objective:** The aim of the present study was to examine the cytotoxicity of a new ring-reduced tetra prenyltoluquinone (TPTQ), [2E,6E,10E]-(+)-4-hydroxy-3-methyl-5-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenyl)-2-cyclohexen-1-one against H-460, MCF-7 and DU-145 cell lines.

**Methods:** Different concentrations of TPTQ were subjected to cytotoxicity study by using MTT method and calculate the percentage of cell viability.

**Results:** The results of this study showed that this compound has IC<sub>50</sub> value 16.3 ± 3.0 µM in H-460 cancer cell lines without any activities towards another two type of cell lines.

**Conclusion:** TPTQ had selective activity against H-460 cancer cell lines.

**Keywords:** *Garcinia cowa*, TPTQ, H-460, MCF-7, DU-145, Cell viability, Cytotoxicity.

INTRODUCTION


Cancer can be defined as a disease in which the disorder occurs in the normal processes of cell division, which are controlled by the genetic material (DNA) of the cell [1]. Several plant-derived compounds are currently successfully employed in cancer treatment such as vincristine (leukemia, lymphoma, breast, lung, solid cancers and others), vinblastine (breast, lymphoma, germ-cell and renal cancer), paclitaxel (ovary, breast, lung, bladder, head and neck cancer), docetaxel (breast and lung cancer) [2].

Anticancer drugs are not selective, because they can damage normal cells that have rapid growth. It encourages researchers to conduct research in order to find an anticancer drug that has selective activity against cancer cells without damaging the normal cells.

Plants are the potential source for discovering biologically active compounds preferably those having unique and new mechanism of therapeutic action. Among of these plants, special attention is paid to genus *Garcinia* (Guttiferae). This genus has been extensively investigated from phyto chemical and biological point of view. Prenylated xanthenes, triterpenes and bio flavonoids have been isolated from this genus. The alcoholic extract of *Garcinia* species were reported to have the wide range of biological activities such as anti-inflammatory, antioxidant, antibacterial and cytotoxicity [3-6].

*Garcinia cowa* Roxb. (Guttiferae), known as kandis in West Sumatra, is a medium-sized tree which attains a height of ca. 30 m and it is widely distributed throughout Indonesia and the Malay peninsula. Traditionally, kandis have been used for many purposes. The bark of kandis is used as pesticide, larva side [7] and antipyretic [8] while the sun dried fruit is used to treat dysentery [9]. Dried leaves of this species are used as a tonic [10]. Biological activity studies on this species showed that the leaves had strong antitumor promoting activity on cells-Raji [11], while the dried steam bark had antimalarial activity [12]. Previous workers have isolated a number of prenylated xanthenes from the stem bark [3, 6, 8, 12]. Previous study showed that the hexane fraction of the steam bark of *G. cowa* was active against small lung cell cancer H-460 and not active against MCF-7 and DU-145 cell lines [13]. A new ring-reduced tetraprenyltoluquinone (TPTQ), [2E,6E,10E]-(+)-4-hydroxy-3-methyl-5-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenyl)-2-cyclohexen-1-one was isolated from this species [14]. Based on these data, further cytotoxicity activity studies of TPTQ on three cancer

cell lines, H-460, MCF-7 and DU-145 cell lines were carried out on this compound.



MATERIALS AND METHODS

Instruments

UV (in absolute ethanol) and IR (KBr) spectra were recorded on a JASCO V-560 spectrophotometer and a Perkin-Elmer 1650 FTIR spectrophotometer, respectively. High resolution fast-atom bombardment mass spectrometry (HRFAB-MS) was obtained on a JEOL JMS HX-110A spectrometer. <sup>1</sup>H and <sup>13</sup>C NMR spectra (CDCl<sub>3</sub>) were recorded on a Varian 500 MHz NMR Spectrometer at 500 MHz (<sup>1</sup>H) and 125 MHz (<sup>13</sup>C), respectively. The spectra was interpreted with the aid of the 1H-1H COSY, HMBC, and HMQC techniques.

Holten Lamin Air microbiological safety cabinet class II was obtained from Heto-Holten (Allerød, Denmark), and Galaxy CO<sub>2</sub> incubator was purchased from RS Biotech (Ayrshire, Scotland). A micro plate reader equipped SOFTmax® Pro software (Versamax, Molecular Devices, California, USA) was used to measure of the formazan solution.

Plant material

*Garcinia cowa* Roxb (stem bark) was collected at Sarasah Bonta, Harau Valley, and West Sumatra at an altitude of 500 m. Plant materials were cut into small pieces (3-5 mm thickness) and air-dried under shade. The dried plant materials were ground to powder before extraction.

Cell lines

The cell lines, NCI-H460 were purchased from the American Type Culture Collection (Manassas, VA, USA). The cancer cells were cultured in RPMI 1640 medium (Life Technologies, Paisly, UK) with 10% v/v fetal calf serum (PAA Laboratories, Linz, Austria), 100 IU/ml penicillin and 100 µg/ml streptomycin (Life Technologies,

Paisly, UK) whereas the solution trypsin-EDTA were purchased from GIBCO (Auckland, New Zealand). Dimethylsulfoxide (DMSO) was purchased from BDH Laboratory (England). Culture flask (25 cm<sup>2</sup> and 75 cm<sup>2</sup>) and 10 ml serological pipettes were purchased from Becton Dickinson (New Jersey, USA).

#### Isolation of TPTQ from the stem bark of *G. cowa*

The ground air-dried stem bark (2.2 Kg) was percolated three times with methanol (10 l). The extract was evaporated to dryness under reduced pressure to give a dark mass (113.54 g). The methanol extract was partitioned between water 200 ml and hexane (4 x 100 ml) and the combined hexane extracts were evaporated to yield a gum (5.7 g). The aqueous layer was further partitioned successively with ethyl acetate and followed by butanol, and upon evaporation 43.0 g and 45.0 g of gums were obtained from the respective solvents.

A portion (4.0 g) of the hexane extract was subjected to CC (silica gel, Merck 7734) with hexane-ethyl acetate mixture as eluent in increasing polarity manner, followed by methanol to give three sub-fractions HxA-HxC. Sub-fraction HxB (0.896 g) was further purified by silica gel CC (Merck 9385) eluting with a mixture of hexane-ethyl acetate mixture in increasing polarity mode to give four sub-fractions (Habra-HxBd). Further purification of HxBb (0.550 mg) was achieved using radial chromatography using the same eluent to yield 310 mg of [2E,6E,10E]-(+)-4-hydroxy-3-methyl-5-(3,7,11,15-tetramethyl-2,6,10,14-hexadecatetraenyl-2-cyclohexen-1-one) as a yellowish oil. High-resolution FABMS gave [M+H]<sup>+</sup> at: 399.3263 C<sub>27</sub>H<sub>43</sub>O<sub>2</sub>, requires *m/z* 399.3359, I<sup>10</sup> +50.0°. UV (MeOH) max (log<sub>e</sub>) nm: 230 (4.04); IR max(nujol) cm<sup>-1</sup>: 3391, 2982, 2909, 2834, 1652, 1436, 1375, 1330, 1289, 1186, 1144, 1110, 1055, 1021, 1005, 3435, 3310, 2980, 2925, 2860, 1653, 1437, 1377, 1111, 1008, 886, 667. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ<sub>H</sub>: 2.03 (3H, s, 3-CH<sub>3</sub>), 5.80 (1H, s, H-2), 4.09 (1H, d, *J* = 6.0 Hz, H-4), 5.2 (1H, t, *J* = 6.0 Hz, H-2'), 5.09 (3H, d, *J* = 6.0 Hz, H-6', H-10', H-14'), 2.1 (1H, d, *J* = 8 Hz, H-1'), 2.35 (1H, d, *J* = 8.0 Hz, H-1'), 2.10 (1H, m, H-5), 2.51 (1H, s, H-6), 1.96 (3H, m, H-4', H-8', H-12'), 2.14 (3H, m, H-5', H-9', H-13'). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ<sub>C</sub>: 199.9 (C-1); 126.4 (C-2), 164.1 (C-3), 20.4 (C-3Me), 73.1 (C-4), 43.5 (C-5), 41.1 (C-6), 30.5 (C-1'), 120.5 (C-2'), 138.2 (C-3'), 39.6 (C-5'), C-8'), 39.8 (C-4'), 26.4 (C-13'), 26.5 (C-12'), 26.6 (C-9), 123.8 (C-14'), 124.1 (C-10'), 124.3 (C-6').

#### Culture of cells

The lung cancer (H-460), breast cancer (MCF-7), and prostate cancer (DU-145) cell lines had been used in this study. All cell lines were maintained in RPMI-1640 medium with 10% v/v foetal calf serum, 100 IU/ml penicillin and 100 g/ml streptomycin as a complete medium. Cells were grown in 25 cm<sup>2</sup> flask with 10 ml of complete media at 37°C with 5% CO<sub>2</sub>. Every 3 days the cells were subcultured by splitting the culture with fresh complete media. Cells were attached by incubating suspension cells (180 ul in each well except blank) for overnight.

#### MTT assay [15]

Varying concentrations of TPTQ were prepared from the stock solutions by serial dilution (100 μM, 10 μM, 1 μM, 0.1 μM) in RPMI-1640 to give a volume of 200 μl in each well of the microtiter plate (96-well). The assay for each concentration was performed in quadruplicate and the culture plate was incubated at 37 °C with 5 % (v/v) CO<sub>2</sub> for 96 hours. After of incubation, 50 μL of 2 mg/ml MTT solution was added to each well and allowed to incubate for a further 4 hours, after which all supernatant were discarded. Subsequently, 100 μL DMSO was added to each well and vigorously mixed to dissolve the formazan crystals. Absorbance values at 550 nm were measured with a microplate reader. Cytotoxicity was expressed as IC<sub>50</sub>, i. e. The concentration of the test sample to reduce the absorbance of treated cells by 50% with reference to the control (untreated cells). To determine the IC<sub>50</sub> value, viability vs. dose response curve is plotted. The viability was calculated according to this formula:

$$\% \text{ Viability} = (\text{Optical density of treated cells} / \text{Optical density of control}) \times 100\%$$

The relation, between the concentrations of the test solution with viability cell was displayed in graphical form. Of the graph was a determined price IC<sub>50</sub> (concentrations that inhibit 50% living cells) test solution.

The data of relation between the concentration of the test preparations and the absorbance were statistically analyzed using a one-way variance analysis (ANOVA) followed by of Duncan's multiple regions (Duncan's Multiple Range Test).





Fig. 1: FTIR spectrum of TPTQ in nujol

## RESULT AND DISCUSSION

## TPTQ

Chromatography of the hexane soluble portion of the methanol extract yielded yellowish oil, which was assigned structure **TPTQ** on the following grounds. The high-resolution mass spectrum (HRMS) indicated the molecular formula  $C_{27}H_{42}O_2$ , implying seven unsaturation.



The infrared spectrum (Fig. 1) of compound TPTQ indicated the presence of a hydroxyl function ( $\nu_{\max}$  3435  $\text{cm}^{-1}$ ) and an  $\alpha,\beta$ -unsaturated ketone ( $\nu_{\max}$  1653  $\text{cm}^{-1}$ ) in keeping with the  $^{13}\text{C}$  NMR spectrum (Fig. 2) at  $\delta_{\text{C}}$  199.0 for ketone, a tertiary vanillic carbon ( $\delta_{\text{C}}$  126.4) and a quaternary vanillic carbon ( $\delta_{\text{C}}$  161.1).

The  $^1\text{H}$  NMR (fig. 3) spectrum of side ring contain six allylic methylenes as complex multiplets between  $\delta_{\text{H}}$  1.96 – 2.14, three vinyl protons as multiplet at  $\delta_{\text{H}}$  5.09 and five vinyl methyl's at  $\delta_{\text{H}}$  1.59, 1.60, 1.60, 1.64, 1.67. The present of the geranyl-geranyl side chain was further confirmed by  $^{13}\text{C}$  NMR, HSQC and HMBC data. Confirmation of the structure was carried out by comparing the spectroscopic data ( $^1\text{H}$  and  $^{13}\text{C}$  NMR) of isolate with spectroscopic data of TPTQ [14].




Fig. 2:  $^{13}\text{C}$ -NMR and DEPT spectra of TPTQ in  $\text{CDCl}_3$




Fig. 3:  $^1\text{H}$ -NMR spectrum of TPTQ in  $\text{CDCl}_3$



Table 1: Cytotoxic activity of TPTQ towards cancer cell lines

Cell lines	IC <sub>50</sub>
MCF-7	>100
H-460	16.3 ± 3.0
DU-145	>100

MCF-7= Breast cancer cells, H-460= Lung cancer cell cells, DU-145= Prostate cancer cells

TPTQ showed good and selective activity towards H-460 cell lines and not active against MCF-7 and DU-145 (Table 1). Similar results were also observed on the crude extract [13].

#### ACKNOWLEDGMENT

The author thank the Malaysian Ministry of Science, Technology and the Environment are thanked for financial support under the Intensified Research in Priority Areas Research Grant (IRPA, EAR No. 09-02-04-0313), International Foundation for Science (IFS no F/3967-1), National L'Oreal Fellowship Grant.

#### CONFLICT OF INTEREST

We declare that we have no conflict of interest.

#### REFERENCES

- Reddy BS, Rao CV. Chemoprevention of colon cancer by thiol and other organosulfur compounds. Food phytochemicals for cancer prevention I: Fruits and vegetables. ACS Symposium series 546. Washington: American Chemical Society; 2003. p. 164-72.
- Rocha AB, Lopes RM, Schwartzman G. Natural products in anticancer therapy. *Curr Opin Pharmacol* 2001;1:364-9.
- Prasanth NV, Rasheed SP, Thomas T, Joseph S, Varghese CP. Evaluation of *in vitro* and *in vivo* anti-inflammatory activity of *Garcinia cambogia* L. *Int J Pharm Pharm Sci* 2013;5 Suppl 2:263-4.
- Shivakumar S, Sandhiya S, Subhasree N, Agrawal A, Dubey GP. *In vitro* assessment of antibacterial and antioxidant activities of fruit rind extracts of *Garcinia cambogia* L. *Int J Pharm Pharm Sci* 2013;5:254-7.
- Ghanekar KG, Dixit V. Antidiarrhoeal evaluation of aqueous extracts of *Garcinia cyminum* and a polyherbal formulation. *Int J Pharm Pharm Sci* 2014;6:329-31.
- Momo IJ, Kuetec V, Dufata H, Michela S, Wandji J. Antimicrobial of the methanolic extract and compounds from the stem bark of *Garcinia lucida* vesque (clusiaceae). *Int J Pharm Pharm Sci* 2011;3:215-7.
- Maikhuri RK, Gangwar AK. Ethnobotanical notes on the khasi and garo tribes of meghalaya northeast India. *Econ Bot* 1993;47:345-57.
- Lihitwitayawuid K, Phadungcharoen T, Mahidol C, Ruchirawat S. 7-O-Methylgarcinone E from *Garcinia cowa*. *Phytochem* 1997;45:1299-301.
- Rao RR. Ethnobotany of meghalaya: medicinal plants used by khasi and garo tribes. *Econ Bot* 1981;35:4-9.
- Ilham M, Yadav M, Norhanom AW. Tumor promoting activity of plants used in Malaysian traditional medicine. *Nat Prod Sci* 1995;1:31-42.
- Murakami A, Jiwajiinda S, Koshimiz K, Ohigashi H. Screening for *in vitro* antitumor promoting activities of edible plants from Thailand. *Cancer Lett* 1995;1:137-46.
- Likhitwitayawuid K, Phadungcharoen T, Krungkrai J. Antimalarial xanthenes from *Garcinia cowa*. *Planta Med* 1998;64:70-2.
- Jabit ML, Wahyuni FS, Khalid R, Shaari K, Lajis NH, Johnson Stanslas J. Cytotoxic and nitric oxide inhibitory activities of methanol extracts of *Garcinia* species. *Pharm Biol* 2009;47:1019-26.
- Wahyuni FS, Lindsay TB, Dachriyanus, Dianita R, Jubahar J, Lajis NH, *et al.* A new ring-reduced tetraprenyltoluquinone and a prenylated xanthone from *Garcinia cowa*. *Aust J Chem* 2004;57:223-6.
- Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 1983;16:55-63.

# International Journal of Pharmacy and Pharmaceutical Sciences Vol 7, Issue 3, 2015

---

## ORIGINALITY REPORT

---

6%

SIMILARITY INDEX

%

INTERNET SOURCES

6%

PUBLICATIONS

0%

STUDENT PAPERS

---

## MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

8%

★ Rohaya Ahmad, Abdul Manaf Ali, Daud A. Israf, Nor Hadiani Ismail, Khozirah Shaari, Nordin Hj. Lajis. "Antioxidant, radical-scavenging, anti-inflammatory, cytotoxic and antibacterial activities of methanolic extracts of some Hedyotis species", Life Sciences, 2005

Publication

---

Exclude quotes On

Exclude matches < 3%

Exclude bibliography On