

NK-1

by Nila Kasuma

Submission date: 16-Jun-2019 08:36AM (UTC+0700)

Submission ID: 1144011349

File name: NILA-1-Analysis_of_Breastfeeding_Pattern_with_Early.pdf (253.75K)

Word count: 3271

Character count: 16150

Analysis of Breastfeeding Pattern with Early Childhood Caries

¹Susi Susi, ²Murniwati Murniwati, ³Nila Kasuma, ⁴Minarni Minarni

ABSTRACT

Aim: This study aimed to investigate the relationship between breastfeeding pattern and early childhood caries (ECC) based on a case in Bukittinggi.

Materials and methods: This was a cross-sectional study located in Integrated Post Service (Posyandu) in Bukittinggi City. In this study, 86 pairs of parents and children aged 2 to 3 years were selected. Breastfeeding pattern was observed using questionnaire and decayed-extract-filled teeth (deft) index was used for ECC. Data were analyzed by chi-square test.

Results: The average of the deft index was 2.33. Samples with exclusive breastfeeding have an average deft index of 1.42 and nonexclusive had an average deft index of 2.48 ($p = 0.783$). This study also found that samples with exclusive breastfeeding and complementary foods had a deft index of 2.27. Nonexclusive breastfeeding and complementary foods had a deft index of 2.31 ($p = 0.038$).

Conclusion: Children with breastfeeding had a lower ECC degree. Breastfeeding with complementary foods at 6 months to 2 years had a significant relationship with ECC.

Clinical significance: Breastfeed could lower the occurrence of ECC. Hence, breastfeeding needs to be promoted and the mothers guided in breastfeeding.

Keywords: Breastfeeding, Decayed-extract-filled teeth, Early childhood caries.

How to cite this article: Susi S, Murniwati M, Kasuma N, Minarni M. Analysis of Breastfeeding Pattern with Early Childhood Caries. *World J Dent* 2018;9(3):197-200.

Source of support: This research was funded by the Ministry of Education, Research and Technology, Indonesia.

Conflict of interest: None

INTRODUCTION

Caries is an infectious disease, which has been ¹major problem in oral health that happens at all ages. ^{2,3}Early childhood caries is defined as caries in primary teeth among children under 71 months. The prevalence of ECC for children under 5 years is high in several countries. For instance, the prevalence of ECC for children under

5 years in the United States was 90%. ³The prevalence of ECC for children ³aged 2 to 5 years in Southern Taiwan was 68.9%. ⁴The prevalence of ECC for children aged 2 to 3 years in Jakarta was 52.75%. These facts show that the prevalences are still far from the World Health Organization (WHO) target that 90% children under 5 years old should be caries free. ⁵

Early childhood caries is a major health problem that needs serious attention, disrupting the function of mastication and affecting the growth and development of children, causing speech impairment and the children to become inferior. ^{6,7}The American Academic of Pediatric Dentistry states that drinking milk patterns, such as frequency, duration, addition of sweetener, and duration of breastfeed are risk factors for ECC. ⁸Carbohydrate content, acidity, and frequency of drinking milk affect the occurrence of caries. Carbohydrates contained in breast milk and formula milk are cariogenic foods that will be fermented by bacteria to acid and cause demineralization of enamel. ¹

Misinformation is one of the causes of ECC. ¹²Breast milk and bottle milk are risk factors for caries because they contain fermentable carbohydrate sucrose, which is a cariogenic food. ¹To address this problem, the WHO provides feeding guidelines for reducing caries prevalence. ⁵

In Indonesia, there is an increasing number of people of low economic status; to meet the needs of households, mothers help by going to work. Mothers with lack of education and health knowledge rarely breastfeed or only breastfeed for a short period of time. Breast milk is replaced with formula or sweet drinks like sweet tea and condensed milk. Mothers with low economic status have a tendency to substitute breast milk with bottled milk and high-sugar foods too early. This is due to the price of very expensive formula. ⁹The United Nations International Children's Fund reports the prevalence of mothers who breastfeed until the child is 12 months in developed countries is 74% while in developing countries it is 12%. ⁵

The American Academic of Pediatric Dentistry stated that breastfeeding pattern causes ECC. The aim of this research is to analyze breastfeeding pattern with ECC in Bukittinggi, Indonesia. To prevent the occurrence of ECC, the WHO makes guidelines for breastfeeding patterns. The WHO recommends that breastfeeding should be given exclusively until the child is 6 months old and

^{1,3}Faculty of Dentistry, Andalas University, Padang, Indonesia

⁴Department of Public Health, Health Polytechnic, Ministry of Health, Indonesia

Corresponding Author: Susi Susi, Faculty of Dentistry, Andalas University, Padang, Indonesia, Phone: +628126752834, e-mail: susiabin@gmail.com, susi@dent.unand.ac.id

followed by breastfeeding plus complementary food until the age of 2 years.⁵ This study aims to analyze the pattern of drinking milk with ECC in children aged 2 to 3 years.

MATERIALS AND METHODS

This research was a cross-sectional study in Posyandu in the city of Bukittinggi. Bukittinggi is a city with diverse socioeconomic status. Samples are taken from all districts in Bukittinggi. The number of samples was determined by the WHO manual.¹⁰ Research subjects were chosen by purposive sampling technique of 66 pairs of mothers and children ranging in age from 2 to 3 years with criteria of four upper anterior teeth already erupted. Research subjects were observed once about risk factors and effects according to current circumstances. The data were taken by six young dentists of the Faculty of Dentistry, Andalas University who were trained at first. The data were collected under the supervision of the head of the researcher.

Permission for the research was obtained from Research Ethics, Faculty of Medicine, Andalas University, Padang West Sumatera. Informed consent was given before retrieving the data. The pattern of breastfeeding in children aged 0 to 6 months and milk drinking patterns of children aged 6 months to 2 years were observed using questionnaires that have been tested for reliability and validity. Early childhood caries was examined using the WHO index deft criteria for epidemiological studies. Early childhood caries is categorized into three groups, i.e., mild (deft index 0–2.6), moderate (≥ 2.6 –4.4), and high (≥ 4.4). Data analysis was performed using Statistical Package for the Social Sciences, version 15. Significance of an effect was stated when $p < 0.05$ (significance level 95%). The prevalence of caries between two groups was tested using chi-square test.

RESULTS

A sample of 66 people was selected, of which 34 people had caries (51.5%), with an average deft index of 1.00 ± 2.76 [confidence interval (1.61–2.97)]. The sample consisted of 25 males (37.8%) and 41 females (62.0%) of age 2 to 3 years with an average age of 2.5 years. Children with exclusive breastfeeding (13, 19.6%) and children who received breast milk and complementary food from the age of 6 months to 2 years (37, 56%) (Table 1) were selected.

Males have higher average deft than females and children aged 3 years have a higher average deft than those aged 2 years. The prevalence of ECC was highest in children in the 3-year age group (Table 1). Children with exclusive breastfeeding have a lower average deft than who are not exclusively breastfed. Children with breastfeeding, bottle milk, and complementary foods have the highest average deft. The highest prevalence of ECC is in

Table 1: Prevalence of ECC (\times deft) based on characteristics

Variable	n	Prevalence of ECC	\times deft	SD
Sex				
Male	25 (37.8%)	13 (52%)	2.56	3.203
Female	41 (62.2%)	20 (48.8%)	2.21	2.48
Age (months)				
24	33 (50%)	13 (39.4%)	1.45	2.166
36	33 (50%)	22 (66.7%)	3.12	3.059

SD: Standard deviation

Table 2: Prevalence of ECC (\times deft) based on breastfeeding

Variable	n	Prevalence of ECC	\times deft	SD
Exclusive breastfeeding	12 (18.2%)	5 (41.7%)	1.42	2.065
Nonexclusive breastfeeding	54 (81.8%)	29 (53.7%)	2.48	2.875
Breastfeeding with complementary foods	37 (56.1%)	23 (62.1%)	2.27	2.446
Nonbreastfeeding with complementary foods	29 (43.9%)	9 (31%)	2.48	2.875

SD: Standard deviation

Table 3: Prevalence of ECC (\times deft) based on feeding pattern from 6 to 24 months

Variable	n	Prevalence of ECC	\times deft	SD
Frequency $\leq 7 \times$	53 (80.3%)	21 (39.6%)	2.17	2.827
Frequency $> 7 \times$	13 (19.7%)	13 (100%)	2.77	2.522
Duration ≤ 15 minutes	42 (63.6%)	19 (45.2%)	2.02	2.580
Duration > 15 minutes	24 (36.4%)	14 (58.3%)	2.75	3.054
Not as lullaby	2 (3.3%)	1 (50.0%)	4.00	3.567
As lullaby	64 (96.7%)	33 (51.6%)	2.23	2.695
No formula feed	36 (54.5%)	19 (52.7%)	2.19	2.505
Formula feed	30 (45.4%)	17 (56.7%)	2.40	3.081
Stop breastfeeding < 24 months	44 (66.7%)	23 (52.3%)	2.27	2.732
Stop breastfeeding > 24 months	22 (33.3%)	11 (50.0%)	2.32	2.901
Stop drinking milk ≤ 14 months	33 (50%)	19 (57.6%)	2.18	2.551
Stop drinking milk > 14 months	33 (50%)	15 (58.1%)	2.09	2.989

SD: Standard deviation

children with breastfeeding and complementary feeding groups (Table 2).

Based on breastfeeding pattern, the prevalence of ECC is highest in children who stop drinking bottled milk when they are older than 14 months. The highest deft index was found in children who were not lactating (Table 3).

The ECC distribution by WHO category is 59.1% low, 16.7 medium, and 24.2 high. Exclusive breastfeeding group was with the category of ECC 66.7% low, 16.7%

Table 4: Breastfeeding pattern and ECC

Variable	Mild	Moderate	High	p-value
Exclusive breastfeeding	8 (66.6%)	2 (16.7%)	2 (16.7%)	0.783
Nonexclusive breastfeeding	21 (38.8%)	9 (16.7%)	14 (25.9%)	
Breastfeeding with complementary foods	19 (51.4%)	10 (27.0%)	8 (21.6%)	0.038*
Nonbreastfeeding with complementary foods	20 (69.0%)	1 (3.4%)	8 (27.6%)	
Frequency <7 times/day	33 (62.3%)	8 (15.1%)	12 (22.6%)	0.565
Frequency ≥7 times/day	6 (46.2%)	2 (23.1%)	4 (30.8%)	
Not as lullaby	27 (64.3%)	5 (11.9%)	10 (23.8%)	0.349
As lullaby	12 (50.0%)	6 (25.0%)	6 (25.0%)	
No formula feed	19 (52.8%)	9 (25.0%)	8 (22.2%)	0.138
Formula feed	20 (66.7%)	2 (6.7%)	8 (26.7%)	
Stop breastfeeding <24 months	26 (59.1%)	8 (18.2%)	10 (22.7%)	0.138
Stop breastfeeding >24 months	13 (59.1%)	3 (13.8%)	6 (27.3%)	
Stop drinking milk ≤14 months	15 (45.5%)	10 (30.3%)	8 (24.2%)	0.009*
Stop drinking milk >14 months	24 (72.7%)	1 (3.0%)	8 (24.2%)	

*Significance difference, $p < 0.05$

medium, and 16.7% high. Exclusive breastfeeding group with ECC category is 57.4% low, medium 16.7%, and 25.9% high. The ECC with high category was found more in the nonexclusive breastfeeding group. There was a significant relationship between breastfed children plus complementary foods from 6 months to 2 years of age compared with those with breast milk, bottles, and complementary foods ($p < 0.05$) (Table 4).

DISCUSSION

The exclusive breastfeeding coverage in this study (19.6%) was lower than the national exclusive breastfeeding coverage (52.3%) in 2004 while the exclusive breastfeeding coverage of West Sumatra Province was 73.6%. World Health Organization in 2011 recommended exclusive breastfeeding until the age of 6 months and then breastfeeding should be accompanied by complementary food until the age of 2 years. Exclusive breastfeeding according to Government Regulation No. 33 of 2012 is to provide breast milk to infants from birth to 6 months of age, without adding or replacing with food or drink except essential drugs, vitamins, and minerals. Breast milk is the ideal nutrition for babies and provides immunological protection. Intestines not only provide nutritional intake but also psychosocial care through the bonding of affection with mother and health based on immunological substances contained in breast milk.^{11,12}

The results showed that 56% children got breast milk until the age of 2 years. The results of this study were lower than the previous study conducted by Ragama, in which they got 71.6% as result.¹¹ The survey results of Hellen Keller in Jakarta showed that only 1% of children who got breast milk until age 6 months.¹⁴ At this time, many women are working to help the family's economy. This causes a limited breastfeeding to the baby. According to the National Health Survey 2012, 47.9% of women who

work in Indonesia in Jakarta, of 44.8% of women who work, 28.7% had children under the age of 5 years.^{12,14}

The prevalence of ECC in this study was 51.5% with an average deft of 1.00 ± 2.76 , similar to that of Sugito et al⁹ in DKI Jakarta in children under 3 years of age. The prevalence was 52.7% with an average deft of 2.85, and research results at Mount Anyar¹⁵ showed an ECC prevalence of 30.8% for children aged 6 months to 3 years. This is still far from the WHO target of 90% of children under 5 years of age who are caries free. Primary teeth are more susceptible to caries because the enamel structure is less dense and thinner than permanent teeth.² Boys have an average deft higher than girls and children aged 3 years have an average deft higher than the age of 2 years but statistical test results have no significant relationship.

Children with exclusive breastfeeding had a lower deft average than no breastfeeding, but statistical test results showed no significant association, this result was similar to that of Perera et al.¹⁵ Breastfed children and with complementary foods from 6 months to 2 years had a lower deft average and from statistical tests, there was a significant association. Calcium and phosphate in breast milk will protect teeth from caries by stimulating remineralization in enamel.¹

The frequency of drinking milk of less than seven times in a day and the duration of drinking milk less than 15 minutes has a lower deft average, but from the statistical test results, there is no significant relationship. The results of Lida et al¹⁶ in the United States are also the same.

Children who were breastfed or had infant formula at night had a higher deft average than those who did not, but statistical tests showed no significant association. The results of Perera et al¹⁵ and Bahuguna et al¹⁷ showed that daytime milk feeding is a risk factor for ECC. When sleeping, saliva production is very little and the remaining milk in the mouth will be fermented by bacteria,

resulting in demineralization of enamel especially in the upper anterior teeth. Breastfeeding at night will be stopped when the first tooth erupts.¹

Children who drink bottle milk and sweetened milk had a higher deft index but no statistically significant test results. Bahuguna et al's¹⁷ results suggest that bottle-feeding and sweetener addition are risk factors for ECC. In the process of making formula milk sweeter like breast milk, sucrose is added. Sucrose and other sweeteners are cariogenic, causing dental demineralization.^{18,19}

Children who stop breastfeeding at age 2 years and stop drinking bottle milk before the age of 14 months have a lower deft average, but statistical test results have no significant relationship. The results of Tyagi and Bahuguna showed that there was a relationship between duration of drinking of breast milk and ECC. Breastfeeding period will affect the child's diet.^{18,20}

The main limitation of this study is the oral glass that was used in the study was unacceptable to some children. The study also used headlamp in assessing caries presence.

CONCLUSION

Children who get exclusive breastfeeding and breast milk plus complementary foods aged 6 months to 2 years have a lower deft average. Exclusive breastfeeding has no significant association with ECC but breastfeeding plus complementary feeding from 6 months to 2 years of age has a significant relationship with ECC.

ACKNOWLEDGMENTS

Authors would like to thank all parents and children who have participated in this research and also the head of health care center in Bukittinggi and Posyandu team who coordinated in this research in data retrieval.

REFERENCES

- Harris R, Nicoll AD, Adair PM, Pine CM. Risk factors for dental caries in young children: a systemic review of the literature. *Community Dent Health* 2004 Mar;21(Suppl 1):71-85.
- McDonald RE, Avery DR, Dean JA. *Dentistry for the child and adolescent*. 8th ed. Mosby; 2004.
- Tinanoff N, O'Sullivan DM. Early childhood caries: overview and recent findings. *Pediatr Dent* 1997 Jan-Feb;19(1):12-16.
- Hsiao SY, Huang ST, Chao LH, Liu HY, Chang CP. The dental caries status of 0-6-year-old children in Southern Taiwan. *Chin Dent J* 2007;26:52-60.
- World Health Organization. *Global strategy for infant young child feeding*. UNICEF; 2003.
- Ramos-Gomez FJ, Weinstraub JA, Gansky SA, Hoover CI, Featherstone JD. Bacterial, behavioral and environmental factors associated with early childhood caries. *J Clin Pediatr Dent* 2002 Winter;26(2):165-173.
- Fejerkof O, Kidd E. *Dental caries: the disease and its clinical management*. 2nd ed. Blackwell Munksgaard; 2008.
- Tang RS, Huang ST, Chen HS, Hsiao SY, Hu HY, Chuang FH. The association between oral hygiene behavior and knowledge of caregivers of children with severe early childhood caries. *J Dent Sci* 2014;9:277-282.
- Sugito PS, Djoharnas H, Darita RW. Breast feeding and early childhood caries (ECC) severity of children under three years old in DKI Jakarta. *Makara J Dent Res* 2008;12(22).
- Lwanga SK, Lemeshow S. *Sample size in health studies*. Geneva: World Health Organization; 1991.
- Kramer MS, Kakuma R. The optimal duration of exclusive breastfeeding: a systematic review. Geneva: Department of Nutrition for Health and Development, Department of Child and Adolescent Health and Development, WHO; 2002.
- Ministry of Health Indonesia. *Health Profile*, 2014.
- Perera PJ, Fernando MP, Warrakulasooriya TD, Ranathunga N. Effect of feeding practice on dental caries among preschool children: a hospital-based analytical cross-sectional study. *Asia Pacific J Clin Nutr* 2014;23(2):272-277.
- Weinstein P. Public health issues in early childhood caries. *Community Dent Oral Epidemiol* 1998;26(Suppl 1):84-90.
- Sutjipto RW, Kuntari HS. The prevalences of early childhood caries and severe early childhood caries in preschool children at Gunung Anyar Surabaya. *Dent J* 2014;37(4):186-189.
- Lida H, Aunger P, Billing RJ, Weitzman M. Association between infant breastfeeding and early childhood caries in the United States. *Pediatrics* 2007 Oct;120(4):e944-e952.
- Bahuguna R, Younis Khan S, Jain A. Influence of feeding practices on dental caries: a case-control study. *Eur J Pediatr Dent* 2013 Mar;14(1):55-58.
- Cameron AC, Widmer RP. *Handbook of pediatric dentistry*. 2nd ed. Mosby; 2003.
- World Health Organization. *Guideline: sugar intake for adults and children*. Geneva; 2015.
- Tyagi R. The prevalence of nursing caries in Davangere preschool children and its relationship with feeding practices socioeconomic status of the family. *J Indian Soc Pedod Prev Dent* 2008 Dec;26(4):153-157.

ORIGINALITY REPORT

5%

SIMILARITY INDEX

2%

INTERNET SOURCES

2%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to University of Oklahoma Health Science Center 1%
Student Paper
- 2** Submitted to Laureate Higher Education Group <1%
Student Paper
- 3** "Early childhood caries : Definition and epidemiology", Early Childhood Oral Health, 2015. <1%
Publication
- 4** apjcn.nhri.org.tw <1%
Internet Source
- 5** alexandriadentaljournal.com <1%
Internet Source
- 6** www.ncbi.nlm.nih.gov <1%
Internet Source
- 7** Submitted to Riga Stradins University <1%
Student Paper
- 8** Submitted to Higher Education Commission Pakistan <1%

9 globalresearchnurses.tghn.org <1%
Internet Source

10 whqlibdoc.who.int <1%
Internet Source

11 onlinelibrary.wiley.com <1%
Internet Source

12 Submitted to Universidad Nacional Mayor de San Marcos <1%
Student Paper

Exclude quotes On

Exclude matches Off

Exclude bibliography On