

Limited effectiveness of household sand filters for removal of arsenic from well water in North Vietnam

by Dr. Cimi Ilmiawati Dr. Cimi Ilmiawati

Submission date: 29-Aug-2018 04:06PM (UTC+0800)

Submission ID: 994474334

File name: Limited_Effectiveness_1.pdf (397.94K)

Word count: 5126

Character count: 25628

Limited effectiveness of household sand filters for removal of arsenic from well water in North Vietnam

Cimi Ilmiawati, Nguyen Dinh Thang, Machiko Iida, Masao Maeda, Shoko Ohnuma, Ichiro Yajima, Nobutaka Ohgami, Reina Oshino, M. M. Aeorangajeb Al Hossain, Hiromasa Ninomiya and Masashi Kato

ABSTRACT

Since well water utilized for domestic purposes in the Red River Delta of North Vietnam has been reported to be polluted by arsenic, barium, iron, and manganese, household sand filters consisting of various components are used. Information regarding the effectiveness of various sand filters for removal of the four toxic elements in well water is limited. In this study, arsenic levels in 13/20 of well water samples and 1/7 of tap water samples exceeded WHO health-based guideline value for drinking water. Moreover, 2/20, 6/20, and 4/20 of well water samples had levels exceeding the present and previous guideline levels for barium, iron, and manganese, respectively. Levels of iron and manganese, but not arsenic, in well water treated by sand filters were lower than those in untreated water, although previous studies showed that sand filters removed all of those elements from water. A low ratio of iron/arsenic in well water may not be sufficient for efficient removal of arsenic from household sand filters. The levels of barium in well water treated by sand filters, especially a filter composed of sand and charcoal, were significantly lower than those in untreated water. Thus, we demonstrated characteristics of sand filters in North Vietnam.

Key words | arsenic, barium, iron, manganese, sand filter, well water

Cimi Ilmiawati
Machiko Iida
Masao Maeda
Ichiro Yajima
Nobutaka Ohgami
Reina Oshino
M. M. Aeorangajeb Al Hossain
Hiromasa Ninomiya
Masashi Kato (Corresponding author)
Departments of Occupational and Environmental Health,
Nagoya University Graduate School of Medicine,
Nagoya,
Aichi 466-8550,
Japan
E-mail: katomasa@med.nagoya-u.ac.jp

Nguyen Dinh Thang
Machiko Iida
Ichiro Yajima
Nobutaka Ohgami
Masashi Kato
Units of Environmental Health Sciences,
Department of Biomedical Sciences, College of
Life and Health Sciences,
Chubu University,
Kasugai,
Aichi 487-8501,
Japan

Cimi Ilmiawati
Machiko Iida
Masao Maeda
Shoko Ohnuma
Ichiro Yajima
Nobutaka Ohgami
Reina Oshino
M. M. Aeorangajeb Al Hossain
Hiromasa Ninomiya
Masashi Kato
Voluntary Body for International Healthcare in
Universities,
Nagoya,
Aichi 466-8550,
Japan

Cimi Ilmiawati
Department of Pharmacology,
Faculty of Medicine Andalas University,
Padang,
West Sumatra,
Indonesia

Nguyen Dinh Thang
University of Science, Vietnam National University,
Hanoi,
Vietnam

INTRODUCTION

Well water derived from groundwater is important as a domestic water source, including drinking water in Asian countries. However, well water may have a problem of naturally occurring arsenic (As) due to As-enriched aquifers. In fact, arsenic pollution of drinking water from wells has been found in many countries, including Bangladesh, India, Cambodia, Myanmar, and Vietnam (Argos *et al.* 2012; Kumasaka *et al.* 2013). Previous studies showed that chronic exposure to As via drinking water can cause various diseases, including cancers, cutaneous pigmented disorders, and diabetes (Nizam *et al.* 2013; Yajima *et al.* 2015).

The World Health Organization (WHO) has provided guideline values for the quality of drinking water for barium (Ba), iron (Fe), and manganese (Mn) in addition to As in the present and/or previous editions (WHO 1984, 2008, 2011). Barium naturally coexists with other toxic elements in arsenic-polluted well water in Bangladesh and Vietnam (Frisbie *et al.* 2009; Kato *et al.* 2013). Our previous fieldwork studies showed a correlation between As and Ba in well drinking water and in human samples (urine, nails, and hair) in Bangladesh (Kato *et al.* 2013). Our previous experimental studies *in vitro* showed that Ba promotes a malignant characteristic of human non-tumorigenic keratinocytes (Thang *et al.* 2015) and that Ba suppresses As-mediated anti-cancer effects (Yajima *et al.* 2012). Moreover, Ba was shown to promote hearing loss in mice (Ohgami *et al.* 2012) and humans (Ohgami *et al.* 2016) in our experimental study and epidemiological study, respectively. Previous studies showed that Fe promotes all steps of carcinogenesis (Weinberg 1996). In fact, previous studies have shown effects of iron on tumor initiation and tumor growth (Kumasaka *et al.* 2013; Torti & Torti 2013). A previous epidemiological study also showed that an elevated level of serum iron increases the risk of several carcinomas in humans (Wen *et al.* 2014). Exposure to Mn in drinking water has been reported to be associated with neurobehavioral disorders in mice (Krishna *et al.* 2014; Kumasaka *et al.* 2014). Exposure to Mn from well drinking water is also associated with an increase in infant mortality, cognitive deficit, memory deficit, and lower intelligence scores in humans (O'Neal & Zheng 2015). Thus, high uptake is a

problem, which necessitates WHO to provide guidelines for the quality of drinking water, although some of the four elements are needed at trace levels. In addition, high levels of As, Ba, Fe, and Mn are harmful to aquatic life, biota, and the environment as well as human health (Wang 1988; Korte & Fernando 1991; Ventura-Lima *et al.* 2011; Kalantzi *et al.* 2013).

Approximately 13 million people in Vietnam (16.5% of the population), who mostly reside in the Red River Delta, are drinking water from wells (Nguyen *et al.* 2009; Winkel *et al.* 2011). There have been many reports on pollution of As and other toxic elements in well drinking water in North Vietnam (Agusa *et al.* 2006, 2014; Berg *et al.* 2006; Nitzsche *et al.* 2015). Levels of As in well water are more than 1,000 µg/L (Berg *et al.* 2001), more than 100-fold higher than the value in WHO health-based guidelines for drinking water. Previous studies also showed increased levels of Fe and Mn in well water in North Vietnam (Agusa *et al.* 2006; Berg *et al.* 2006; Nitzsche *et al.* 2015). However, there is little information on Ba levels in well water in North Vietnam compared to the information on As, Fe, and Mn levels.

Another source of drinking water in Vietnam is harvested rainwater. The quality of harvested rainwater in the Mekong Delta was investigated in a previous study (Wilbers *et al.* 2013). The quality of well water and that of rainwater were compared in another study in Hanoi (Agusa *et al.* 2006). However, there has been no report comparing the quality of well water, rainwater, and tap water in North Vietnam.

Household sand filters are commonly used to remediate well water in North Vietnam in order to obtain drinkable water by removing Fe, which causes discoloration and a metallic taste (Berg *et al.* 2006). Previous studies showed that sand filters are effective for removing As, Fe, and Mn from well water (Berg *et al.* 2006; Nitzsche *et al.* 2015). However, information on the effectiveness of sand filters for removal of Ba in well water is limited despite the fact that previous studies showed high levels of Ba in well water in North Vietnam (Agusa *et al.* 2006; Winkel *et al.* 2011). To our knowledge, there is also limited information on the

effectiveness of sand filters consisting of different components for removal of toxic elements in well water.

In order to assess the health risk of domestic water in North Vietnam, levels of four toxic elements (As, Ba, Fe, and Mn) were compared among well water, rainwater, and tap water in this study. The effectiveness of household sand filters consisting of different components for removal of toxic elements was also investigated.

METHODS

Water sampling

Water samples were collected at Vinh Tru Town (six hamlets), Nhan Khang commune (three hamlets), and Dong Ly commune (one hamlet) in Ly Nhan District of Ha Nam Province, where pollution of As in well water was previously reported (Nguyen *et al.* 2009). A total of 20 untreated and 77 sand filter-treated well water samples were collected. In addition, 105 harvested rainwater and seven tap water samples were collected. All samples were collected in polyethylene bottles. Each bottle was filled with sampled water after rinsing out the inside of the bottle with sampled water. Tightly capped bottles were promptly sent from Vietnam to Japan by airplane. Samples were kept at 4 °C and measurements of total levels of As, Ba, Fe, and Mn were completed within 2 weeks after arrival at Nagoya University. Free and informed consent of the participants or their legal representatives was obtained, and the study protocol was approved by the Ethical Committee of Chubu University, Aichi, Japan (approval No. 20008 on July 9, 2008; approval No. 260019 on July 8, 2014) and the Ethical Committee of Nagoya University, Aichi, Japan (approval No. 2013-0070 on July 22, 2013) by following the regulations of the Japanese government.

Analytical method

Levels of As, Ba, Fe, and Mn in water samples were quantified by using an inductively coupled plasma-mass spectrometer (ICP-MS; 7500cx, Agilent Technologies Inc., CA, USA) according to the method previously described

(Ohgami *et al.* 2012; Kato *et al.* 2013). The limits of detection for As, Ba, Fe, and Mn by the ICP-MS are 0.1, 0.1, 3.0, and 0.1 µg/L, respectively.

Statistical analyses

The Mann–Whitney *U* test and chi-square test were used for statistical analyses to compare differences between groups. The JMP Pro software package (version 11.0.0; SAS Institute, Cary, NC, USA) was used for all statistical analyses.

RESULTS

Levels of the four elements in various sources of water in the Red River Delta of North Vietnam

Median levels of As, Ba, Fe, and Mn in well water samples ($n = 20$), rainwater samples ($n = 105$), and tap water samples ($n = 7$) collected in Ha Nam Province, Vietnam are shown in Figure 1. Levels (mean \pm SD) of As, Ba, Fe, and Mn in well water samples ($n = 20$) collected in Ha Nam Province, Vietnam are shown in Table 1. Boxplots (Figure 1) show that the levels of As, Ba, Fe, and Mn in rainwater and tap water were lower than those in well water. However, there was one tap water sample containing 27.8 µg/L of As (Figure 1(a)).

Since the levels of the four elements in well water were higher than those in rainwater and tap water, pollution of the toxic elements in well water was then focused on in this study. The median level (14.6 µg/L) of As in well water exceeded the value in the current WHO guidelines (10 µg/L) for drinking water, while the median level of Ba in well water was below the value in the current guidelines (700 µg/L). Although there are no guideline values for Fe and Mn in the present edition, the median levels of Fe and Mn in well water were below the guideline values in the previous editions (300 µg/L for Fe and 400 µg/L for Mn). The percentages of unsafe wells with levels of As, Ba, Fe, and Mn exceeding the values of WHO health-based guidelines for drinking water in the present and/or previous editions were 65%, 10%, 30%, and 20%, respectively (Table 1).




Figure 1 | Boxplots of levels of four elements in well water, rainwater, and tap water in North Vietnam. Concentrations of As, Ba, Fe, and Mn in well water, harvested rainwater, and tap water in Ha Nam Province, Vietnam are presented. Boxplots are used to present the first quartile, median, third quartile (each presented by a horizontal line of the box) and minimum (lower whisker) and maximum (upper whisker) values. Broken lines show previous (Fe and Mn) and present (As and Ba) guideline values for drinking water by WHO. * and **, significantly different ($p < 0.05$; $**p < 0.01$) from well water by the Mann-Whitney *U* test.

Table 1 | Levels of four elements in untreated well water and well water treated by sand filters

Element	WHO guideline value (µg/L)	Element concentration (µg/L)		Filter	Unsafe wells (%)
		Mean	SD		
As	10 [#]	15.7	20	No	65 ns
		23.7	28.0	Yes	74 ns
Ba	700 [#]	422.4	240.4	No	10 ns
		322.9	168.6	Yes	2.6ns
Fe	300 ^{###}	186.3	267.2	No	30 ^{**}
		10.2	22.3	Yes	0 ^{**}
Mn	400 ^{**}	236.0	287.7	No	20 ^{**}
		16.2	46.4	Yes	0 ^{**}

Levels (mean ± SD) of As, Ba, Fe, and Mn in untreated well water and well water treated by sand filters in North Vietnam are presented. Unsafe wells (%) are percentages of wells with levels of As, Ba, Fe, and Mn exceeding the values in WHO health-based guidelines in untreated well water and well water treated by sand filters.

[#], ^{**}, and ^{###} WHO guideline values in the previous (^{###}, 1984; ^{**}, 2008) and present ([#], 2011) editions.

^{**}Statistically different ($p < 0.01$) from the untreated control by the chi-square test. ns, not significant.

Levels of the four elements in untreated well water and well water treated by sand filters

In our fieldwork study, 79.4% (77/97) of randomly selected wells water were treated by sand filters (Figure 2), suggesting that the use of sand filters is common in Ha Nam Province. Boxplots of the levels of As, Ba, Fe, and Mn in untreated well water samples ($n = 20$) and well water samples treated by sand filters ($n = 77$) (Figure 2) show that the levels of As in untreated well water and well water treated by sand filters are comparable (Figure 2(a)). Correspondingly, the percentages of unsafe wells with As-polluted untreated water and water treated by sand filters were comparable (Table 1). These results suggest very limited effectiveness of sand filters for removal of As from well water in North Vietnam. Levels of Ba, Fe, and Mn in well water treated by sand filters were significantly lower than those in untreated well water (Figure 2(b)–2(d)), although the difference in Ba levels is




Figure 2 | Boxplots of levels of four elements in untreated well water and well water treated by sand filters. Concentrations of As, Ba, Fe, and Mn in untreated well water (control) and well water treated by sand filters in Ha Nam Province, Vietnam are presented. Boxplots are used to present the first quartile, median, third quartile (each presented by a horizontal line of the box) and minimum (lower whisker) and maximum (upper whisker) values. Broken lines show previous (Fe and Mn) and present (As and Ba) guideline values for drinking water by WHO. * and **, significantly different ($p < 0.05$; $**p < 0.01$) from the untreated control by the Mann-Whitney *U* test.

smaller than the differences in Fe and Mn levels in untreated well water and well water treated by sand filters. Furthermore, the percentages of wells with Fe-polluted and Mn-polluted but not Ba-polluted well water treated by sand filters were significantly lower than those with water not treated by sand filters (Table 1). These results suggest

that the effectiveness of sand filters for removal of Fe and Mn is greater than their effectiveness for removal of Ba in well water in North Vietnam.

Levels of As, Ba, Fe, and Mn in untreated well water and in well water treated by three kinds of sand filters consisting of sand only, sand plus charcoal, and sand plus gravel (Figure 3)




Figure 3 | Cross-sectional graphs of household sand filters used in Ha Nam, North Vietnam. Schemas for three kinds of sand filters composed of sand (a), sand and charcoal (b), and sand and gravel (c) are presented.

were compared. The levels of As in untreated well water and in well water treated by the three kinds of sand filters were comparable (Figure 4(a)), while levels of Fe and Mn in well water treated by the three kinds of sand filters were lower than those in untreated well water (Figure 4(c) and 4(d)). These results suggest that all kinds of sand filters can remove Fe and Mn but not As. Moreover, levels of Ba in well water treated by a sand filter consisting of sand and charcoal were lower than the levels in untreated well water (Figure 4(b)). Levels of Ba in well water treated by a sand filter consisting of sand only and in well water treated by a sand filter consisting of sand plus gravel were comparable to the levels in untreated well water. These results suggest that a sand

filter consisting of sand and charcoal may effectively remove Ba from well water.

DISCUSSION

We first compared levels of As, Ba, Fe, and Mn in well water, rainwater, and tap water in North Vietnam. Since the level of As in 65% of the well water samples exceeded the value in WHO health-based guidelines for drinking water, well water is not suitable for drinking water, as was shown in previous studies (Berg et al. 2007; Agusa et al. 2014). Rainwater and tap water seem to be suitable for drinking from the viewpoint of the four toxic elements.




Figure 4 | Boxplots of levels of four elements in untreated well water and well water treated by three kinds of sand filters. Concentrations of As, Ba, Fe, and Mn in untreated well water and well water treated by various sand filters composed of sand only, sand plus charcoal, and sand plus gravel in Ha Nam Province, Vietnam. Boxplots are used to present the first quartile, median, third quartile (each presented by a horizontal line of the box) and minimum (lower whisker) and maximum (upper whisker) values. * and **, significantly different (* $p < 0.05$; ** $p < 0.01$) from the untreated control by the Mann-Whitney U test.

Unexpectedly, however, As at a level of 27.8 µg/L was detected in one of seven tap water samples. Our results suggest that continuous monitoring for various kinds of water including tap water is important in the future in North Vietnam.

A previous study showed that the level of As in well water from Ly Nhan District, North Vietnam was 420 µg/L and that the level in sand-filtered water was 23 µg/L, with 80% of the filtered water still containing As at a level higher than 10 µg/L (Agusa *et al.* 2014). Other studies in Vietnam also showed that sand filters could remove 80% (Berg *et al.* 2006) to 95% of As (Nitzsche *et al.* 2015). Sand filters have been reported to be effective for the removal of Mn and Fe with 86.2–99.6% efficiency and nearly 100% efficiency, respectively (Nitzsche *et al.* 2015). Correspondingly, our results showed that levels of Fe and Mn in well water treated by sand filters were lower than those in untreated well water, suggesting that sand filters are effective for removal of Fe and Mn in well water. In contrast to previous studies (Agusa *et al.* 2014; Nitzsche *et al.* 2015), however, levels of As in untreated well water and in well water treated by sand filters were comparable in our study. A previous study showed that Fe/As ratios of ≥ 50 and >250 are required to reduce As concentrations to levels below 50 µg/L and 10 µg/L, respectively, indicating that the level of Fe in water affects removal of As by sand filters (Berg *et al.* 2006). The median of Fe/As ratio in well water was about 5 in this study (Figure S1) and this might be a reason why sand filters could not remove As from well water. Our results suggest that the Fe/As ratio in well water should be examined to determine the effectiveness of sand filters for removal of As. Since a sand filter composed of sand and zero-valent iron filings was shown to be effective for arsenic removal (Leupin & Hug 2005; Leupin *et al.* 2005; Mehta & Chaudhari 2015), the use of such a sand filter may be one way to overcome the low performance of a sand filter based on the Fe/As ratio. Proper design of a sand filter may also improve the effectiveness for As removal (Leupin *et al.* 2005). Our results showing a decrease of 24% in Ba in well water treated with sand filters suggest that sand filters can partially remove Ba from well water. To our knowledge, however, there has been no study showing the effectiveness of charcoal and a charcoal-containing sand filter for Ba removal.

Further study on their effectiveness for Ba removal is needed.

CONCLUSIONS

After confirming pollution of As, Ba, Fe, and Mn in well water in North Vietnam, we found that sand filters could remove Fe and Mn but not As, despite the fact that previous studies showed that sand filters could remove As (Berg *et al.* 2006; Nitzsche *et al.* 2015). We then showed that a low ratio of Fe/As in well water in North Vietnam might be one of the reasons for the poor performance of sand filters. Thus, this fieldwork study clarified characteristics of various sand filters being used in North Vietnam.

ACKNOWLEDGEMENTS

This study was supported in part by Grants-in-Aid for Scientific Research (A) (15H01743 and 15H02538), (B) (24390157 and 24406002), and (C) (16K11177), Grant-in-Aid for Challenging Exploratory Research (26670525), Grant-in-Aid for Scientific Research on Innovative Areas (24108001) and Grant-in-Aid for Research Activity Start-up (15H06274) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), the Mitsui & Co., Ltd. Environment Fund, The Mitsubishi Foundation, Ichihara International Scholarship Foundation, Health Sciences Foundation (Kenko-Kagaku Zaidan) and International Fellowship Programs For Foreign Researchers from Takeda Science Foundation. The funding sources had no involvement in the study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. All authors declare that there is no conflict of interest.

REFERENCES

- Agusa, T., Kunito, T., Fujihara, J., Kubota, R., Minh, T. B., Trang, P. T. K., Iwata, H., Subramanian, A., Viet, P. H. & Tanabe, S. 2006 Contamination by arsenic and other trace elements in

- tube-well water and its risk assessment to humans in Hanoi, Vietnam. *Environmental Pollution* **139**, 95–106.
- Agusa, T., Trang, P. T. K., Lan, V. M., Anh, D. H., Tanabe, S., Viet, P. H. & Berg, M. 2014 Human exposure to arsenic from drinking water in Vietnam. *Science of the Total Environment* **488–489**, 562–569.
- Argos, M., Ahsan, H. & Graziano, J. H. 2012 Arsenic and human health: epidemiologic progress and public health implications. *Reviews on Environmental Health* **27** (4), 191–195.
- Berg, M., Tran, H. C., Nguyen, T. C., Pham, H. V., Schertenleib, R. & Giger, W. 2001 Arsenic contamination of groundwater and drinking water in Vietnam: a human health threat. *Environmental Science & Technology* **35** (13), 2621–2626.
- Berg, M., Luzi, S., Trang, P. T. K., Viet, P. H., Giger, W. & Stuben, D. 2006 Arsenic removal from groundwater by household sand filters: comparative field study, model calculations, and health benefits. *Environmental Science and Technology* **40** (17), 5567–5573.
- Berg, M., Stengel, C., Trang, P. T. K., Viet, P. H., Sampson, M. L., Leng, M., Samreth, S. & Fredericks, D. 2007 Magnitude of arsenic pollution in the Mekong and Red River Deltas—Cambodia and Vietnam. *Science of the Total Environment* **372**, 413–425.
- Frisbie, S. H., Mitchell, E. J., Mastera, L. J., Maynard, D. M., Yusuf, A. Z., Siddiq, M. Y., Ortega, R., Dunn, R. K., Westerman, D. S., Bacquart, T. & Sarkar, B. 2009 Public health strategies for Western Bangladesh that address arsenic, manganese, uranium, and other toxic elements in drinking water. *Environmental Health Perspective* **117** (3), 410–416.
- Kalantzi, I., Shimmield, T. M., Pergantis, S. A., Papageorgiou, N., Black, K. D. & Karakassis, I. 2013 Heavy metals, trace elements and sediment geochemistry at four Mediterranean fish farms. *Science of the Total Environment* **444**, 128–137.
- Kato, M., Kumasaka, M. Y., Ohnuma, S., Furuta, A., Kato, Y., Shekhar, H. U., Kojima, M., Koike, Y., Dinh Thang, N., Ohgami, N., Ly, T. B., Jia, X., Yetti, H., Naito, H., Ichihara, G. & Yajima, I. 2013 Comparison of barium and arsenic concentrations in well drinking water and in human body samples and a novel remediation system for these elements in well drinking water. *PLoS One* **8** (6), e66681.
- Korte, N. E. & Fernando, Q. 1991 A review of arsenic (III) in groundwater. *Critical Reviews in Environmental Control* **21** (1), 1–39.
- Krishna, S., Dodd, C. A., Hekmatyar, S. K. & Filipov, N. M. 2014 Brain deposition and neurotoxicity of manganese in adult mice exposed via the drinking water. *Archives of Toxicology* **88** (1), 47–64.
- Kumasaka, M. Y., Yamanoshita, O., Shimizu, S., Ohnuma, S., Furuta, A., Yajima, I., Nizam, S., Khalequzzaman, M., Shekhar, H. U., Nakajima, T. & Kato, M. 2013 Enhanced carcinogenicity by coexposure to arsenic and iron and a novel remediation system for the elements in well drinking water. *Archives of Toxicology* **87** (3), 439–447.
- Kumasaka, M. Y., Yajima, I., Ohgami, N., Naito, H., Omata, Y. & Kato, M. 2014 Commentary to Krishna et al. 2014: Brain deposition and neurotoxicity of manganese in adult mice exposed via the drinking water. *Archives of Toxicology* **88** (5), 1185–1186.
- Leupin, O. X. & Hug, S. J. 2005 Oxidation and removal of arsenic (III) from aerated groundwater by filtration through sand and zero-valent iron. *Water Research* **39**, 1729–1740.
- Leupin, O. X., Hug, S. J. & Badruzzaman, A. B. M. 2005 Arsenic removal from Bangladesh tube well water with filter columns containing zerovalent iron filings and sand. *Environmental Science and Technology* **39**, 8032–8037.
- Mehta, V. S. & Chaudhari, S. K. 2015 Arsenic removal from simulated groundwater using household filter columns containing iron filings and sand. *Journal of Water Process Engineering* **6**, 151–157.
- Nguyen, V. A., Bang, S., Viet, P. H. & Kim, K. W. 2009 Contamination of groundwater and risk assessment for arsenic exposure in Ha Nam province, Vietnam. *Environment International* **35**, 466–472.
- Nitzsche, K. S., Lan, V. M., Trang, P. T. K., Viet, P. H., Berg, M., Voegelin, A., Planer-Friedrich, B., Zahoransky, J., Muller, S. K., Byrne, J. M., Schroder, C., Behrens, S. & Kappler, A. 2015 Arsenic removal from drinking water by a household sand filter in Vietnam – Effect of filter usage practices on arsenic removal efficiency and microbiological water quality. *Science of the Total Environment* **502**, 526–536.
- Nizam, S., Kato, M., Yatsuya, H., Khalequzzaman, M., Ohnuma, S., Naito, H. & Nakajima, T. 2013 Differences in urinary arsenic metabolites between diabetic and non-diabetic subjects in Bangladesh. *International Journal of Environmental Research and Public Health* **10** (3), 1006–1019.
- Ohgami, N., Hori, S., Ohgami, K., Tamura, H., Tsuzuki, T., Ohnuma, S. & Kato, M. 2012 Exposure to low-dose barium by drinking water causes hearing loss in mice. *Neurotoxicology* **33** (5), 1276–1283.
- Ohgami, N., Mitsumatsu, Y., Ahsan, N., Akhand, A. A., Li, X., Iida, M., Yajima, I., Naito, M., Wakai, K., Ohnuma, S. & Kato, M. 2016 Epidemiological analysis of the association between hearing and barium in humans. *Journal of Exposure Science and Environmental Epidemiology* **26**, 488–493.
- O’Neal, S. L. & Zheng, W. 2015 Manganese toxicity upon overexposure: a decade in review. *Current Environmental Health Reports* **2** (3), 315–328.
- Thang, N. D., Yajima, I., Ohnuma, S., Ohgami, N., Kumasaka, M. Y., Ichihara, G. & Kato, M. 2015 Enhanced constitutive invasion activity in human nontumorigenic keratinocytes exposed to a low level of barium for a long time. *Environmental Toxicology* **30** (2), 161–167.
- Torti, S. V. & Torti, F. M. 2013 Iron and cancer: more ore to be mined. *Nature Reviews Cancer* **13** (5), 342–355.
- Ventura-Lima, J., Bogo, M. R. & Monserrat, J. M. 2011 Arsenic toxicity in mammals and aquatic animals: a comparative biochemical approach. *Ecotoxicology and Environmental Safety* **74**, 211–218.
- Wang, W. 1988 Site-specific barium toxicity to common duckweed, *Lemna minor*. *Aquatic Toxicology* **12**, 203–212.
- Weinberg, E. D. 1996 The role of iron in cancer. *European Journal of Cancer Prevention* **5** (1), 19–36.

- Wen, C. P., Lee, J. H., Tai, Y. P., Wen, C., Wu, S. B., Tsai, M. K., Hsieh, D. P., Chiang, H. C., Hsiung, C. A., Hsu, C. Y. & Wu, X. 2014 High serum iron is associated with increased cancer risk. *Cancer Research* **74** (22), 6589–6597.
- Wilbers, G. J., Sebesvari, Z., Rechenburg, A. & Renaud, F. G. 2013 Effects of local and spatial conditions on the quality of harvested rainwater in the Mekong Delta, Vietnam. *Environmental Pollution* **182**, 225–232.
- Winkel, L. H. E., Trang, P. T. K., Lan, V. M., Stengel, C., Amini, M., Ha, N. T., Viet, P. H. & Berg, M. 2011 Arsenic pollution of groundwater in Vietnam exacerbated by deep aquifer exploitation for more than a century. *Proceedings of the National Academy of Sciences* **108** (4), 1246–1251.
- World Health Organization 1984 *Guidelines for Drinking-Water Quality*. WHO, Geneva, Switzerland.
- World Health Organization 2008 *Guidelines for Drinking-Water Quality, Incorporating the First and Second Addenda*. 3rd edn. WHO, Geneva, Switzerland. http://www.who.int/water_sanitation_health/publications/2011/wsh_vol1_1and2_addenda.pdf?ua=1 (accessed 18 May 2016).
- World Health Organization 2011 *Guidelines for Drinking-Water Quality*, 4th edn. WHO, Geneva, Switzerland. http://www.who.int/water_sanitation_health/publications/2011/dwq_guidelines/en (accessed 18 May 2016).
- Yajima, I., Uemura, N., Nizam, S., Khalequzzaman, M., Thang, N. D., Kumasaka, M. Y., Akhand, A. A., Shekhar, H. U., Nakajima, T. & Kato, M. 2012 Barium inhibits arsenic-mediated apoptotic cell death in human squamous cell carcinoma cells. *Archives of Toxicology* **86**, 961–973.
- Yajima, I., Kumasaka, M. Y., Ohnuma, S., Ohgami, N., Naito, H., Shekhar, H. U., Omata, Y. & Kato, M. 2015 Arsenite-mediated promotion of anchorage-independent growth of HaCaT cells through placental growth factor. *Journal of Investigative Dermatology* **135** (4), 1147–1156.

First received 17 June 2016; accepted in revised form 15 August 2016. Available online 4 October 2016

Limited effectiveness of household sand filters for removal of arsenic from well water in North Vietnam

ORIGINALITY REPORT

5%

SIMILARITY INDEX

0%

INTERNET SOURCES

5%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

Masafumi Yoshinaga, Hiromasa Ninomiya, M.M. Aeorangajeb Al Hossain, Makoto Sudo et al. "A comprehensive study including monitoring, assessment of health effects and development of a remediation method for chromium pollution", Chemosphere, 2018

Publication

5%

Exclude quotes On

Exclude bibliography On

Exclude matches < 3%