



Cytotoxic Activity Screening of Fungal Extracts Derived from the West Sumatran Marine Sponge *Haliclona fascigera* to Several Human Cell Lines: HeLa, WiDr, T47D and Vero

Dian Handayani^{1*}, Wildan Rasyid¹, Rustini¹, Elmi Nurhaidah Zainudin², Triana Hertiani³

¹Faculty of Pharmacy, Andalas University, Padang 25163, Indonesia.

²Faculty of Marine Science and Fisheries, Hasanuddin University, Makassar, Indonesia.

³Faculty of Pharmacy, Gadjah Mada University, Sekip Utara, Yogyakarta, Indonesia.

ARTICLE INFO

Article history:

Received on: 07/05/2017

Accepted on: 08/06/2017

Available online: 28/01/2018

Key words:

Cytotoxic activity, *Haliclona fascigera*, marine derived fungi, Brine Shrimp Lethality Test (BSLT), MTT assay

ABSTRACT

Cytotoxic activity of twenty extracts of fungal derived from marine sponge *Haliclona fascigera* collected from West Sumatera was evaluated by using Brine Shrimps Lethality Test (BSLT) and MTT assay on HeLa, WiDr, T47D and Vero cells. In BSLT result, all fungal extracts were cytotoxic because their $LC_{50} < 1000$ ppm, and ranging from 1 to 335 ppm. While sixteen extracts (80%) were toxic ($LC_{50} < 100$ ppm) and further tested its cytotoxic activity against HeLa, WiDr, T47D and Vero cell lines. A single 100 $\mu\text{g/mL}$ dose of each extract was employed with 72 h of incubation for all tests. Doxorubicin was used as the positive control. The fungal extract of WR6 showed the highest MTT cytotoxicity with IC_{50} values of 47.4, 67.1, 118.3 and 163.37 ppm against WiDr, T47D, HeLa and Vero cell lines respectively. Therefore, this extract was chosen to be a good candidate for further studies as a new anticancer drug in the treatment of human cancer.

INTRODUCTION

In recent years, cancer has become the most deadly diseases of man. According to the World Health Organization (WHO), cancer has become one of the major causes of morbidity and mortality throughout the world and there were approximately 14 million new cases and 8.2 million cancer-related deaths in 2012. In Indonesia, the prevalence of cancer was 1.4 per 1000 population, or around 330 thousand people (Basic Health Research, 2013). The number of cancer patients in Indonesia was very high. This can be seen from various cancer data published by both government and cancer institutions. Even according to the WHO in 2030 will be a surge in cancer patients in Indonesia up to seven times. The number of cancer patients who died is also increasingly

apprehensive. For many years cancer researchers around the world have been focused on finding better therapeutic strategies and new molecular approaches to reduce mortality. Nowadays there is a considerable scientific discovery of new anticancer agents from natural products, one of which was marine sponge.

Sponges are one group marine organisms contained in the waters of Indonesia by amount of about 850 kinds (Van Soest, 1989). The content of sponge's secondary metabolites was reported to have wide range of bioactivity which was promising for development as the lead compound (Edrada *et al.*, 2000). Sponges have tissues and organs that are very simple and have varied shapes. Sponge growth is strongly influenced by the chemical and physical environment, such as the: the depth, the waves, and sedimentation. Sponges are also a very fertile host to a wide range symbiotic microorganisms, like bacteria and fungi.

These microbes live and attached to the sponge and interacted symbiotic mutualism. The existence of microbial symbiotic function are in assisting process of acquiring nutrients (especially in the fixation of carbon and nitrogen), the allocation of the com-

*Corresponding Author

Dian Handayani, Faculty of Pharmacy, Andalas University, Padang, Indonesia.

E-mail: dianh_17@yahoo.com

pounds of defense, synergies between the chemical structural, stabilizer skeletons sponge, and assist the process of excretion as well as contribute in the production cycle of secondary metabolites (Hentscel *et al.* 2002). Thus, it can be assumed that symbiotic microbes produce a secondary metabolite similar to a sponge. Microbial symbiotic can produce bioactive compounds with huge potential to develop into a new drug compounds.

This study was a continuous study to investigate the anti-bacterial and cytotoxic activity of ethyl acetate extract obtained from marine derived fungi of the sponge from West Sumatera, Indonesia. Examination of cytotoxic activity was performed by BSLT and MTT assay methods (Handayani *et al.*, 2015a; Handayani *et al.*, 2015b; Handayani *et al.*, 2016a; Handayani *et al.*, 2016b).

MATERIALS AND METHODS

Sponge material

The sponge *H. fascigera* was collected from the Setan Island in Februari 2015, at South Coast of West Sumatra, Indonesia, in depth of ± 13 m. The Sponge was taken 200 g and transferred into a sterilized plastic bag and stored in the ice box. The sample was transported to the laboratory and processed immediately for the isolation of symbiotic fungi.

Isolation of marine derived fungi from sponge *H. fascigera*

The isolation of symbiotic fungi was started by surface sterilization of the sample. The sponge was rinsed with sterile seawater, and then cut into small pieces and was taken as much as 10 grams an inserted into Erlenmeyer flask and 100 mL of sterile seawater was added. Then it was diluted until its concentration 10^{-6} and inoculated on Sabouraud Dextrose Agar (SDA) containing chloramphenicol (0.05 g/L) as medium, and incubated at a temperature of 27-29°C for 5-7 days. Colonies that have different shapes and colors with other colonies could be regarded as different isolates. Then the isolates were purified by the scratch method to obtain pure isolates (Kjer *et al.*, 2010). The Pure isolate was identified based on Brigitte (1980). The isolated fungi have been deposited at our laboratory in Sumatran Biota Laboratory, Andalas University.

Cultivation of isolated fungi in medium of MEB

Each of the pure fungal isolate obtained in the insulating phase was cultured in medium of Malt Extract Broth (MEB). The fungal was taken one loop, then put in a 50 ml MEB medium and incubated at room temperature for 7 days. Furthermore, they were grown on a scale of 50 ml were transferred aseptically similar to the culture medium at a larger scale (500 ml) and incubated at room temperature for 3-4 weeks (Kjer *et al.*, 2010).

Extraction of secondary metabolites from fungal isolates

The fungal isolates at their maximum growth were then extracted with ethyl acetate (EtOAc) in 3 repetitions. The ethyl acetate extract was collected and evaporated *in vacuo* using a rotary evaporator (Kjer *et al.*, 2010). The concentrated EtOAc extracts were tested for cytotoxic activity using method of BSLT. Based on cytotoxic test results on BSLT method, fungal extract which has a $LC_{50} < 100$ ppm was continued to test by determining the percentage of viability against three cancer cells and normal

cells with MTT assay method. The IC_{50} determination was done for extract with a low percentage of viability.

Screening of toxicity and cytotoxic activities

BSLT method

The BSLT assay is non-specific toxicity assay that is used in natural products to detect the presence of bioactive compounds. In this assay was used the *nauplii of Artemia salina* L (*Artemiidae*). Brine shrimp eggs were hatched in 500 mL of filtered seawater under constant aeration for 48 h at $(27 \pm 2)^\circ\text{C}$. After hatching, active *nauplii* of the egg cell are collected and used for testing. Five hundred, fifty, and five μl of all fungal ethyl acetate extracts were added to the vials. After the solvent was dry, 50 μl DMSO was added and supplemented with 5000 μl of seawater, resulting in final concentration of 1000 ppm, 100 ppm and 10 ppm in triplicate. Ten *nauplii* are then inserted into each vial and kept at room temperature for 24 hours. The filtered seawater and DMSO are used as negative controls. The LC_{50} value is calculated using curve method based on probit analysis (Meyer *et al.*, 1982).

MTT assay

Sample screening

4 cell lines, HeLa as cervix cell line, WiDr as a colon adenocarcinoma cell line, T47D as human ductal breast epithelial tumor cell line and Vero as normal cell line have been prepared for cytotoxic assay using MTT. All cell lines were obtained from Laboratory of Parasitology at UGM. HeLa, T47D and WiDR were cultured in RPMI 1650 and Vero was cultured in M199 Medium. All cells were subculture after mild trypsinization with trypsin-EDTA (Sigma-Aldrich, USA), and then determined the cell number and viability. The cells were seeds in 96-well plates at density 6×10^3 cells/well in 100 μl medium and incubated overnight. All media were supplemented at 10% with fetal bovine serum (Gibco) and streptomycin and penicillin (2%, Sigma-Aldrich, USA). The cell line were kept at 37°C, 98% relative humidity with 5% CO_2 atmosphere.

A stock solution was prepared by dissolving the samples in DMSO and was given 100.000 ppm concentration. Cells that had been incubated 24 hours, then divided into several groups, namely treatment, positive control, cell control and media control (blank). Removed medium and washed using PBS sterile which each well was add 100 μl PBS. Then, 100 μl of each material (extract) added to each well with one concentration (100 ppm). As control positive was used by doxorubicin. Then it was incubated for 24 hours in an incubator at 37°C, 5% CO_2 .

Cells that had been treated and incubated 24 hours later dumped throughout the medium and washed using sterile PBS. Then in each well was added 100 μl of MTT (5 mg/ml) followed by a 4 hour incubation in an incubator at 37°C, 5% CO_2 . To each well was added 100 μl of 10% SDS to dissolve the formazan crystals formed and incubated one night in room temperature. The plates were then read by ELISA reader at 540 nm (Permanasari *et al.*, 2016)

% Cell viability then calculated by the equation

$$\frac{\text{OD of treatment} - \text{OD of blank}}{\text{OD of control} - \text{OD of blank}} \times 100\%$$

The aim of this screening was to identify which sample was given cytotoxic activity in cell line.

Cytotoxic assay

Cytotoxic assay was determination by IC_{50} which was made by the same procedure as sample screening. Fungi extracts were made in several concentrations which are 300, 150, 75, 37.5, 18.75 and 9.375 ppm. Doxorubicin as a positive control was created in concentration of 100, 50, 25, 12.5 and 6.25 ppm.

Statistical methods

Value Ln concentration is regressed against cell viability. Regression equations were then used to calculate IC_{50} values. Another method that can be used to calculate the IC_{50} values are probit analysis using SPSS.

RESULTS AND DISCUSSION

Marin fungi that are associated with marine sponge can be derived from the environment in which the sponge arise through the process of filter feeder, and enter into mesophyll. However, some studies show that besides microbial epibiont attached to the surface of the sponge during the growing period (Chelossi *et al.*, 2004), some bacteria and yeasts (fungi) transmitted genetically in the body of the sponge (Maldonado *et al.*, 2005).

21 fungi were successfully isolated from the marine sponge *Haliclona fascigera* collected from the waters of Setan Island, in West Sumatra at a depth of 13 meters. The fungi isolates were cultivated in a liquid medium, Malt Extract Broth (MEB) for 30-40 days. Ethyl acetate extract of fungal isolates ranged from 15.4-551.3 mg.

The examination results the toxicity activity by the method of Brine Shrimp Lethality Test showed that the LC_{50} of ethyl acetate extract of fungi isolates ranges from 1 to 335 ppm. According to Meyer *et al.* (1989), extracts of plants or animal are classified a cytotoxic when the LC_{50} value <1000 ppm. An extract is considered very toxic when it has a LC_{50} value < 30 ppm, and toxic if it has a LC_{50} value 30-1000 ppm and is considered non-toxic if LC_{50} > 1000 ppm. While pure compound was declared toxic if it has a value of LC_{50} <200 ppm (Meyer, 1982). Therefore, extract of WR4, WR8, WR13, and WR19 with LC_{50} value of <10 can be categorized as toxic and should be followed up in order to find a potential anti-cancer compound (Figure 1).




Fig. 1: Brine shrimp toxicity results (LC_{50} in ppm).

The BSLT is found to have a good correlation with cytotoxic activity in some human solid tumors and with pesticide activity, and led to the discovery of new class of natural pesticides and active antitumor agent (Mc Laughlin *et al.*, 1998). The cytotoxic activity of the drug can affect and disrupt the fundamental mechanisms associated with cell growth, mitotic activity, differentiation and function. The results of the cytotoxic activity for this extract may be due to one of these mechanisms.

Further testing of cytotoxic test was conducted by MTT assay. This method is a colorimetric method, in which the MTT reagent is a tetrazolium salt that can be broken down into formazan crystals by succinate tetrazolium reductase. This enzyme is present in active mitochondria and plays a role on cell respiration cycle. Formazan purple crystals can be read its absorbance by using ELISA reader (Doyle and Griffith, 2000). NADH or other reductor might be transfer the electron into MTT and made formazan purple crystal (Berridge *et al.*, 1993).

Cytotoxic activity of the compounds can be determined based on the value of IC_{50} . IC_{50} is the concentration of the extract/test compounds that can kill as many as 50% of the cell population. The determination of IC_{50} value begins with determining of the viability percentage of cancer cells in the culture medium after adding ethyl acetate extract of symbiotic fungal with a concentration of 100 μ g/ml.

Fungi extract with the lowest percentage of viability ($\leq 50\%$) especially against the HeLa cancer cells are extracts WR2, WR4, WR6 and WR9 (Figure 2). The percentage of viability extracts against other cancer cells (WIDR, T47D and Vero) is highly variable and some of which are not toxic to the normal cells (vero).




Fig. 2: Cytotoxic effect of ethyl acetate extracts on various cancer cell lines

Determination of IC_{50} value is then performed to extracts which shows the lowest of the viability percentage of above. Data obtained from several concentrations of the extract was analyzed by linear regression equation to obtain IC_{50} values. IC_{50} extract values of WR2, WR4, WR6 and WR9 against HeLa, WIDR, T47D and Vero cell lines were different for each other (Table 1). Based on Table 1, it can be determined that the most cytotoxic extract against all cancer cell cultures compared with extract of WR2, WR4 and WR9 was extract of WR6. The cytotoxic activity of WR6 extract is quite selective against WIDR only, when it compared with the cytotoxic activity of doxorubicin against all cancer cells even the normal cells (vero cells).

The four fungi above were selected for further characterization and identification based on Brigitte (1980). Macroscopic

examination includes a visual observation to the form colony or hyphal, surface and reverse colony color, and colony texture. While on microscopic examination was carried out by observing the characteristic of the spores or conidia, and reproductive structures (sexual and asexual) under a light-field microscope. Based on the results of the morphological identification of four fungal isolates which showed cytotoxic activity, it is known that the fungus WR2 is *Trichophyton* sp, WR4 is *Aspergillus* sp, WR6 is *Trichophyton* sp and WR9 is *Penicillium* sp.

Table 1. IC₅₀ of selected fungal extracts on various cancer cell lines.

No.	Code of Extract	IC ₅₀ of Cancer and Normal Cell Line (ppm)			
		WIDR	T47D	Hela	Vero
1	WR2	193.95	5,861.67	211.55	357.49
2	WR4	38.21	328.23	598.89	321.54
3	WR6	47.36	67.08	118.29	342.94
4	WR9	284.28	132.74	118.29	342.94
5	Doxorubicin	0.28	10.05	1.25	43.74

In this preliminary study, it can be conclude that the cytotoxic activity can be produced from secondary metabolites contained in the extract of the fungus. The results of this study provide important information for the continued research on isolation, and characterization of cytotoxic compounds from WR6 and other extracts. Therefore, this fungal extract can be developed as a source of candidate for new drugs in overcoming the problem of cancer.

CONCLUSIONS

This study demonstrated that several EtOAc extracts of the fungi derived from marine sponge *Haliclona fascigera* were cytotoxic. The results indicated that fungal isolates of WR2, WR4, WR6 and WR9 may be useful as an alternative to produce the anticancer used in pharmaceutical. However, further research needs to be done in determining of anticancer compounds produced by those marine sponge derived fungi.

CONFLICT OF INTERESTS

Declared none.

ACKNOWLEDGEMENT

This research was funded by Directorate General of Higher Education Ministry of National Education, Indonesia, in project "International Research Collaboration and Publication, No: 059/SP2H/LT/DPRM/IV/2017.

REFERENCES

Berridge MV, Tan AS. Characterization of the cellular reduction of 3-(4,5-dimethylthiazol 2-yl)-2,5-diphenyltetrazolium bromide (MTT): subcellular localization, substrate dependence, and involvement of mitochondrial electron transport in MTT reduction. *Arch Biochem Biophys* 1993; 303(2):474-82.

Brigitte, G. Kompendium der medizinischen mykologie. 1980, Berlin-Hamburg. Verlag Paul Parey.

Chelossi E, Milanese M, Milano A, Pronzato R, & Riccardi G. Characterisation and antimicrobial activity of epibiotic bacteria from *Petrosia ficiformis* (Porifera, Demospongia). *J. Experimental Mar Biol & Ecol*, 2004; 309: 21-33.

Doyle A, and Griffiths JB. 2000. Cell and tissue culture for medical research. New York: John Wiley and Sons Ltd.

Edrada RA, Wray V, Handayani D, Schupp P, Balbin-Oliveros M, and Proksch, P. *Structure-activity relationships of bioactive metabolites from some Indo-Pacific marine invertebrates*. *Studies in Natural Products Chemistry* 21(B). 2000; 251-292. ISSN 1572-5995.

Handayani D, Sandrawaty N, Murniati M, and Regina R. Screening of Endophytic Bacteria Isolated from Marine Sponge *Haliclona fascigera* for Inhibition against Clinical Isolates of Methicillin Resistant *Staphylococcus aureus* (MRSA). *J. of Applied Pharmaceutical Science*, 2015a; 5(9): 139-142.

Handayani D, Ahdinur, RF, and Rustini R. Antimicrobial Activity of Endophytic Fungi from Marine Sponge *Haliclona fascigera*. *Journal of Applied Pharm. Science*, 2015b; 5(10), 154-156

Handayani D, Ormando R and Rustini, Antimicrobial Activity Screening of Symbiotic Fungi from Marine Sponge *Petrosia nigrans* collected from South Coast of West Sumatera Indonesia, *International Journal of Pharmacognosy and Phytochemical research*, 2016a; 8(4): 623-625.

Handayani D, Murniati M, Rustini R, In vitro inhibitory activity of ethyl acetate extract of symbiotic bacteria isolated from the marine sponge *Haliclona fascigera* against multidrug resistant organism (MDRO). *Journal of Applied Pharm. Science*, 2016b; 6(11):218-222.

Hentschel U, Piel J, Degnan, SM, and Taylor MW. Genomic insights into the marine sponge microbiome. *Nature Reviews Microbiology*, 2012;10:641-654.

Kjer J, Debbab A, Aly AH, Proksch P. Methods for isolation of marine-derived endophytic fungi and their bioactive secondary products. *Nature Protocols*, 2010,5(3): 479-490.

Meyer BN, Ferrign RN, Putnam JE, Jacobson JE, Nicholas DE, McLaughlin JL, Brineshrimp: a convenient general bioassay for active plant constituents. *Planta Medica*, 1982, 45, 31-34.

Maldonado M, Corrtadellas N, Trillas MI, & Rützler C. Endosymbiotic yeast maternally transmitted in a marine sponge. *Biol Bull*. 2005, 209: 94-106.

McLaughlin JL, Rogers LL, Anderson JE, The Use of Biological Assays to Evaluate Botanicals. *Drug Information Journal*, 1998, 32, 513-524.

Permanasari P, Hertiani T, Yuswanto A. Immunomodulatory Effect of Massoia Bark Extract and The Cytotoxicity Activiy Against Fibroblast and Vero Cells in Vitro. *International Journal of Pharmaceutical and Clinical Research*, 2016; 8(5) Suppl:326-330.

Reports the Results of Basic Medical Research. 2013. Research and Development Agency for Health, Ministry of Health, Republic of Indonesia.

Van Soest RW. 1989. The Indonesian Sponge Fauna: A Status Report. Institute of Taxonomic Zoology, University of Amsterdam.

How to cite this article:

Handayani D, Rasyid W, Rustini, Zainudin EN, Hertiani T. Cytotoxic Activity Screening of Fungal Extracts Derived from the West Sumatran Marine Sponge *Haliclona fascigera* to Several Human Cell Lines: Hela, WiDr, T47D and Vero. *J App Pharm Sci*, 2018; 8 (01): 055-058.