



Abstract  
Book

# International Conference on Food Security During Challenging Times

5-7th July 2010

Universiti Putra Malaysia,  
Serdang, Selangor, Malaysia

Jointly  
Organised by:



UPM  
UNIVERSITI PUTRA MALAYSIA



Supported by



AGRICULTURE

FOOD SECURITY

FACULTY OF AGRICULTURE  
UNIVERSITI PUTRA MALAYSIA

ALL RIGHT RESERVED

No part of this publication may be reported, stored in a retrieval system or transmitted, in any means, electronic, mechanical, photocopying, recording or otherwise without the prior written permission of the copyright owner.

Published by  
FACULTY OF AGRICULTURE  
UNIVERSITI PUTRA MALAYSIA  
43400 UPM SERDANG, SELANGOR  
MALAYSIA

Perpustakaan Negara Malaysia Cataloging-in-Publication Data

International Conference on Food Security During Challenging Times (ICFSDCT 2010)  
Book of extended abstracts of ICFSDCT 2010  
Universiti Putra Malaysia, Serdang, Selangor, Malaysia, 5-7 July 2010  
Edited by Rosli B. Mohamad et al., 2010  
ISBN 978-983-3280-01-8

Abstracts and papers published in this book have not been refereed. The responsibility of the content rests with the authors.

J

## CONTENTS

### KEYNOTES

Special address	Positioning Malaysia in global food security issues <i>Zakri Abdul Hamid</i>	2
K1	Rethinking agriculture: Addressing food security and climate change issues <i>Lim Li Ching</i>	4
K2	Ensuring food security by meeting current challenges <i>Yusof Basiron</i>	8
K3	From food vulnerability to nutritional (in)security <i>Berger Jacques, Martin-Prével Yves, Delpeuch Francis and Maire Bernard</i>	12

### ORAL PAPERS

#### SESSION 1: POLICY AND ECONOMIC PERSPECTIVES

S1.1	Perspectives on food security and freer trade in agriculture <i>YM Tengku Mohd Ariff Tengku Ahmad</i>	18
S1.2	The new European food regulation on precaution, traceability, food safety (application on January 1 <sup>st</sup> , 2005 of the regulation EC 178/2002) <i>Max Reynes and Didier Montet</i>	20
S1.3	Global food crisis: Implications to Malaysia's food security <i>Mad Nasir Shamsudin and Fatimah Mohamed Arshad</i>	21
S1.4	Food safety and security situation in Pakistan <i>Mian Kamran Sharif, Faqir Muhammad Anjum, Masood Sadiq Butt, Moazzam Rafiq Khan and Sarfraz Ahmad</i>	22
S1.5	Malaysian rice production sustainability and security in view of population increase, rice consumption and global warming trends for the next 50 years duration <i>Ahmad Selamat and Yahya Awang</i>	26
S1.6	Thailand agriculture's tree of alternatives <i>Roland Poupon</i>	32

#### SESSION 2: ENVIRONMENT AND CLIMATE CHANGE

S2.1	Defining imperatives and megatrends on ASEAN agriculture: Food and energy scenario 2050 <i>Makhdir Mardam and Rosli B. Mohamad</i>	38
------	---	----

82	S5.4	Threat of <i>Streptococcus suis</i> infection <i>Tsutomu Sekizaki</i>	121
86	S5.5	Halal principles and food safety: The complementary effect <i>R. Golnaz, M. Zainalabidin Mohamad and S. Mad Nasir</i>	124
90	S5.6	Prospects for sustainable crop production technologies in East Timor <i>Edmundo Viegas and Peter Kemp</i>	127

<b>SESSION 6: FOOD SAFETY TECHNOLOGY</b>			
--	--	--	--

96	S6.1	Novel PCR-based genotyping method, using genomic variability between repetitive sequences of toxigenic <i>Vibrio cholerae</i> O1 El Tor and O139 <i>Ro Osawa</i>	130
102	S6.2	Antibiotic resistance of bacteria in aquaculture: Development of novel antimicrobials based on essential oils <i>S. Sarter and P. Danthu</i>	132
106	S6.3	Triploidy verification and vitellogenesis induction in African catfish ( <i>Clarias gariepinus</i> ) <i>A. Karami, A. Christianus, S.C. Courtenay, Z. Ishak, S. Mohd. Arif and K.M.S. Zamri</i>	134
107	S6.4	Contribution of irradiation to food security and safety: An update <i>Zainon Othman, Norimah Yusof and Ahmad Zainuri Mohd Dzomir</i>	137
111	S6.5	Food security in Bangladesh: Present status and reality <i>M.M. Alam, M.K. Rahman and S. Khatun</i>	138
	S6.6	Microfiltration, a processing technology to have safe and sure buffalo dairy products with their natural quality <i>Sarfraz Ahmad, Faqir Muhammad Anjum, Nuzhat Huma, Mian Kamran Sharif, Moazzam Rafiq Khan, Gaelle Tanguy, Nadine Leconte and Frederic Gaucheron</i>	142

<b>POSTERS</b>			
----------------	--	--	--

118	P1	Ability of light intensity to enhance medicine component and anti-oxidant activities in Malaysia young ginger variety ( <i>Zingiber officinale</i> var. <i>halia bentong</i> ) under controlled environment <i>A. Ghasemzadeh Daghigh, H. Z. Jaafar and A. Rahmat</i>	148
119	P2	The effect of different spraying at different sowing date to pest control and management of sweet corn <i>A.R. Saberi, H. Shamsabadi and Z. Barati</i>	152
120			



## YIELD POTENTIAL OF TROPICAL MAIZE HYBRIDS IN ACID SOILS

P.K.D. Hayati<sup>4</sup>, G.B. Saleh<sup>1</sup>, J. Shamshuddin<sup>2</sup> and S. Napis<sup>3</sup>

<sup>1</sup>Dept. of Crop Science

<sup>2</sup>Dept. of Land Management,

<sup>3</sup>Dept. of Cell and Molecular Biology

University Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia

<sup>4</sup>Dept. of Agrotechnology, Faculty of Agriculture University of Andalas, Kampus Limau Manis, Padang, 25163 Indonesia

### INTRODUCTION

Maize is one of the three major cereal crops grown in the world. In Malaysia, it is an important commodity as most of the grain maize used to support the local livestock feed industry is imported in the past three decades (USDA/FAS, 2009). Meanwhile, most of the arable soils in the country are weathered and acidic in nature, comprising 72% or 23.7 million ha of land area in Malaysia (IBSRAM, 1985; FAO, 2006). The soils have low cation exchange capacities (CEC), low basic cations, especially Ca and/or Mg, and high aluminum (Al) concentration in the soil solution (Shamshuddin et al., 1991). Among these constraints, Al toxicity is the major factor limiting plant growth and yield potential in acid soils, including maize (Shamshuddin et al., 1991). Even though the application of lime has proven to overcome the soil acidity problems, a number of limitations in use have been reported (Kamprath, 1971; Shamshuddin et al., 1998; Hede et al., 2002). Hence, planting high yielding varieties, in particular, maize hybrids tolerant to acid soils along with the use of sustainable agronomic practices offers an effective strategy for improving maize productivity in acid soils in the country. The objective of this study was to select maize hybrids for high yield potential in acid soils.

### METHODOLOGY

The 36 maize hybrids obtained from a 9x9 diallel cross and the two check varieties, the composite variety *Sukmaraga* as acid soil tolerant check variety and the hybrid *Putra J-58* were evaluated in 2007 in Puchong and Serdang, in Split Plots arranged in a RCB design, with three replications. Two soil conditions (unlimed and limed) and genotypes were assigned to the main-plots and the sub-plots, respectively. The unlimed treatment was the naturally acid soil, while the limed treatment was the soil treated with ground magnesium limestone (GML) at the rate of 2 t ha<sup>-1</sup>. The data were subjected to the analysis of variance (ANOVA) using the General Linear Models (Proc GLM) of the Statistical Analysis System (SAS) version 9.1 (SAS Institute Inc., 2003).

### RESULTS AND DISCUSSION

Application of GML at 2 t ha<sup>-1</sup> in Serdang series soil in Puchong decreased exchangeable Al on the topsoil from 3.05 to 0.9 cmol.kg<sup>-1</sup>, while that on mixed-tin tailings soil in Serdang decreased exchangeable Al from 1.58 to 0.52 cmol.kg<sup>-1</sup>. Application of GML also increased the basic cations and the soil pH. The relative grain yields of the hybrids in this study decreased exponentially with the increasing amount of exchangeable Al (Fig.1). Critical value for exchangeable Al was 0.60

cmol<sub>c</sub>kg<sup>-1</sup>, indicating that application of lime at the rate of 2 t ha<sup>-1</sup> was enough to alleviate exchangeable Al in Serdang; however, it was not enough in Puchong. For a better growth of maize in acid soils, the exchangeable Ca and Mg have to be increased to about 1.22 and 0.42 cmol<sub>c</sub>kg<sup>-1</sup>, respectively, while the soil pH has to be raised to about 5.1.




Fig.1. Effects of exchangeable Al on relative grain yield of 36 maize hybrids

The reduction of grain yields in acid soil in Puchong was high, ranging from 49 to 91%, while in Serdang, it ranged from 24 to 64%. Grain yields of the hybrids varied greatly within acid soil conditions and locations, in which different hybrids were found to have high yield performance in the different acid soil conditions and locations. The top ten high yielding hybrids in each acid soil condition and across environments (the combined data of both soil conditions and locations) out-yielded both the check varieties (Table 1). With regards to heterosis, the hybrids exhibited significant and high heterosis estimates for grain yield, indicating that the hybrids performed higher grain yield over their parents. Only H29 was consistently high yielding in each soil condition at the two locations. The hybrid also showed relatively high critical level of Al (0.62 cmol<sub>c</sub>kg<sup>-1</sup>). This hybrid was progeny of a cross between acid soil tolerant inbred line (IPB-12) and an elite inbred line introduced from CIMMYT (CML-2). Furthermore, its parental inbred lines had the highest genetic distance (0.72) based on the SSR markers (data were not shown), indicating that high diverse genetic background was possessed by the hybrid. Thus, high heterosis was expected from the hybrid.

Table 1. The top ten high yielding maize hybrids and the two check varieties evaluated on each soil condition from the combined data of two locations, heterosis estimates and critical level of AI

Hybrid	Yield (kg ha <sup>-1</sup> )	MPH (%)	Hybrid	Yield (kg ha <sup>-1</sup> )	MPH (%)	Hybrid	Yield (kg ha <sup>-1</sup> )	MPH (%)	Critical level of AI (cmol/kg <sup>-1</sup> )
Acid soils			Limed soils			Across environments			
H18	2662	252.7	H36	6001	582.4	H21	4275	153.8	0.64
H24	2875	328.3	H21	6164	325.6	H29	4117	133.3	0.62
H29	2871	395.5	H25	5665	156.3	H36	4007	420.1	0.60
H17	2816	204.7	H26	5633	210.7	H25	3980	103.1	0.60
H9	2772	263.5	H8	5594	394.7	H30	3914	121.8	0.64
H23	2668	300.7	H34	5576	290.5	H26	3873	97.6	0.56
H30	2660	449.5	H14	5461	206.5	H17	3827	127.3	0.63
H4	2587	327.0	H10	5404	99.0	H8	3814	278.2	0.55
H7	2533	771.1	H33	5401	280.2	H4	3805	115.6	0.74
H28	2450	235.4	H29	5364	174.2	H9	3766	123.6	0.44
Putra J-58	2179	n.a.	Putra J-58	5310	n.a.	Putra J-58	3744	n.a.	0.64
Sukmaraga	2563	n.a.	Sukmaraga	5088	n.a.	Sukmaraga	3826	n.a.	0.64
Hybrid mean	2210	-	Hybrid mean	4925	-	Hybrid mean	3576	-	-

n.a.=not available

Regarding to tolerance to acid soils, hybrids generally showed a lower relative yield (63%) than Sukmaraga as acid soil tolerant check variety (66.9%); however, they had higher relative yield than Putra-J58 (56%). This indicates that the hybrids revealed in the study had high tolerance to acid soils than commercial maize hybrid variety established in the country. The highest relative yield in Puchong was shown by H29 (72.8%), while in Serdang was shown by H24 (86.2%). To distinguish hybrids corresponding to their tolerance to acid soils, a quadrant model was developed according to Sangalang and Bouwkamp (1988) and Howeler (1991). Five hybrids, H29, H24, H17, H18 and H9 were found to be tolerant to acid soil and had high yielding potential in acid soil conditions (Quadrant III) (Fig.2).




Fig.2. Relationship between relative yield and yields of hybrids in acid soils of 36 maize hybrids and two check varieties

## CONCLUSION

This study has revealed hybrids that have high genetic potential for tolerance to acid soils. Among the hybrids, H29 was identified to be tolerant to acid soils and high potential on both acid and limed soils. Hybrid H29 should be given the opportunity for further evaluation before it is released as a hybrid maize variety for commercial production.

## ACKNOWLEDGEMENTS

The authors would like to record their appreciation to the Ministry of Science, Technology and Innovation, Malaysia (MOSTI) through the IRPA Grant. This research was partly supported by funds from the SEARCA through the PhD Research Scholarship awarded to the senior author.

## References

- FAO, (2006). Food and Agricultural Indicator Malaysia. [http://www.fao.org/es/ESS/compendium\\_2006/pdf/MAL\\_ESS\\_E.pdf](http://www.fao.org/es/ESS/compendium_2006/pdf/MAL_ESS_E.pdf).
- Hede, A. R., Skovmand B., Ribaut J. M., Gonzáles-de-León D. and Stølen O. (2002). Evaluation of aluminum tolerance in a spring rye collection by hydroponic screening. *Plant Breeding* 121:241-248.
- Howeler, R.H. (1991). Identifying plants adaptable to low pH conditions. In R. J. Wright, V. C. Valigar and R. P. Murrmann (eds.), *Plant-Soil Interactions at Low pH*. Kluwer Academic Publishers. Dordrecht. pp.885-904.
- IBSRAM, (1985). Report of the Inaugural Workshop and Proposal for Implementation of the Acid Tropical Soils Management Network. Bangkok. International Board for Soil Research and Management.
- Kamprath, E. J. (1971). Potential detrimental effects from liming highly weathered soils to neutrality. *Soil Crop Sci. Soc. Florida Proc.* 31:200-204.
- Sangalang, J. B. and Bouwkamp J.C. (1988). Selection of sweet potato for tolerance to aluminum toxicity: Screening procedures and field test. *J.Amer.Soc.Hortic.Sci.* 113:277-281.
- SAS Institute Inc., 2003. *SAS/STAT® User's Guide*. Version 9.1. SAS Institute Inc. Cary, NC.
- Shamshuddin, J., Che Fauziah I. and Sharifuddin H. A. H. (1991). Effects of limestone and gypsum applications to a Malaysian Ultisol on soil solution composition and yields of maize and groundnut. *Plant and Soil* 134:45-52.
- Shamshuddin, J., Sharifuddin H. A. H. and Bell L. C. (1998). Longevity of ground magnesium limestone applied to an Ultisol. *Commun. Soil Sci. and Plant Anal.* 26(9&10):1299-1313.
- USDA/FAS. (2009). Grain: World Markets and Trade. Foreign Agricultural Services. The United States of Department of Agriculture.