

# Oral Contraceptive Used More than 5 Years is Associated with Increased Risk of Breast

*by* Wirsma Arif Harahap

**Submission date:** 06-Jan-2020 04:49PM (UTC+0800)

**Submission ID:** 1239493101

**File name:** dr.Wirsma-JurnalInternasional9.pdf (3.02M)

**Word count:** 785

**Character count:** 4158

## Oral Contraceptive Used More than 5 Years is associated with Increased Risk of Breast Cancer: A Meta-Analysis of 28,776 South East Asian Women

RICHAN DIANA NINDIRAH<sup>1</sup>, SUGAIDI LUKMAN ANWAR<sup>2</sup>, WIRMA<sup>3</sup>, ANI HARAHAP<sup>4</sup>, LUTFAN LAZUARDI<sup>5</sup>,  
YAN LAMPRAHASTO<sup>6</sup>, UGGI D. ARIANDONO<sup>7</sup>

<sup>1</sup>Doctoral Program, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia.

<sup>2</sup>Department of Public Health and Community Medicine, Faculty of Medicine, Universitas Andalas, Padang, Indonesia.

<sup>3</sup>Department of Surgical Oncology, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia.

<sup>4</sup>PILAR Research and Education, 20 Station Road, Cambridge CB1 2JD, UK

<sup>5</sup>Department of Surgical Oncology, Faculty of Medicine, Universitas Andalas, Padang, Indonesia.

<sup>6</sup>Department of Health Policy and Management, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia.

<sup>7</sup>Department of Pharmacology and Therapy, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia.

Article History:

Submitted: 17.08.2019

Revised: 22.10.2019

Accepted: 15.11.2019

### ABSTRACT

Breast cancer associated with variety of hormonal, metabolic, and reproductive factors. Duration of oral contraceptive use with breast cancer risk is not well understood in Southeast Asian women. A comprehensive literature search of published articles from January 2001 to June 2019 (PubMed, ProQuest, and EBSCO) on the article databases and meta-analyses were performed to ask the duration of oral contraceptive application with risk of breast cancer among women in Southeast Asia. We divided the reference category being <5 years' duration and >5 years' duration of oral contraceptive application. We use the fixed and random-effect models to calculate odds ratios (OR), Egger's and Begg's test used to detect publication bias was presented with funnel plots. All analyzed data in Review Manager 5.3 (RevMan 5.3) and Stata version 16.2. A total of 388 studies were reviewed and 10 studies involving a total of 28,776 women were included in meta-analysis. This study found a slight increase in breast cancer risk in Southeast Asian women with oral contraceptive application >5 years with OR = 1.21 (95% CI 0.99-1.52, p=0.05). A higher risk of breast cancer was found in women with oral contraceptive application >5 years with OR = 2.69 (95% CI 1.70-3.94, p<0.0001).

Publication bias and heterogeneity were not found concerning a group of Southeast Asian women with oral contraceptive application for more than 5 years. Oral contraceptives use more than 5 years are at a higher risk of breast cancer among women in Southeast Asia. Although other reproductive factors including age at first childbirth, menarche, menopause, and abortion might influence the risk of breast cancer.

**KEY WORDS:** Breast cancer, Five years, Oral contraceptive, Southeast Asia

**Correspondence:** Richan Diana Nindirah, Doctoral Program, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia/ Department of Public Health and Community Medicine, Faculty of Medicine, Universitas Andalas, Padang, Indonesia.  
Email: richandiana78@gmail.com

Tel: + 62 271 832422  
Fax: + 62 71 545455

DOI: 10.55302/sr.v10i2.2719.2.22

© Adnanul Sholikhin, Nindirah. All rights reserved.

### INTRODUCTION

Breast cancer is the most frequent cancer and the second leading cause of cancer-associated mortality among women worldwide. A total of 2,088,849 new cases and 626,679 deaths have been related to breast cancer in 2018 [1]. Multifactorial etiology has been identified as one of the risk factors known is oral contraceptive application [2],[3].

Previous studies have revealed that combined oral contraceptives reduce endometrial and ovarian cancer risks through ovulation suppression [4],[5]. These studies found contrary results of contraceptive pills use with occurrence of breast cancer. An increased breast cancer risk correlated with used in oral contraceptive pills [4,5,6], meanwhile other studies have shown inconclusive results [7,8,9,10].

Other determinants of oral contraceptive use and the occurrence of breast cancer are the length of the use, dose-response, and age of the users. Another study showed a significant elevate in the number of young women who use oral hormonal contraceptive as well as women who begin to use oral contraceptives before the age of 20 [2]. Furthermore, the studies found there is no association between breast cancer and oral contraceptives use [2,4]. Increased breast cancer risk particularly in women with oral contraception application for more than 4 years before the first pregnancy [5]. Another study

reported breast cancer risk increasing in the oral contraceptives pills use for more than eight years [6].

Oral contraception has recently emerged as the third most common method for family planning and the most widely distributed across the globe compared to other contraceptive methods. It is estimated that 6.4% of Asian women use contraceptive pills [11]. Meanwhile, the proportion of oral contraceptive users in Southeast Asia is 12.8% [12]. Oral contraceptive users by country in Southeast Asia revealed that Thailand as the highest users (35%), followed by Indonesia (13.6%) and Vietnam (8.6%) [11,13]. Therefore, oral contraceptive application in Southeast Asia is relatively high yet the correlation with breast cancer risk is still under study.

### MATERIALS AND METHODS

#### Study design and research sample

In this meta-analysis study, we followed the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) Statement [14]. The samples of this research included published research articles which were published from January 2001 to June 2019 in PubMed, ProQuest, and EBSCO of online article databases. In each study we identified the reference category, being <5 years' duration and >5 years' duration of the application of oral contraceptive.

#### Operational definitions

This study comprised of independent variables, (a)  $\leq 5$  years' duration and (b)  $> 5$  duration of oral contraceptive use, and a dependent variable, i.e. breast cancer risk.

#### Research procedure

The procedure of this research was started by collecting data through published research articles identification on the oral contraceptive application length and the risk of breast cancer in Southeast Asian women on PubMed online article databases, ProQuest, and EBSCO (Figure 1).

The following keywords are used treating as title or abstract for the literature search: ("oral contraceptive" OR "hormonal oral") AND ("breast cancer" OR "breast neoplasms"). We found 385 articles were identified by examining the article titles, abstract and full-text. Furthermore, this study classified the research articles into two groups based on the oral contraceptive application length with  $\leq 5$  years' duration and  $> 5$  years' duration.

The article will be excluded if it is: (a) not breast cancer outcome, (b) conducted by using other study designs other than case-control or cohort study (c) included insufficient data for extraction.

#### Data collection technique

Online searching was performed in data collection. The collected data was limited by the articles written in English. The article type was limited to original research one. The publication date of the articles was limited from January 2000 to May 2019. The research subject was limited to humans only. Conceivably significant title unique articles were inspected, while the insignificant articles were prohibited. Those possibly significant unique articles will be evaluated in full-content structure, while then the unessential articles were excluded. The sample inclusion criterias were researched on the duration of oral contraceptive application, including  $\leq 5$  years' duration or  $> 5$  years' duration, and breast cancer in Southeast Asian women with retrospective study and prospective study design. The exclusion criterias were (a) the inclusion criterias were unsatisfyingly fulfilled, (b) the articles were unavailable in full-text form, and/or (c) the data provided in the full-text form was not sufficient for extraction of data. The following data were also collected from the articles: the name of the original writer, study location, study type, and a number of several samples.

The information from all of the articles that fulfilled inclusion criterias per under a standardized protocol carefully extracted by two investigator, while contradictions were settled by three different investigators. Newcastle Ottawa Quality Assessment Scale (NOS) was performed for evaluation quality of research articles. Articles were indicated poor (score 0-3), moderate (score 4-6) and high quality (score 7-9) [15].

#### Data analysis

Data analysis was conducted to get pooled and combined odd ratios of the collected articles. Odd ratio with 95% confidence intervals (CI) was utilized to pool the results. These test presented the minimum statistical power article with limited numbers and sample sizes was

indicated by  $I^2 > 50\%$  significantly heterogeneity. Assessment significant heterogeneity used a random-effect model and homogeneity used a fixed-effect model. Review Manager 5.3 (RevMan 5.3) was used to analyze the data. Publication bias was performed funnel plots and Egger's and Begg's tests, statistically significant publication bias p-value  $< 0.05$ . Stata version 14.2 was conducted to analyze publication bias. A two tailed P value of  $< 0.05$  was considered statistically significant.

#### RESULTS

Reviewing process was held to select 10 studies related to the duration of oral contraceptive application with breast cancer risk in Southeast Asian women with a total 28,776 samples patients (Table 1) [13,16-24].

A meta-analysis study was performed to determine the duration of oral contraceptive application and breast cancer risk in Southeast Asian women (Figure 2). The figure presented an insignificant result of  $\leq 5$  years' duration of oral contraceptive application and breast cancer risk in Southeast Asian women with OR = 1.21 (95% CI 0.96-1.52,  $p > 0.05$ ). The Heterogeneity test in these studies ( $P_{heterogeneity} = 0.0004$ ;  $I^2 = 70\%$ ) resulted in a variety of heterogeneous results on the breast cancer risk. In contrast, oral contraceptive application for  $> 5$  years and breast cancer risk in Southeast Asian women had a significant result with OR = 2.66 (95% CI 1.79-3.94,  $p < 0.00001$ ). Different from the  $\leq 5$  years' application duration, heterogeneity tests in these studies ( $P_{heterogeneity} = 0.14$ ;  $I^2 = 45\%$ ) resulted in multiformousness of homogeneous research on breast cancer risk.

Publication bias in articles collected described by presenting Funnel plots on the duration of oral contraceptive application with breast cancer risk among women in Southeast Asia (Figure 3). In figure 3, results showed as significant publication bias detected in articles on oral contraceptive  $\leq 5$  years' duration of oral contraceptive and risk of breast cancer among women in Southeast Asia with Egger's test ( $P = 0.011$ ) and Begg's test ( $P = 0.09$ ). In contrast, in articles on  $> 5$  years' duration of oral contraceptive use and risk of breast cancer among women in Southeast Asia (Figure 4), our study found that there was no significant publication bias with Egger's test ( $p = 0.270$ ) and Begg's test ( $p = 0.308$ ).

Subgroup analysis for oral contraceptive use and breast cancer risk in Southeast Asia (Table 2). Ten studies assessing oral contraceptive application and breast cancer risk, seven were case-control studies [13,16-18,21,22] with Pooled Odds Ratio (POR) 1.52 (1.06-2.18), heterogeneity test in these studies ( $P_{heterogeneity} = < 0.001$ ;  $I^2 = 90\%$ ) resulted in various heterogeneous research on breast cancer risk, and three were cohort studies [19,20] with POR 1.34 (0.98-1.84), heterogeneity test in these studies ( $P_{heterogeneity} = 0.28$ ;  $I^2 = 21\%$ ) resulted in a variety of homogeneous research on the breast cancer risk.

Study quality found eight studies with high NOS scores [13,18-24], POR 1.51 (1.09-2.10), heterogeneity test in these studies ( $P_{heterogeneity} = < 0.001$ ;  $I^2 = 88\%$ ) resulted various heterogeneous research on the breast cancer risk. Two studies with low NOS scores [16,17], POR 1.35

(0.80-2.27), heterogeneity test in these studies ( $I^2 = 73.1\%$ ;  $P < 0.001$ ;  $I^2 = 48\%$ ) resulted various homogeneous research. The oral contraceptive use duration and breast cancer risk in Southeast Asian countries was presented in Table 3. Our study found that there is an association of >5 years' duration of oral contraceptive application and breast cancer risk in Southeast Asian women, while <5 years' duration of oral contraceptives is not associated to breast cancer risk in Southeast Asian women.

Indonesia has the largest figure of women used oral contraceptives for >5 years with POR for breast cancer risk was 2.67 (1.62-4.40), followed by Thailand and Malaysia with POR for breast cancer risk were 2.56 (1.08-6.08) and 2.28 (0.84-6.16), respectively.

#### DISCUSSION

This study result showed that >5 years' length of oral contraceptive pills use increased breast cancer risk in Southeast Asian women with risk opportunity was more than doubled. The results were also homogeneous which means the studies will give consistent or insignificantly different if the study performs at the same time and place.

Furthermore, Indonesia has been known that this country has the greatest risk opportunities for breast cancer, followed by Thailand and Malaysia. Our study findings described possibly relate to the high rates of oral contraceptive application in those country mentioned in advance [11,13,24].

The previous studies found breast cancer incidence related to oral contraceptive application in several ethnicities. Non-Hispanic Caucasians have the highest percentage of oral contraceptive application (55%) as well as the highest number of incidence of breast cancer (6%). Interestingly, the lowest proportion of oral contraception application (37.5%) yet being the second-highest in the percentage of breast cancer incidence (5.7%) in non-Hispanic/Pacific Islanders. Non-Hispanic African-Americans and Hispanics have the same percentage of oral contraceptive application (52.2%) with a relatively lower figure of breast cancer incidence with 3.8% and 4.7%, respectively [25]. Our study had found the percentage of oral contraceptive application with breast cancer incidence followed by a range of 3.7-13.9%. The figure was higher in some ethnic than others as we had described, possibly due to oral contraception application percentage is known was also relatively high in the Southeast Asian region with 12.8% [12].

Based on the risk opportunities found, it was demonstrated that there was a higher breast cancer risk in Southeast Asian regions with < 5 years' duration of oral contraceptive application was OR = 1.21 (95% CI 0.96-1.52), while >5 years' duration of oral contraceptive use was OR = 2.66 (95% CI 1.79-3.94). Previous study also had found risk opportunities of breast cancer related to oral contraceptive included non-Hispanic Caucasians (HR = 1.09 (95% CI 1.01-1.18), non-Hispanic African Americans (HR = 0.95 (95% CI 0.63-1.42), and non-Hispanic Asian / Pacific Islander (HR = 0.93 (95% CI 0.63-1.39) [25].

Data from 39 case-control studies led from 1960 to 2006 have discovered a association of contraceptive pills with breast cancer among premenopausal [26]. Another study recommended that present utilization of contraceptive pills which substance is estrogen appears to somewhat elevate the breast cancer risk. A few factors adding to the advancement of breast disease incorporate hormones, which is estrogen itself [27-29]. The use of contraceptive pills which contains estrogen the breast tissue to be presented to large amounts of hormones for longer periods. That estrogen stimulates growth factors that exist in breast cancer cells resulting in tumor progression [30,31].

The previous research on the impact of age at beginning of utilization of the pill on the rate of breast cancer growth presumed that ladies that had begun utilizing the pill before 18 years old contracted disease 4 years sooner [32]. In a study, the researcher accepts prolonged use increases the risk. Another study found a relative risk of 2.2 with over 10 years of length of using [33]. The other study found the affinity between breast cancer and contraceptive pills use (OR=2.11). The investigation results show that women that utilization preventative pills have more noteworthy requirements for preventive and screening measures [34].

Based on data oral contraceptive is the most common means of contraception in the world and Southeast Asian countries, utilized by millions of women [35-37]. According to the duration of oral hormonal contraceptives use more than 5 years will increase breast cancer risk which will be higher compared to those who have never used hormonal contraception. If a person stops taking hormonal contraception for 5 years then she will have no risk of breast cancer. This result is supported by a theory found about the imbalance of the hormones estrogen and progesterone used in hormonal contraception [4,38,39]. This hormonal imbalance has resulted in a feedback mechanism that can physiologically control the number of hormones in the body when it is not functioning properly [40]. It was triggered by the estrogen receptors up regulation so that the number of hormones continues to increase. Exposure to sexual hormones over a period of more than 5 years will increase the proliferation of breast cells and increase breast cancer stem cell mitosis [41].

A meta-analysis data in this study has several limitations. First, there were 2 potential articles should be included in this study but the full-text articles are unavailable. Second, there were 2 studies are also potentially included in data analysis yet the data presented were incomplete and different risk factors in the calculations were also performed.

Based on our analysis, as we found of the significant association of oral hormonal contraceptives use with breast cancer risk as well as increase for more than five years' duration of contraceptive pills use, we suggest people to select effective and efficient long-term contraceptive methods to prevent pregnancy, yet safe in accordance with infertility and breast cancer risk. Another suggestion is breast cancer screening programs are necessary for Southeast Asian countries.

## CONCLUSION

This study confirms the significant correlation of more than five years' duration of oral hormonal contraceptives use and breast cancer risk in Southeast Asia. Therefore, we recommend fertile women who want to prevent pregnancy for a relatively long period, should use long-term contraception methods.

## ACKNOWLEDGMENTS

We express our gratitude to Mac Arit Hamdanas, MA for proofreading the manuscript draft.

## ABBREVIATIONS

CI: Confidence Intervals  
HR: Hazard Ratio  
NOS: Newcastle–Ottawa Quality Assessment Scale  
OR: Odds Ratios  
POR: Pooled Odds Ratios  
PRISMA: Preferred Reporting Items for Systematic Reviews and Meta Analysis  
RevMan: Review Manager

## REFERENCES

1. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for Clinicians* [Internet]. Wiley; 2018 Sep 12;68(6):394–424. Available from: <http://dx.doi.org/10.3322/caac.21492> Feedback: <https://doi.org/10.1111/1365-3113.12525>
2. Breast cancer and hormonal contraceptives: collaborative reanalysis of individual data on 53 297 women with breast cancer and 100 239 women without breast cancer from 51 epidemiological studies. *The Lancet* [Internet]. Elsevier BV; 1996 Jun;347(9017):1713–27. Available from: [https://doi.org/10.1016/S0140-6736\(96\)03100-0](https://doi.org/10.1016/S0140-6736(96)03100-0)
3. Barnard ME, Boeke CJ, Tamimi RM. Established breast cancer risk factors and risk of intrinsic tumor subtypes. *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer* [Internet]. Elsevier BV; 2015 Aug;186(1):73–85. Available from: <https://dx.doi.org/10.1016/j.bbcan.2015.06.002>
4. Haile RW, Thomas DC, McGuire V, Fellberg A, John EM, Milne RL, et al. BRCA1 and BRCA2 Mutation Carriers, Oral Contraceptive Use, and Breast Cancer Before Age 50. *Cancer Epidemiology Biomarkers & Prevention* [Internet]. American Association for Cancer Research (AACR); 2006 Oct 1;15(10):1863–70. Available from: <http://dx.doi.org/10.1158/1078-2967.ept1051478>
5. Urban M, Banks E, Egger S, Canfell K, O'Connell D, Beral V, et al. Injectable and Oral Contraceptive Use and Cancers of the Breast, Cervix, Ovary, and Endometrium in Black South African Women: Case-Control Study. Franco LL, editor. *PLoS Medicine* [Internet]. Public Library of Science (PLoS); 2012 Mar 6;9(3):e1001182. Available from: <https://doi.org/10.1371/journal.pmed.1001182>
6. Hunter DJ, Colditz GA, Hankinson SE, Malspeis S, Spiegelman D, Chen W, et al. Oral Contraceptive Use and Breast Cancer: A Prospective Study of Young Women. *Cancer Epidemiology Biomarkers & Prevention* [Internet]. American Association for Cancer Research (AACR); 2010 Aug 27;19(10):2496–502. Available from: <https://doi.org/10.1158/1078-2967.ept1051478>
7. Marchbanks PA, McDonald JA, Wilson HG, Folger SG, Mandel MG, Daling JR, et al. Oral Contraceptives and the Risk of Breast Cancer. *New England Journal of Medicine* [Internet]. Massachusetts Medical Society; 2002 Jun 27;346(26):2025–32. Available from: <https://doi.org/10.1056/NEJMoa012707>
8. Belhica TN, Rosenberg L, Hong C-C, Troester MA, Lunetta KL, Bandera EV, et al. A case-control analysis of oral contraceptive use and breast cancer subtypes in the African American Breast Cancer Epidemiology and Risk Consortium. *Breast Cancer Research* [Internet]. Springer Science and Business Media LLC; 2015 Feb 21;17(1). Available from: <https://doi.org/10.1186/s12874-015-0172-5>
9. Veisy A, Lotfnejad S, Salehi K, Zhiban F. Risk of Breast Cancer in Relation to Reproductive Factors in North west of Iran, 2013–2014. *Asian Pacific Journal of Cancer Prevention* [Internet]. Asian Pacific Organization for Cancer Prevention; 2015 Feb 25;16(2):151–5. Available from: <https://doi.org/10.7324/apjcp.2015.162151>
10. Zare N, Haem E, Loukarami KB, Heydari SJ, Barooti E. Breast Cancer Risk factors in a Defined Population: Weighted Logistic Regression Approach for Rare Events. *Journal of Breast Cancer* [Internet]. Korean Breast Cancer Society (KAMJEB); 2013;16(2):214. Available from: <http://dx.doi.org/10.4048/jbc.2013.16.2.214>
11. Mansour D, Gemzell Danielsson K, Inki P, Tensen JT. Fertility after discontinuation of contraception: a comprehensive review of the literature. *Contraception* [Internet]. Elsevier BV; 2011 Nov;84(5):165–77. Available from: <https://doi.org/10.1016/j.contraception.2011.07.007>
12. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. *Combined Estrogen-progestogen Contraceptives and combined estrogen-progestogen menopausal therapy*. IARC Monogr Eval Carcinog Risks Hum 2007;91:1–528.
13. Nguyen J, Le QH, Duong BH, Sun P, Pham HT, Ta VT, et al. A Matched Case-Control Study of Risk Factors for Breast Cancer Risk in Vietnam. *International Journal of Breast Cancer* [Internet]. Hindawi Limited; 2016;2016:1–7. Available from: <https://doi.org/10.1155/2016/1612312>
14. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JPA, et al. The PRISMA Statement for Reporting Systematic Reviews and Meta Analyses of Studies That Evaluate Health Care Interventions: Explanation and Elaboration. *PLoS Medicine* [Internet]. Public Library of Science (PLoS); 2009 Jul 21;6(7):e1000100. Available from: <http://dx.doi.org/10.1371/journal.pmed.1000100>
15. Wells GA, Shea B, O'Connell D, Peterson I, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of non randomised studies in meta-analysis; 2009 [cited 2018 June 10]. Available from: [http://www.chcrl.ca/files/resources/publications/assessment\\_tools/nos/](http://www.chcrl.ca/files/resources/publications/assessment_tools/nos/)

39. Dall GV, Britt KL. Estrogen Effects on the Mammary Gland in Early and Late Life and Breast Cancer Risk. *Frontiers in Oncology* [Internet]. Frontiers Media SA; 2017 May 26;7. Available from: <http://dx.doi.org/10.3389/fonc.2017.00110>
40. Colditz GA, Hohlke K, Beckey CS. Breast cancer risk accumulation starts early: prevention must also. *Breast Cancer Research and Treatment* [Internet]. Springer Science and Business Media LLC; 2014 May 13;143(2):567-79. Available from: <http://dx.doi.org/10.1007/s10548-014-0299-0>
41. Nindrea RD, Aryandono T, Lazuardi L, Dwipradasto I. Association of Dietary Intake Ratio of n-3/n-6 Polyunsaturated Fatty Acids with Breast Cancer Risk in Western and Asian Countries: A Meta-Analysis. *Asian Pacific Journal of Cancer Prevention* [Internet]. EpiSmart Science Vector Ltd; 2019 May 1;20(5):1321-7. Available from: <http://dx.doi.org/10.31838/apjcp.2019.2005.1321-1327>

Table 1: Systematic review of duration of oral contraceptive use with breast cancer risk in Southeast Asian women

First Author	Region	Study Type	Patients Characteristic	Duration (years)	Number of Sample				NOS
					Cases		Control		
					Total	Exposure	Total	Exposure	
Norsaslah et al [16]	Malaysia	Case control	Aged 26-70 years	≤5	147	53	147	36	6
Gibson et al [17]	Philippines	Case control	Aged 35-64 years	≤5	123	16	978	125	6
Matalqah et al (a) [18]	Malaysia	Case control	Aged 25-83 years, three ethnic groups (Malay, Chinese and Indian)	<5	150	27	150	21	7
Matalqah et al (b) [18]	Malaysia	Case control	Aged 23-83 years, three ethnic groups (Malay, Chinese and Indian)	>5	150	13	150	6	7
Pousari et al (a) [19]	Thailand	Cohort	Aged 30-69 years	<5	3,664	14	2,153	11	8
Pousari et al (b) [19]	Thailand	Cohort	Aged 30-69 years	>5	5,397	45	2,153	11	8
Lee et al [20]	Singapore	Cohort	Aged 45-74 years, Chinese population	<5	411	39	1,213	81	7
Nguyen et al [13]	Vietnam	Case control	Aged 25-75 years	≤5	294	25	294	13	7
Trien et al [21]	Vietnam	Case control	Aged 27-74 years	≤5	269	79	519	134	7
Chaveepojunkomjorn et al (a) [22]	Thailand	Case control	Aged <45 years	≤5	257	42	257	41	7
Chaveepojunkomjorn et al (b) [22]	Thailand	Case control	Aged <45 years	>5	257	125	257	51	7
Tan et al [23]	Malaysia	Case control	Aged 40-74 years, three ethnic groups (Malay, Chinese and Indian)	<5	3,683	935	3,980	1,145	8
Wahidi et al (a) [24]	Indonesia	Case control	Aged 40-49 years	≤5	381	77	381	53	7
Wahidi et al (b) [24]	Indonesia	Case control	Aged 40-49 years	>5	381	58	381	21	7
<b>Total</b>					<b>15,764</b>	<b>1,548</b>	<b>13,012</b>	<b>1,755</b>	

Abbreviation: NOS, Newcastle–Ottawa Quality Assessment Scale

Table 2: Subgroup analysis for oral contraceptive use and breast cancer risk in Southeast Asia

Subgroups	Number of studies	Pooled OR (95% CI)	Heterogeneity	
			I <sup>2</sup> (%)	p
<b>Study design</b>				
Cohort	3	1.34 (0.98-1.84)	21	0.28
Case control	7	1.52 (1.06-2.18)	90	<0.001
<b>Study quality</b>				
High (NOS scores ≥7)	8	1.51 (1.09-2.10)	88	<0.001
Low (NOS scores <7)	2	1.33 (0.80-2.27)	18	0.17

Abbreviation: CI, confidence interval; OR, odds ratio; p < 0.05 considered statistically significant; significant heterogeneity >=0%

Figure 13: Funnel plots for <5 years' duration of oral contraceptive use and breast cancer risk in Southeast Asian women







Figure 14. Funnel plots of >5 years' duration of oral contraceptive use and breast cancer risk in Southeast Asian women



# Oral Contraceptive Used More than 5 Years is Associated with Increased Risk of Breast

## ORIGINALITY REPORT

14%

SIMILARITY INDEX

14%

INTERNET SOURCES

9%

PUBLICATIONS

14%

STUDENT PAPERS

## MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

15%

★ Submitted to Udayana University

Student Paper

Exclude quotes

Exclude bibliography

Exclude matches < 3%