

**Proceeding  
11<sup>th</sup> FDI IDA  
Continuing Dental Education 2015**

**Bumi Minang Hotel, Padang, Indonesia  
14-15 November 2015**

*LSKI*

ISBN 602180828-2



9 786021 808283

PROCEEDING  
11<sup>th</sup> FDI-IDA CONTINUING DENTAL EDUCATION 2015

Editor :

Dhona Afriza  
Rifani  
Abu Bakar  
Okmes Fadriyanti

Bumi Minang Hotel, Padang November 14<sup>th</sup>-15<sup>th</sup> 2015

**PROCEEDING  
11<sup>th</sup> FDI-IDA CONTINUING DENTAL EDUCATION  
PADANG**

Proceeding 11th Fdi Continuing Dental Education  
Publisher by Lembaga Studi Kesehatan Indonesia (LSKI)

Bandung, November 2015

**Editor** Dhona Afriza  
Rifani  
Abu Bakar  
Okmes Fadriyanti

**Setting** Siti Mariam  
**Production** Agus Sono  
**Printed by** Sono Offset  
**Copyright** © 2015 11th FDI Continuing Dental Education Committee  
**ISBN** 978 602 18082 8 3

Perpustakaan Nasional: Katalog Dalam Terbitan (KDT)

**PROCEEDING  
11<sup>th</sup> FDI-IDA CONTINUING DENTAL EDUCATION 2015/Editor: Dhona  
Afriza, Rifani, Abu Bakar, Okmes Fadriyanti**

-- Bandung : LSKI (Lembaga Studi Kesehatan Indonesia) 2015 viii +  
214 hlm; 21 cm

ISBN 978 602 18082 8 3

## CONTENT

- Solution of cancer of the mouth cavity et protein conformational disorder which it is prefixed by the presence of an exaggerated expression heat shock proteins by a virus-infected cell** 1-9  
Adi Prayitno
- Analysis of the difference of caries occurrence between resin based and glass ionomer based pit and fissure sealant on young permanent teeth oral presentation** 10-16  
Nila Kasuma, Aida Fitriana
- Oral lesions associated with systemic lupus erythematosus mimicking oral lichen planus oral presentation (case report)** 17-25  
Fitria Mailiza, Irma Sufiawati
- Acyclovir as prophylactic antiviral therapy in recurrent herpes simplex infection** 26-33  
Indra Gunawan, Irma Sufiawati
- The Effect of CPP-ACP Application on Surface Hardness and Roughness of Tooth Enamel after Immersed in Carbonated Drinks** 34-39  
Ega Sofiana, Sulistiyani, Nadie Fatimatuzzahro
- Oral conditions in patients with major thalassemia In thalassemia centre at cipto mangunkusumo hospital** 40-46  
Surya Nelis, Harum Sasanti
- The effectiveness of longan (euphoria longana) seed extract against bacterial growth of streptococcus  $\alpha$ -haemolyticus Post tooth extraction** 47-53  
Elsya Octavianti, Efa Ismardianita, Resa Ferdina
- Antibacterial effect of 80% mahkota dewa (phaleria macrocarpa. Scheff [boerl]) extract in root canal irrigation** 54-59  
Syahrul Fatah Maulady, Siti Rusdiana Puspa Dewi, Rini Bikarindrasari
- Cutting section of putty impression matrices for optimum tooth preparation index Oral presentation (literature review)-proceeding fdi padang 2015** 60-66  
Veni Takarini, Zulia Hasratiningsih
- Immediate Dental Implant Placement Post Radicular Cyst Enucleation with an Application of Alloplastic Bone Graft and L-PRF Membrane** 67-76  
Setiawan

<b>Management of oral candidiasis</b>	<b>77-81</b>
IGN. Putra Dermawan	
<b>The correlation between smoking habit and blood pressure towards pre-clinic dentistry students of prof. Dr. Moestopo (beragama) university jakarta</b>	<b>82-92</b>
Retno Damayanti, Rizfira Ananda	
<b>The effectivity of platelet rich plasma (prp) gel on blood clot formation process After tooth extraction on young balinese dog</b>	<b>93-98</b>
Putu Sulistiawati Dewi	
<b>The application of color in dental room treatment to reduce patient anxiety</b>	<b>99-107</b>
Marta Juslily	
<b>The effect of trauma from occlusion on periodontal tissue</b>	<b>108-115</b>
Abdul Gani Soulisa	
<b>Allergy to dental implant as a reason for implant failure</b>	<b>116-121</b>
Durra Mufida	
<b>Salivary flow rate and total protein concentration in asthmatics, treated with corticosteroids and sympathomimetics drugs</b>	<b>122-126</b>
Nur Isya Sagita, Shanty Chairani, Sulistiawati	
<b>Esthetic complex treatment, an interdisciplinary of periodontology and dentalconservative</b>	<b>127-132</b>
Risya Cilmiaty, Anton Kusumo Widagdo, Adisty Restu Poetri	
<b>Calsium hydroxide as intracanal medication for non surgical endodontic treatment on a maxillary right lateral incisor with periapical lesion</b>	<b>133-140</b>
Deli Mona	
<b>The difference between saliva ph and blood glucose levels before and after consuming white rice (oriza sativa) and casava (manihot esculenta crantz)</b>	<b>141-151</b>
Fahmi Hamjah Siregar, DwiPrijatmoko, Sulistiyani	
<b>The ph effect toward the micro structure and surface hardness of hybrid composite</b>	<b>152-159</b>
Elin Karlina, Zulia Hasratiningsih, RennyFebrida	

Keyword : virus, PCDo, Misfolding, Hsp70, Herb

<b>Prosthetic rehabilitation post enucleation in patients with eye injury due use ocular prosthesis to increase aesthetic</b>	<b>160-167</b>
Nining Handayani	
<b>Paradigm in Esthetic Dental Restoration</b>	
<b>Oral Presentation (Literature Review)</b>	<b>168-173</b>
Zulia Hasratiningsih, Veni Takarini	
<b>Lip line position on three different facial types of young adult population in Makassar</b>	<b>174-179</b>
Susilowati, Rika Damayanti, Nur Infaq Ridal	
<b>The Importance of good 'dental teamwork' in supporting successful dental practice</b>	<b>180-186</b>
Mita Juliawati	
<b>Comparison of effectiveness between poster media and flipchart media for deaf student of sdlb-b yptb malang knowledges in the context of oral health promotion</b>	<b>187-193</b>
Trining Widodorini, Neny Roeswahjuni, Ambar Puspitasari, Tiazani Sholichah	
<b>Extracellular matrix types and distribution of pattern differentiation of ameloblastoma based on histopathological type appearance</b>	<b>194-201</b>
Galang Rikung Edy Santoso, Mei Syafriadi, Dwi Merry Christmarini Robin	
<b>The analysis of composite resin color change after cigarette smoke exposure</b>	<b>202-207</b>
Diana Setya Ningsih, Viona Diansari, Maya Rizki Mauliani	
<b>Analysis of il-1<math>\beta</math> levels in male wistar rats induced by porphyromonas gingivalis with intervention of dadiah isolate bacteria pediococcus pentosaceus</b>	<b>208-214</b>
Aida Fitriana, Nila Kasuma	

# Analysis of the difference of caries occurrence between resin based and glass ionomer based pit and fissure sealant on young permanent teeth oral presentation

Nila Kasuma, Aida Fitriana

Faculty of Dentistry, Andalas University Indonesia

## ABSTRACT

**Background :** Caries is a major problem in dentistry. One of caries preventive method is pit and fissure sealant. Pit and fissure sealant is effective in area with narrow anatomical shape which facilitate plaque and bacterial growth. Materials used for this action may take the form of resin based and glass ionomer cement. **Objective :** To compare the white spot index (ICDAS-II) after application of resin based and glass ionomer fissure sealant and see the types of materials that are more effective in the long term **Method :** Design of this study was experimental pre-test and post-test. Population were students of class I, II, and III in the Elementary School 02, Cupak Tengah, Padang. Samples were taken by using purposive sampling. There are two groups of samples, each consisted of 30 children given application resin based sealant and glass ionomer. After a month post application, white spot index were measured with ICDAS-II index. Data were analyzed with SPSS 17 using unpaired t-test. **Result :** There is no significant difference between applications resin based sealant and glass ionomer cements after 1 month ( $p= 0,45$ ). **Conclusion :** Resin based sealant and glass ionomer cement are effective pit and fissure sealant material

**Keyword :** pit and fissure sealant , resin based, glass ionomer, white spot index

## INTRODUCTION

Caries and periodontal disease is the oral cavity disease which ranked highest in Indonesia. According to Household Health Survey (SKRT), caries prevalence in 2004 reached 90.05%. According to Basic Health Research (RISKESDAS) in 2013, DMF-T index (decayed, missing and filled teeth) reached 4.6, for West Sumatra province were 4,7. DM-T increases as age. At the age of 12 years DMFT index is 1.4, while at the age of 15 years reaches 1.5, and at the age of 18 years is 1.6. Data of Padang based RISKESDAS 2007 showed that caries is the first of ten most common diseases suffered with prevalence 52%. Recent data showed prevalence in 2013 reached 56%.

White spot lesions are the initial clinical manifestations of caries. White spot is the damage of enamel surfaces. White spot on the enamel surface appear in 4 weeks. Changes



from white spot into caries happen in 6 months.<sup>1</sup> The incidence of white spot into caries in permanent dentition is 59.1% , while in primary dentition is 71.7%.<sup>2</sup>

Caries can damage the body health either locally or systemically. In children, caries incidence is due to mother's lack of knowledge about mixed dentition phase and mechanical capabilities of immature children to clean their teeth. Caries affects chewing food and aggravates the digestive system, so the nutrients that are needed by the body for growth will not be completely absorbed. Caries in the oral cavity will increase the number of bacteria retention and affect the body systemically through the bloodstream causing vital organs infection e.g heart. Efforts should be made to reduce the prevalence of caries include promotive, preventive, curative and rehabilitative . Curative and rehabilitative efforts are generally carried out in the health service, but in reality, communities who received treatment are only 31.1%.

Prevention of dental caries can be done through fluoridation of drinking water, application of topical fluoride on enamel development phase, and the plaque control program for each individual. It is not proven effective in reducing the incidence of caries in pits and fissures that are the most vulnerable parts of caries, because of its anatomical narrow shape.<sup>3</sup>

Administering topical and systemical fluoride does not cause much effect on reducing the incidence of caries pit and fissure because of pit and fissure is a protected basin area. Condition of pits and fissures is the influencing factor of caries development process. Preventive way to prevent pit and fissure caries is called pit and fissure sealant.<sup>4</sup> Giving pit and fissure sealant material at the beginning of tooth eruption is expected to prevent bacteria colonization in pit and fissure.<sup>5</sup>

The aim of sealant is the penetration of ingredients into the pits and fissures and polymerizes and sealed off the area of bacteria and debris.<sup>2</sup> Sealant material must have a good retention capability and durability, low solubility of the oral fluid, biocompatible with the tissue cavity mouth, and easy to apply.<sup>6</sup>

Indication of pit and fissure sealant is a young new permanent tooth eruption around 1-3 years in the age range of 6-9 years, free of caries and have anatomical structure with deep pits and fissures. Molar teeth of the most often damaged even loss occurs at an early age. The teeth are erupted permanent teeth first so often ordinary people pay less attention because they are still the primary teeth, also first molar is occlusion key .

Two sealant material that is often used is resin-based sealant and glass ionomer cement based sealant (GIC). Resin-based sealant materials polymerize through autopolymerization and photopolymerization. While glass ionomer cement usually autopolymerized.<sup>5</sup>

Resin-based sealants last longer and stronger because it has the ability to penetrate better. This is because the process of etching the tooth enamel which results in better contact between the surface of the resin material with enamel.<sup>7</sup> Etching eliminates mineral tooth enamel and produces resin tags and clinically appears whiter and fade. Sealant material given in the etched area will penetrate into the resin tag. This can improve the mechanical retention of sealant material to the surface of the tooth enamel.<sup>8</sup>

Glass ionomer sealants have the ability to prevent caries as well as the resin-based sealant. Glass ionomer cement sealant is easier to manipulated because it doesn't require stages of etching on the surface of the tooth.<sup>9</sup> Glass ionomer cement sealant materials perform specific interactions with tooth enamel by releasing calcium, strontium and fluorine ions that are cariostatic also reduce the caries development in areas where sealant were given.<sup>10</sup>

### **objectives**

The authors are interested to compare the caries index in the application of resin-based fissure sealant with sealant glass ionomer cements as pit and fissure sealing material on the surface of the young permanent teeth for a month usage

### **METHODS**

Design of this study was an experimental pre-test and post-test. The study population were students of class I, II, and III in the Elementary School 02, Cupak Tengah, Padang. Samples were taken by using purposive sample has met the inclusion criteria, first molars maximum of 4 years after the eruption; free of caries; deep pit and fissure; ages 6-9 years. Sample exclusion criteria are students are not at school while doing research; there is a congenital disease; students are not willing to be a subject of research. There are two groups of samples. The first group consisted of 30 children given application-based pit and fissure sealant resin. The second group consisted of 30 children given a pit and fissure sealant applications based glass ionomer. Before the application of pit and fissure sealants, teeth cleaned with a brush and pumice. Teeth were isolated with cotton rolls, dry surface for 20-30 seconds.

In the first group, the tooth surfaces were etched, rinsed for 60 seconds, then dried with air. Check the etching with air, etched surface will appear whiter. Dry with air for 20 seconds. Sealant material is applied, polymerization with light curing. Check occlusion with articulating paper.

In the second group, dentin conditioner is applied for 10-20 seconds to remove plaque and pellicle and prepare cement to adapt well toward the tooth surface and provide good adhesion. Tooth is rinsed with water for 60 seconds. Drying with air conditioner dentin surface after application of pit and fissure done flushing for 20-30 seconds. GIC applications in pit and fissure then apply varnish. Check occlusion with articulating paper, if there is excessive contact, carried out spot grinding.

After a month post pit and fissure sealant application, white spot index were measured with ICDAS-II index. Score of ICDAS-II :

- 0 = sound tooth surface, no evidence of caries after 5 seconds air drying
- 1 = first visual change in enamel : Opacity or discoloration (white or brown) is visible at the entrance to the pit or fissure seen after prolonged air drying
- 2 = Distinc visual change in enamel visible when wet , lesion must be visible when dry

- 3 = localized enamel breakdown (without clinical visual signs of dentinal involvement) seen when wet and after prolonged drying
- 4 = underlying dark shadow from dentine
- 5 = distinct cavity with visible dentine
- 6 = extensive (more than half the surface) distinct cavity with visible dentine

Data were analyzed with SPSS 17 unpaired t-test to analyze comparative pit applications and fissure sealant resin with glass ionomer-based.

## RESULT

Subject of the study consisted of 63% male, average age  $14.18 \pm 7.15$  and 27% female, average age of  $8.20 \pm 5.91$ . The number of samples is 30 samples were applied based pit and fissure sealant resin and 30 samples of pit and fissure sealant were applied glass ionomer-based glass.

Table 1. The average age of the subjects by gender

Age	Gender	n	%	Mean
6-9 years old	Male	38	63	7,15±14,8
	Female	22	27	8,20±5,91

From the collected data, ICDAS-II index was measured based white spot on usage-based fissure sealants with glass ionomer resin after 1 month of use.

Table 2. Index-II ICDAS white spot on usage-based fissure sealants with glass ionomer resin after 1 month of use.

Variable	n	Mean ±SD
F.S. Resin	30	0,00±0,00
F.S. Glass Ionomer Cement	30	0,00±0,00

Further analysis test conducted by using test t to see the difference between the usage-based fissure sealants with glass ionomer resin after 1 month of use.

Table 3. Results of statistical unpaired T-Test of ICDAS-II

Variable	n	Mean ±SD	p- value
F.S. Resin	30	0,00±0,00	0,45*
F.S. Glass Ionomer Cement	30	0,00±0,00	

significance  $p < 0,05$

Based on statistical test results as shown in Table 3, the value  $p = 0.45$  where  $p < 0.05$ , which means there is no significant difference ICDAS-II between the use of 1 month fissure sealants and glass ionomer-based resin.

## DISCUSSION

Research conducted on elementary school students Cupak Tengah 02 on the age of 6-9 years old. Thirty students were applied resin-based pit and fissure sealant and thirty students were applied glass ionomer cement pit and fissure sealant.

Based on the results of research on the index of caries in children with fissure sealant and glass ionomer-based resin-based fissure sealant, it is known that there is no significant differences between the glass ionomer-based fissure sealants and fissure sealant based resin with value of  $p = 0,45$  ( $p < 0,05$ ).

The occlusal surfaces of posterior teeth are the areas most prone to the occurrence of caries. Anatomical form allows plaque retention and maturation. Bacteria activity causes pH fluctuations. Conditions of pH rising causes teeth remineralization, while the drop in pH will result in the loss of tooth mineral. Loss of these minerals is a process of demineralization of hard tissue that becomes the signs and symptoms of caries.<sup>5</sup>

Early symptoms of an enamel caries can be seen macroscopically in the form of white patches. These patches have colors that appear very different from the surrounding healthy enamel. Sometimes it would appear brown lesions caused by the material around that is absorbed into the pores. Both patches of white or brown spots can last annual duration.<sup>11</sup> The damage begins with an overview in the form of white lesions on the surface of the tooth enamel which is called white Spot lesions, measured using ICDAS-II (international caries detection and assessment system). The occurrence of white spot on the enamel surface for 4 weeks. Changes into a white spot caries will occur for 6 months.<sup>1</sup> After the enamel were exposed with caries, it takes about 3-4 years to grow until reaching the dentine caries. The development of clinically detectable caries depending on the thickness of the enamel loss and morphological pit and fissure formation.<sup>12</sup>

Pits and fissures of posterior teeth are considered to be highly susceptible to the adhesion of micro-organisms and consequently, to caries. Therefore, a significant amount of tooth decay occurs at these sites. Fissure sealants are used to prevent occlusal caries, 71% percent of occlusal decay being preventable after a once-off fissure sealant application.<sup>13</sup> The most commonly used sealant material is resin composite.<sup>14</sup> Its caries-preventive effect relies on the sealing of pits and fissures through micro-retention, created through tags after enamel acid etching. However, these are easily destroyed by saliva contamination, which reduces micro-retention and, consequently, the caries-preventive effect.<sup>15</sup>

Under the generally wet conditions in the oral cavity, Glass Ionomer Cement (GIC) offers an alternative. Owing to its hydrophilic properties, GIC is not as moisture-sensitive as hydrophobic resin.<sup>15</sup> Many researchers confirm that the Glass ionomers are separately be preferable for sealing newly erupted teeth. Glass ionomer sealants offer similar caries-

preventive effects as resin-based sealants, with easier manipulation and without the use of acid etching. The glass ionomer may be valuable as a sealant in cases of difficult operating conditions i.e. difficulty with moisture control in partially erupted teeth or in children with management problems or in very young children.<sup>9</sup>

Within the limitation of this study, it is concluded that the ICDAS-II index is not different between application of resin based and glass ionomer cement pit and fissure sealant. Each material has the different effectiveness over 1 months. Etching roughens the tooth surface and produces a honeycomb-like structure so that tags of sealant can penetrate deeply into the enamel and form an effective mechanical bond, thus retaining the sealant. While glass ionomer cement could be a good alternative when and where resources for resin sealant placement are not readily available, and difficult to isolate the teeth due to hypersalivation.


## CONCLUSION

There is no significance different of caries index ICDAS-II between application of resin based and glass ionomer cement pit and fissure sealant for 1 month usage. Both of materials are effective as pit and fissure sealant

## REFERENCES

1. Livas C, Kuijpers-Jagtman AM, Bronkhorst E, Derks A, Katsaros C. Quantification of white spot lesions around orthodontic brackets with image analysis. *Angle Orthod* 2008 ;78(4):585-90.
2. Anusavice KJ. Ilmu Bahan Kedokteran Gigi 2004. Jakarta: EGC.
3. Craig, Robert G. Dental Materials. London: Mosby Co. 1979.
4. Andlaw, RJ, Rock. 1992. Perawatan Gigi Anak. Alih bahasa: Agus Djaya dari A Manual of Pedodontics. Jakarta: EGC
5. Kervanto, Sari. 2009. Arresting Occlusal Enamel Caries Lesions with Pit and Fissure Sealants. Academic Dissertation Faculty of Medicine, University of Helsinki. Diakses dari <https://oa.doria.fi/bitstream/handle/10024/43707/arrestin.pdf?sequence=1> pada 8 Juni 2009
6. Lesser, Donna, RDH, BS. 2001. An Overview of Dental Sealants. [http://www.adha.org/downloads/sup\\_sealant.pdf](http://www.adha.org/downloads/sup_sealant.pdf) ; June 8th 2009
7. Ganesh, Mahadevan MDS, et al. 2007. Comparative Evaluation of The Marginal Sealing Ability of Fuji VII and Concise as Pit and Fissure Sealants. *The Journal Contemporary Dental Practice*, diakses dari <http://www.thejcdp.com/issue033/ganesh/ganesh.pdf> pada 8 Juni 2009.
8. Paarmann, Carline, RDH, MEd. 1991. Application of Pit and Fissure Sealants. [http://www.ptc.idaho.gov/Forms\\_Publications/Health/Curriculum/Dental Application Of Pit And Fissure Sealants.pdf](http://www.ptc.idaho.gov/Forms_Publications/Health/Curriculum/Dental_Application_Of_Pit_And_Fissure_Sealants.pdf) ; June 6th 2009.
9. Subramaniam P. 2008. Retention of Resin Based Sealant and Glass Ionomer used

- as a Fissure Sealant: a Comparative Study. *Jurnal Indian Soc. Pedodontics Prevent Departemen*. [http://www.jisppd.com/temp/JIndianSocPedodPrevDent263114-3280171\\_090641.pdf](http://www.jisppd.com/temp/JIndianSocPedodPrevDent263114-3280171_090641.pdf); June 8th 2009
10. Walsh, Laurence J, Prof. 2006. Pit and Fissure Sealant: Current Evidence and Concepts. *Dental Practice Journal*. [https://espace.library.uq.edu.au/eserv/UQ:13804/Sealants\\_2006.pdf](https://espace.library.uq.edu.au/eserv/UQ:13804/Sealants_2006.pdf). June 8th 2009
  11. Kidd, Edwina AM and Bechal, Sally Joyston. 1992. *Dasar-Dasar Karies Penyakit dan Penanggulangannya*. Terjemahan Narlan Sumawinata dan Safrida Faruk dari *Essential of Dental Caries* (1992). Jakarta: EGC
  12. Pinkham, J.R. 1994. *Pediatric Dentistry, Infancy Trough Adolescence* second edition. Philadelphia: W.B Saunders Co Locker D, Jokovic A, Kay EJ.
  13. Prevention. Part 8: The use of pit and fissure sealants in preventing caries in the permanent dentition of children. *Br Dent J* 2003 11;195(7):375-8.
  14. Adair SM. The role of sealants in caries prevention programs. *J Calif Dent Assoc*. 2003;31:221-227.
  15. Bishara SE, Oonsombat C, Ajlouni R, Denehy G. The effect of saliva contamination on shear bond strength of orthodontic brackets when using a self-etch primer. *Angle Orthod*. 2002;72:554-557.
  16. Mickenautsch S, Yengopal V. Caries-preventive effect of glass ionomer and resin-based fissure sealants on permanent teeth: An update of systematic review evidence. *BMC Research Notes*. 2011;4:22.



# Certificate

11<sup>th</sup>

**11<sup>th</sup> FDI-IDA CONTINUING DENTAL EDUCATION 2015**

**"DENTISTRY TOWARDS GLOBALIZATION: THE NEW CHALLENGES TO IMPROVE  
COMMUNITY AWARENESS OF ORAL HEALTH SERVICES"**

Presented to

.....  
**NILIA KRASUMA**  
.....

as

**SPEAKER**

**IN THE 11<sup>TH</sup> FDI-IDA CONTINUING DENTAL EDUCATION 2015**

**NOVEMBER 14 - 15<sup>TH</sup> 2015**

**Bumi Minang Hotel - Padang**

**drg. Farichah Hanum, M.Kes**  
President of IDA

**drg. Syffianti Syaharman, MMKes**  
Chairman of Organizing Committee

Accredited by : Indonesian Dental Association (IDA)

No : SKP-1/401rev/PB PDGI/X/2015

Participant	:	6,5	SKP
Invited Speaker (1-2 hour)	:	4	SKP
Speaker (< 1 hour)	:	3	SKP
Presentation Jury	:	5	SKP
Poster Jury	:	5	SKP
Poster Presenter	:	4	SKP
Moderator	:	3	SKP
Committee	:	3	SKP
Hands on/Workshop participant	:	3,5	SKP
Hands on/Workshop Instructor	:	5	SKP