

Turnitin Originality Report

- Processed on: 30-Jan-2020 9:23 PM +08
- ID: 1248708310
- Word Count: 2400
- Submitted: 1

Isolation, Characterization and Production of Phytase from Endophytic Fungus its Application for Feed

Similarity Index

29%

Similarity by Source

Internet Sources:

N/A

Publications:

29%

Student Papers:

N/A

15% match (publications)

[Xiao-Wei Shi, Ming-Lv Sun, Bo Zhou, Xiao-Yun Wang. "Identification, characterization, and overexpression of a phytase with potential industrial interest", Canadian Journal of Microbiology, 2009](#)

7% match (publications)

[Chun-Shan Quan, Wen-Jie Tian, Sheng-Di Fan, Jun-Ichi Kikuchi. "Purification and properties of a low-molecular-weight phytase from Cladosporium sp. FP-1", Journal of Bioscience and Bioengineering, 2004](#)

4% match (publications)

[Ida Chapaval Pimentel, Chirlei Glienke-Blanco, Juarez Gabardo, Rodrigo Makowiecky Stuart, João Lúcio Azevedo. "Identification and colonization of endophytic fungi from soybean \(Glycine max \(L.\) Merrill\) under different environmental conditions", Brazilian Archives of Biology and Technology, 2006](#)

3% match (publications)

[Martin, J.A.. "Purification and physico-chemical characterisation of genetically modified phytases expressed in Aspergillus awamori", Bioresource Technology, 200609](#)

Pakistan Journal of Nutrition 9 (5): 471-474, 2010 ISSN 1680-5194 © Asian Network

for Scientific Information, 2010 **Isolation, Characterization and**

Production of Phytase from Endophytic Fungus its

Application for Feed Yetti Marlida¹, Rina Delfita², Peri Adnadi² and Gita

Ciptaan¹ ¹Department of Animal Nutrition, Faculty of Animal Science, Andalas

University, Padang, Indonesia ²Department of Biology, Andalas University, Padang,

Indonesia Abstract: Thirty four isolates of endophytic fungus produce phytases were

isolated from leaf, stem and root fragments of soybean. Two isolates were the best of

phytases enzyme producer and identified as Rhizoctonia sp. and Fusarium

verticillioides. [The phytase production was](#) induced [by phytate in medium used.](#)

[The](#) crude [preparations were used in subsequent](#) characterization

[studies,](#) pH and temperature optimum and [compared to other phytases tested](#)

[and is thus a promising candidate for animal feed applications.](#)

The results showed that optimal production of phytase from Rhizoctonia sp. were pH 4.0 and temperature 50oC and pH 5.0, temperature 50oC for Fusarium verticillioides. Key words: Endophytic fungus, phytase, soybean, Rhizoctonia sp., Fusarium verticillioides

INTRODUCTION Phytate (myo-inositol-hexaphosphate) is the major form of

phosphorus stored in cereals, pollens, legumes and oil seeds. Phytate is known as an

anti-nutrient factor, since it chelates various metal ions such as Mg⁺², Ca⁺², Zn⁺², Fe⁺², Fe⁺³ and forms complex with proteins (Pallauf and Rimbach, 1996; Martin et al., 2005; Cao et al., 2007; Liu et al., 2007). Moreover, phytate is not

metabolized by monogastric animals, which have low levels phytate- degrading enzymes in their digestive tracts, Thereby, inorganic phosphate has to be added to feeds to ensure a sufficient phosphate supply for these animals. Consequently, the

phytate in animal feeds is discharged in feces of these animals into waterways,

which contributed to eutrophication for surface waters, particularly in areas

of livestock production (Takizawa, 1998). One way to enhance phosphate utilization from phytate is the use of phytase. To obtain a good source of phytase, a variety of microorganisms, animals tissue and plant have been screened for enzyme. Several plant phytases in wheat, barley, bean, corn, soybean, rice and cotton have been studied extensively (Greiner and Konietzny, 2006). Microbial sources are Bacillus sp. (Poward and Jagannathan, 1982), Eschericia coli (Greiner et al., 1993a,b), Enterobacter (Yoon et al., 1996) Raoutella sp. (Greiner et al., 1997; Shah and Parekh, 1990), Aspergillus niger, Aspergillus fumigatus, Aspergillus terreus (Howson and Davis, 1983) and ruminal bacteria (Yanke et al., 1998). Several types of fungal phytase are available on the market from several companies such as Gist Brocades Co, Novo Nordisk Co and Kyowa Hakko Kogyo Co (Takizawa, 1998). At the end of 20th

century, annual sales of phytase as feed additive were estimated at US\$ 500

million and are continuing to rise (Vats and Banerjee, 2004). In Indonesia, the

phytase has been new research. The possibility of using these phytases in industry has not investigated. However, more work needs to be done to obtain

superior enzyme for industrial applications. It includes screening for strain that produce high phytase with better physicochemical properties, including high

thermostability and suitable pH, along with gen cloning. The objectives of this study were to isolate, characterization production of phytase enzyme from endophytic fungus and its applications for feed. MATERIALS AND METHODS Calcium phytate was made in the laboratory by adding phytic acid into a saturated calcium hydroxide solution. Sodium phytate and sodium dodecyl sulfate were sourced from

Sigma. All other reagents were domestic products of analytical grade. Isolation

of endophytic fungus: Isolation of phytase producers was performed by the agar plate of method Quan et al. (2001). Leaf, stem and root fragments sample of soybean [Glycine max (L.) Merril] were obtained from a farmer garden in Padang, Indonesia.

All leaf stem and root samples were washed twice in distilled water then surface sterilized by immersion for 1 minute in 70% (v/v) ethanol, 4 minutes in

sodium hypochlorite [3% (v/v) available chlorine] and 30 sec in 70% (v/v)

ethanol and then washed three times in sterilized distilled water for 1 min each time. After surface sterilization, the samples were cut into 5-7 mm pieces and

aseptically transferred to plates containing 0.1% Ca-phytate; 1.5% glucose;
0. 2% NH₄NO₃; 0.05% KCl; 0.05% MgSO₄•7H₂O; 0.03% MnSO₄•4H₂O; 0.03%
FeSO₄•7H₂O and 1,5% agar. The final pH was adjusted Corresponding
Author: Yetti Marlida, Department of Animal Nutrition, Faculty of Animal Science,
Andalas University, Padang, Indonesia to 5.5. Cultivation carried out at 28oC for 2-5
days. Fungal colonies, capable of hydrolyzing Ca-phytate which can be recognized by
their surrounding clear halo, were selected and repeatedly streaked onto solid Potato
Dextrose Agar (PDA) plates. Colonies which developed on the plates were inspected for
their morphology. Pure colonies were obtained by replating single colonies.
Identification of fungal phytase was determined with using of methods Samson and
Van Reenen-Hoekstra (1988); Barnett and Hunter (1972). Screening of endophytic for
phytase produser: Each of isolated strains was grown in 50 ml of liquid medium (0.1%
Ca-phytate; 1.5% glucose; 0. 2% NH₄NO₃; 0.05% KCl; 0.05% MgSO₄•7H₂O;
0.03% MnSO₄•4H₂O; 0.03% FeSO₄•7H₂O, pH 5.5) in 500-ml Sakaguchi flask
and incubated at 28oC for 48 h on reciprocal shaker (200 rpm). Cells collected from 1
ml of culture by centrifugation at 5000 x g for 10 min in cool room (4oC). Then, the
collected cells were resuspended in acetate buffer (0.2 M, pH 5.5) and used for the
phytase activity assay. Measurement of enzymatic activity: The phytase
activity assay was determined by measuring the amount of liberated inorganic
phosphate according to a method of Quan et al. (2001). Reaction mixture
consisted of 0.8 ml acetate buffer (0.2 M, pH 5.5) containing 1 mMNa -phytate
and 0.2 ml of cell suspension. After incubation for 30 min at 37oC, the
reaction was stoped by adding 1 ml of trichloroacetic acid. A 1 ml aliquot
was analyzed for inorganic phosphate liberated by method Kim and Lei
(2005). One unit of enzyme activity was defined as the amount of enzyme
liberating 1 nmol of inorganic phosphate per minute. Enzymatic characterization
studies: The effect of the pH on the activity of phytase was examined from pH 2
.0- 8.0 in 100 mM buffer. The buffers used were as follows: pH 1.0-
3.5: Gly-HCl; pH 3.5-6.0: NaAc-NaOH; pH 6.0-7.0: Tris- HAc; pH 7.0-
8.0: Tris-HCl. Temperature versus enzyme activity was measured over a range of
28-80oC. RESULTS Identification of isolates: A total 34 endophytic fungal strains were
screened for their ability to produce extracellular phytase. Only two strains,
forming clear peripheral zones on turbid agar plate, were isolated from root samples
and their activities were determined using liquid culture. According to the results of
morphological observation were classified as fungi. They are Rhizoctonia sp. and
Fusarium verticillioides (Fig. 1). The changes of phytase activity in fermentation were
shown in Fig. 2. The phytase activity of Rhizoctonia sp. was 0.77-2.72 U/ml and F.
verticillioides was 0.79-6.11 U/ml. a b A B b c d C D Fig. 1: Morphology of endophytic

fungal phytases. A-B = *Rhizoctonia* sp.; C-D = *F. verticillioides*; (100x); a = sclerotia; b = hifa; c = phialid; d = macroconidia [Fig. 2: pH versus activity profiles of](#)

[phytase](#) from *Rhizoctonia* sp. and *F. verticillioides* Enzymatic characteristics: Phytase production in culture of *Rhizoctonia* sp. and *F. verticillioides* by submerged fermentation reached its stationary growth phase after cultivation for 24 h. During exponential growth phase, its produced small amounts of phytase. Phytase production occurred in late stage of the exponential growth phase and the activity phytase increased gradually with increasing incubation time. Both of enzyme activity increased abruptly and reached the maximal value of 0.46 U/ml and 0.77 U/ml respectively. pH: The phytase activity of *Rhizoctonia* sp. reached the highest point at pH 4.0 for 15 min and phytase activity of *F. verticillioides* at pH 5.0 (Fig. 2). Temperature: Both fungal phytase displayed maximum activity were at temperature of 50oC (Fig. 3). At pH 8.0 and at temperature 80oC almost all enzymatic activity of both fungal were lost. DISCUSSION Endophytic fungal phytase was isolated from leaf, stem and root fragments sample of soybean and identified as *Rhizoctonia* sp. and *F. verticillioides* was identified for further study. This is the first report of *Rhizoctonia* sp. and *F. verticillioides* are extracellular phytase and exhibiting high phytase activity. The phytase synthesis in *Rhizoctonia* sp. and *F. verticillioides* by phytate in the culture medium. It may be concluded that only phytate induced these enzymes. Many other phytase producing strains as *Escherichia coli*, *Pseudomonas* sp and *Raoutella* sp. also were induced by phytate (Shah and Parekh, 1990; Konietzny and Greiner, 2004). Enzyme induction is due to physiological change in a whole microbial population and it involves an accelerated rate of renewed formation of enzyme in response to a relatively specific nutritional stimulus (Rhodes and Fletcher, 1966). The phytase was induced early phase of cultivation. It seems that the phosphates are released from phytate. Phytase often has a low-pH optimum range (pH 4.5-6.0) with a rapid drop in activity at pH value above 6.0. Yeast Fig. 3: Temperature versus activity profiles of phytase from *Rhizoctonia* sp. and *F. verticillioides* phytases also have an optimal range 4.0-5.0 (Cao et al., 2007; Quan et al., 2001; Nakamura et al., 2000). The phytase in both *Rhizoctonia* sp. and *F. verticillioides* have pH 4.0 and 5.0 respectively and most stable at pH range 2.0-7.0. were very [compatible with the internal environment of](#)

[monogastric animals'](#) stomach [such as](#) in pigs and [poultry](#). Compared to many other phytase producing strains which exhibit low enzyme activity at pH values associated with the upper digestive tract, the *Rhizoctonia* sp. and *F. verticillioides* phytase activity is significantly higher, reaching levels of commercial acceptability. The optimum temperature of the these phytase did not reveal differences

between [phytase from *Aspergillus niger* N-3](#) and Natuphos phytase, [the](#) latter exhibiting maximum activity at 50oC (Martin et al., 2005). [At present, a major drawback to the widespread use of phytases in animal feed is the constraint of thermal stability required for these enzymes to withstand inactivation during the feed-pelleting](#)

[or expansion processes](#) (Cao [et al.](#), 2007). Both of phytases exhibited maximum activity as high as Natuphos and pGP209 phytase, were at 50oC (Martin et al., 2005). This phytase reached levels of commercial acceptability. [This phytase is worthy of further research as retains activity over a wide range of pH values characteristic of digestive tract and could conceivably be more suited to increasingly higher feed processing temperatures currently employed in the animal feed industry.](#)

ACKNOWLEDGEMENT This work was supported by Grant from DIKTI (Hibah Bersaing). The authors are grateful to the laboratories staff at Microbiology Lab. Universitas Andalas for their support during the experiment. REFERENCES Barnett, H.L. and B.B. Hunter, 1972. Illustrated Genera of Imperfect Fungi. 3rd Edn. Burgess Publishing Company. USA. Cao, L., W. Wang, C. Yang, Y. Yang, J. Diana, A. Yakupitiyage, Z. Luo and D. Li, 2007. Application of Microbial Phytase in Fish Feed. J. Enzyme and Microbial Technol., 40: 497-507. Greiner, R. and U. Konietzny, 2006. Phytase for Food Application. Food Technol. Biotechnol., 44: 125-140. Greiner, R., U.

Konietzny and K.D. Jany, 1993a. Purification and Characterization of Two Phytases From *Escherichia coli*. Arch. Biochem. Biophys., 303: 107-113. Greiner, R., U. Konietzny and K.D. Jany, 1993b. Purification and Characterization of Two Phytases From *Escherichia coli*. Arch. Biochem. Biophys., 303: 107-113. Greiner, R., E. Haller, U. Konietzny and K.D. Jany, 1997. Purification and Characterization of A Phytase From *Klebsiella Terrigena*. Arch. Biochem. Biophys., 341: 201-206. Howson, S.J. and R.J. Davis, 1983. Production of Phytate-Hydrolysing Enzyme by Fungi. J. Enzyme Microbiol. Technol., 5: 377-382. Kim, T.W. and X.G. Lei, 2005. An Improved Method for a Rapid Determination of Phytase activity in animal feed. J. Anim. Sci., 83: 1062-1067. Konietzny, U. and R. Greiner, 2004. Bacterial Phytase: Potential Application, In vivo Function and Regulation of Its Synthesis. Braziliat. J. Microbiol., 35: 11-18. Liu, Z., H. Wang, E. Xiu Wang, H. Xu, D. Gao, G. Zhang, P. Chen and D. Liu, 2007. Effect of Wheat Perling on Flour Phytase Activity, Phytic Acid, Iron and Zinc Content. Food Sci. and Technol. Swiss. Martin, J.A., R.A. Murphy and R.F.F. Power, 2005. Purification and Physico-Chemical Characterisation of Genetically Modified Phytases Expressed in *Aspergillus awamori*. Bioresource Technol., 97. Irlandia. Nakamura, Y., H. Fukuhara and Sano, 2000. Secreted Phytase Activities of Yeast. Bioscience Biotechnol. and Biochem., 64: 841-844. Pallauf, J. and G. Rimbach, 1996. Nutritional Significance of Phytic Acid and Phytase. Arch. Anim. Nutr., 50: 301-319. Pimentel, I.C., C. Glienkeblanco, J. Gabardo, R.M. Stuard and J.L. Azevedo, 2006. Identification and Colonization of Endophytic Fungi from Soybean [*Glycine max* (L.) Merril] under Different Environmental Conditions. J. Brazilian Archives of Biol. and Technol., 45: 705-711. Powar, V.K. and V. Jagannathan, 1982. Purification and Properties of Phytase-Specific Phosphatase from *Bacillus subtilis*. J. Bacteriol., 151: 1102-1108. Quan, C., L. Zhang, Y. Wang and Y. Ohta, 2001. Production of Phytase in Low Phosphate Medium by a Novel Yeast *Candida krusei*. J. Biosci. and Bioeng., 9: 154-160. Rhodes, A. and D.L. Fletcher, 1966. Principles of Industrial Microbiology. Pergamon Press. New York. Shah, V. and L.J. Parekh, 1990. Phytase From *Klebsiella* sp. No. Pg-2: Purification and Properties. Indian J. Biochem. Biophys., 27: 98-102. Samson, R.A. and E.S. van Reenen-Hoekstra, 1988. Introduction to Food-Borne Fungi. Centraalbureau voor Schimmelcultures. Wageningen. Takizawa, N., 1998. Utilization of Recombinant Phytase on the Avoidance of Water Pollution by Phosphorus of Excretory Organ of Liverstock. Seitbutsu-kakagu, 76. Vats, P. and U.C. Banerjee, 2004. Production Studies and Catalytic Properties of Phytase. Enzymes Technol. J., 35: 3-14. Yanke, L.J., H.D. Bae, L.B. Selinger and K.J. Cheng, 1998. Phytase Activity of Anaerobic Ruminant Bacteria. Microbiol., 144: 1565-1573. Yoon, S. J., Y. J. Choi., H. K. Min., K. K. Cho., J. W. Kim., S. C. Lee and Y. H. Jung. 1996. Isolation and Identification of Phytase-Producing Bacterium, *Enterobacter* sp. 4, and Enzymatic Properties of Phytase Enzyme. Enzyme Microbiol. Technol., 18: 449-454. Pak. J. Nutr., 9 (5): 471-474, 2010 Pak. J. Nutr., 9 (5): 471-474, 2010 Pak. J. Nutr., 9 (5): 471-474, 2010 471 472 473 474