

## Turnitin Originality Report

- Processed on: 23-Feb-2020 8:35 PM +08
- ID: 1262260558
- Word Count: 4878
- Submitted: 1

# A Review on Potential of Glutamate Producing Lactic Acid Bacteria of West Sumatera's Fermented Food Origin, as Feed Additive for Broiler Chicken.

Similarity Index

7%

### Similarity by Source

Internet Sources:  
N/A

Publications:  
7%

Student Papers:  
N/A

1% match (publications)

[Vabera Maslami, Yetti Marlida, Mirnawati., Jamsari., Yuliaty Shafan Nur. "Optimization of Glutamate Production from Lactobacillus plantarum Originating from Minangkabau Fermented Food as a Feed Supplement for Broiler", Pakistan Journal of Nutrition, 2018](#)

1% match (publications)

["Ethnic Fermented Foods and Alcoholic Beverages of Asia", Springer Science and Business Media LLC, 2016](#)

< 1% match (publications)

[Jinap, S., M.S. Mohd-Mokhtar, A. Farhadian, N.D.S. Hasnol, S.N. Jaafar, and P. Hajeb. "Effects of Varying Degrees of Doneness on the Formation of Heterocyclic Aromatic Amines in Chicken and Beef Satay", Meat Science, 2013.](#)

< 1% match (publications)

[Maryoris E.S. Lopez, Marco T.P. Gontijo, Delaine M.G. Boggione, Luiz A.A. Albino, Laís S. Batalha, Regina C.S. Mendonça.](#)

["Microbiological Contamination in Foods and Beverages: Consequences and Alternatives in the Era of Microbial Resistance", Elsevier BV, 2018](#)

< 1% match (publications)

[Frederick Adzitey. "Antibiotic resistance of Escherichia coli and Salmonella enterica isolated from cabbage and lettuce samples in Tamale metropolis of Ghana", International Journal of Food Contamination, 2018](#)

< 1% match (publications)

[Belal J. Muhiadin, Hana Kadum, Mohammad Zarei, Anis Shobirin Meor Hussin. "Effects of metabolite changes during lacto-fermentation on the biological activity and consumer acceptability for dragon fruit juice", LWT, 2020](#)

< 1% match (publications)

[Lili Anggraini, Yetti Marlida, Wizna Wizna, Jamsari Jamsari, Mirzah Mirzah, Frederick Adzitey, Nurul Huda. "Molecular identification and phylogenetic analysis of GABA-producing lactic acid bacteria isolated from indigenous dadih of West Sumatera, Indonesia", F1000Research, 2019](#)

< 1% match (publications)

[Li-Oon Chuah, Ahamed Kamal Shamila-Syuhada, Min Tze Liong, Ahmad Rosma, Kwai Lin Thong, Gulam Rusul. "Physio-chemical, microbiological properties of tempoyak and molecular characterisation of lactic acid bacteria isolated from tempoyak", Food Microbiology, 2016](#)

< 1% match (publications)

[Hiroyuki KOBAYASHI. "Regulation of muscular glutamate metabolism by high-protein diet in broiler chicks : FEED AND MUSCULAR GLU METABOLISM", Animal Science Journal, 02/2011](#)

< 1% match (publications)

[K.Y Sim, F.Y. Chye, A. Anton. " Chemical composition and microbial dynamics of fermentation, a traditional Malaysian fish sauce ", Acta Alimentaria, 2015](#)

< 1% match (publications)

[Mardijah, Guisheng Zhai, Dieky Adzkiya, Lutfi Mardianto, Muhammad Ikhwan. "Modified T2FSMC approach for solar panel systems", Systems Science & Control Engineering, 2019](#)

< 1% match (publications)

["Nutraceuticals in Veterinary Medicine", Springer Science and Business Media LLC, 2019](#)

< 1% match (publications)

[Audina Putri, Komang Gede Wiryawan, Toto Toharmat, Sri Suharti. "Potential of lactic acid bacteria from asam durian as a probiotic candidate for chicken", IOP Conference Series: Materials Science and Engineering, 2019](#)

< 1% match (publications)

[Bezerra, R. M., F. G. P. Costa, P. E. N. Givisiez, E. R. Freitas, C. C. Goulart, R. A. Santos, J. G. Souza, P. A. Brandão, M. R. Lima, M. L. Melo, V. P. Rodrigues, E. T. Noqueira, and D. V. G. Vieira. "Effect of l -glutamic acid supplementation on performance and nitrogen balance of broilers fed low protein diets", Journal of Animal Physiology and Animal Nutrition, 2015.](#)

< 1% match (publications)

[C. Feurer. "Taxonomic characterization of nine strains isolated from clinical and environmental specimens, and proposal of Corynebacterium tuberculostearicum sp. nov.", INTERNATIONAL JOURNAL OF SYSTEMATIC AND EVOLUTIONARY MICROBIOLOGY, 07/01/2004](#)

< 1% match (publications)

[Ramesh C. Ray. "Traditional and novel fermented foods and beverages from tropical root and tuber crops: review", International Journal of Food Science & Technology, 06/2009](#)

< 1% match (publications)

[Theron, . "Biopreservation", Organic Acids and Food Preservation, 2010.](#)

< 1% match (publications)

[Sabine Greisinger, Stefan Jovanovski, Gerhard Buchbauer. " An Interesting Tour of New Research Results on and Compounds ", Natural Product Communications, 2016](#)

< 1% match (publications)

[Jazila El Malti, Hamid Amarouch. "Microbial and Physicochemical Characterization of the Horse Meat in Fermented Sausage", Food Biotechnology, 2008](#)

< 1% match (publications)

[Khashayar Morshed-Behbahani, Pooria Najafisayar, Mahmoud Pakshir, Nika Zakerin. "The effect of surface modification on the corrosion protection ability of the passive films of sensitized UNS S31803 duplex stainless steel", Corrosion Engineering, Science and Technology, 2018](#)

< 1% match (publications)

[P. Dietrich. "Physiology and biophysics of plant ligand-gated ion channels : Physiology and biophysics of ligand-gated ion channels", Plant Biology, 09/2010](#)

JWPR 2018, Scieline Publication J. World Poult. Res. 8(4): 120-126, December 25, 2018 Journal of World's Poultry Research Review Paper, PII: S2322455X1800017-8 License: CC BY 4.0 A Review on Potential of Glutamate Producing Lactic Acid Bacteria of West Sumatera's Fermented Food Origin, as Feed Additive for Broiler Chicken Vebera Maslami<sup>1</sup>, Yetti Marlida<sup>2\*</sup>, Mirnawati<sup>2</sup>, Jamsari<sup>3</sup>, Yuliaty Shafan Nur<sup>2</sup>, Frederick Adzitey<sup>4</sup> and Nurul Huda<sup>5</sup> <sup>1</sup>Department of Agricultural Science, Faculty of Agriculture, Andalas University, 25163, Indonesia <sup>2</sup>Department of Animal [Nutrition and Feed Technology,](#)

[Faculty of Animal Science, Andalas University,](#) 25163, Indonesia <sup>3</sup>Department of Crop Science, Faculty of Agriculture, Universitas Andalas, 25163, Indonesia <sup>4</sup>Department [of Veterinary Science, University for Development Studies, Box TL](#)

[1882, Tamale, Ghana](#) <sup>5</sup>Faculty [of Food Science and Nutrition, Universiti Malaysia Sabah, 88400 Kota Kinabalu,](#)

[Sabah, Malaysia](#) \*Corresponding author' s Email: [yettimarlida@ansci.unand.ac.id](mailto:yettimarlida@ansci.unand.ac.id); ORCID: 0000-0001-5395-1297

ABSTRACT [Received: 30 Oct 2018 Accepted: 02 Dec 2018](#) Increasing broiler populations must be supported by cheap and high quality feed. Improving the quality of feed can be done by adding feed additives. Glutamate is a non-essential amino acid that can be used as a feed additive in the form of flavoring agents in broiler feed which functions as a

neurotransmitter of taste, basic structure of proteins, and in metabolism [of the](#) body. [Lactic Acid Bacteria](#)

(LAB) are one [of the](#) microbes that [are](#) considered faster and safe in producing glutamate. Fermented foods of West Sumatera, Indonesia origin serve as sources of LAB include dadih (fermented milk), asam durian (fermented durian), ikan budu (fermented fish) and tapai (fermented rice and cassava). The West Sumatra's fermented foods are

potential sources of glutamate. Supplementation [of glutamate in broiler](#) diet [can increase body weight,](#) protein digestibility, reduce faecal ammonia and improve carcass quality (improve umami taste, and reduce bruises and abdominal fat). Key words: Carcass quality, Feed additive, Fermented food, Glutamate, Lactic acid bacteria, Performance INTRODUCTION Broiler is a meat-producing animal that has enough potential to meet people's needs for animal protein. This is because broiler meat is relatively inexpensive and easily obtained compared to other animal proteins. High broiler populations must be supported by high quality feed to improve the performance and quality of broiler carcasses. One important component of feed needed to increase growth and feed efficiency is feed additive. According to Slyamova et al. (2016), feed additives has function in improvement of immune system against disease attacks and effects of stress, increase appetite, stimulate growth, and increase the production of meat and eggs. Feed additives can be flavoring agents, antibiotics, enzymes, antioxidants, hormones, probiotics and anticoccidials (Lesson and Summers, 2001). One of the feed additives that can be used in animal feed is glutamate. Glutamate is a feed additive that serves as flavoring agent which helps to increase feed consumption. According to Fujimura et al. (2001), glutamate functions as a neurotransmitter for taste, where glutamate is serves as umami, flavors or savory. Besides functioning as a glutamate flavoring agent, it functions as a protein building block, a substrate in the synthesis of protein, precursor of glutamine and helps the body's metabolism (Young and Ajami, 2000). Giving glutamate to broilers has been carried out by several researchers. According to Berres et al. (2010), administering glutamate to broilers decreases abdominal fat, improves meat examination by reducing red bruises on carcasses and increasing meat protein content. The administration of 0.75% glutamate in feed increases

muscle free glutamate and improves the [taste of meat \(Fujimura et al., 2001; Imanari et al., 2008\)](#). Furthermore, administration of glutamate improves broiler performance, reduce the feed conversion ratio, affect the length of intestinal

villi, reduce raw ration protein and reduce faecal ammonia (Zulkifli [et al., 2016;](#) Ribeiro [et al., 2015;](#) Bezerra

[et al., 2016](#)). [To cite this](#) paper: [Maslami](#) V, [Marlida](#) Y, [Mirnawati, Jamsari, Shafan Nur](#)

Y, Adzitey F and Huda N (2018). **A Review on Potential of Glutamate Producing Lactic Acid Bacteria of West Sumatera's Fermented Food Origin, as Feed Additive for Broiler Chicken.** J. World Poult. Res., 8 (4): 120-126. <http://jwpr.science-line.com> 120

The sources of glutamate are animals, plants and microbes. Microbes are mostly used to produce glutamates probably because it is easier, faster, safer and relatively inexpensive compared to other sources. Glutamate production can be done by fermentation or by chemical processes. At present, most glutamates are produced by fermentation because it is superior to chemical processes. The fermentation process produces L-glutamate while the chemical process produces racemic glutamate (D- and L-glutamic acid) (Sono, 2009). The difference of L-glutamate and D-glutamate is their uses in the body. According to Wijayasekara and Wansapala (2017), L-glutamate is the form glutamate is used in the body. Lactic Acid Bacteria (LAB) are examples of microbes that produce glutamate. According to Lucke (2000), LAB is one of the gram-positive bacterium that produces glutamate which is considered safe and environmentally friendly. A number of LAB isolated from different sources have been reported to produce glutamate. These include LAB from tapai (Ishak et al., 2017), Lactobacillus plantarum MNZ from fermented food (Zareian et al., 2012), Bacillus spp from vegetable protein (Lawal et al., 2011) and Lactobacillus from skim

milk (Zalan et al., 2010). West Sumatera is rich [in the production of fermented foods](#) which [are](#) sources of

[LAB](#). Fermented foods in West Sumatera that are widely consumed contain dadih (fermented milk), asam durian (fermented durian), ikan budu (fermented fish) and tapai (fermented rice and cassava) (Mustopa and Fatimah, 2014). Purwandhani et al. (2018) isolated LAB from dadih of West Sumatera origin. Yusra (2014) reported of Bacillus spp from ikan budu in Pariaman. Furthermore, Chuah et al. (2017) found six LAB strains in asam durian (*Fructobacillus durionis*, *Lactobacillus plantarum*, [Lactobacillus fructivorans](#), [Leuconostoc dextranicum](#), [Lactobacillus collinoides](#) and [Lactobacillus](#)

[paracasei](#)) and Sujaya et al. (2001) *Weissella* spp in tapai. There is high diversity of fermented foods in West Sumatera, therefore, it is necessary to explore their nature and potentials as sources of LABs capable of producing glutamate, which can serve as feed additive for broilers. Food Fermentation from West Sumatra Fermentation was carried out in the past based on non-scientific studies and much was not known about the role of microbes in changing food characteristics. It was based on traditional techniques of storing and handling food ingredients which turns out to produce new food products that are different from the original ones. The goal of initial food fermentation was to preserve seasonal foods that were easily damaged. During fermentation, the digestibility and nutritional quality of foods are improved. There is

the addition of vitamins, and [essential amino acids and fatty acids during fermentation](#) (Steinkraus, 2002). Indonesia has many fermented foods scattered throughout the region. Some fermented foods in Indonesia include: Brem (fermented rice) from Madiun and Wonogiri, danke (fermented buffalo milk) from Sulawesi, lemea (fermented bamboo) from Bengkulu, oncom (fermented tofu waste) from West Java, pakasam and wadi (fermented fish) from various regions of Indonesia, tempe (fermented soybeans) from various regions of Indonesia, gatot and growol (fermented dried yam) from Yogyakarta, Tempoyak (fermented durian) from various regions of Indonesia and urutan (pig fermentation) from Bali (Nuraida, 2015). Even though, the same fermented foods may be produced in different regions, the manufacturing process are slightly different. West Sumatera also has a lot of fermented foods, among them are dadih, asam durian, ikan budu and tapai. Dadih is a fermented food from milk which is a potential functional food and a source of probiotic. Dadih processing is still done traditionally without any standard processing procedure. Curds are made from buffalo milk poured into a bamboo tube to ferment naturally for 24 to 48 hours at room temperature. The natural fermentation process in making curds involves various types of microbes found on the surface of the inner bamboo tube, the surface of the cover leaves and from buffalo milk used. According to Hasim et al. (2017), dadih produced in West Sumatera are made from buffalo milk by relying on natural microbes as the inoculum. The bamboo is usually covered with taro leaves, banana leaves, plastic, or left without a cover, depending on the region (Maskiyah and Broto, 2011). Isolation and use of LAB in dadih have been reported by several researchers. In a study by Surono (2003), LAB in curds were dominated by *Lactococcus* bacteria consisting of *Lactococcus lactis*, *Leuconostoc mesenteroides*, *Lactococcus brevis*, *Lactococcus casei*, *Lactococcus plantarum*, and *Enterococcus faecium*. Pato et al. (2005) found the following LAB, *Lactobacillus casei* subsp *casei*, *Leuconostoc paramesenteroides*, *Enterococcus faecalis* subspecies *liquefaciens*, and *Lactococcus lactis* subsp *lactis* in dadih. Asam [durian Asam durian is a](#)

[fermented product](#) made [from](#) durian. Fermented foods are found in several regions (West Sumatra, Jambi and Riau) of Sumatera, Indonesia. The process of making asam durian and the name given to it differ among each region. In Jambi of Indonesia, durian fermentation is called tempoyak while in West Sumatera and Riau it is known as asam durian. Fermentation of durian is done by the Jambi community by adding salt (Anggraini and Widawati, 2015). LAB found in salt-containing asam durian includes *Lactobacillus* species, *Lactobacillus plantarum*, *Pediococcus acidilactici* and *Weissella paramesenteroides* (Yuliana and Dizon, 2011). Asam durian found in West Sumatera Indonesia has differences from other regions. The manufacturing process is spontaneous fermentation, by natural microorganisms found in nature without adding salt during incubation. Six types of bacteria have been identified in asam durian. They are: *Lactobacillus plantarum*, *Pediococcus acidilactici*, *Leuconostoc mesenteroides*, *Staphylococcus saprophyticus*, *Lactobacillus curvatus*, and *Micrococcus variance* (Hasanudin, 2010). Ikan budu Ikan budu is a product made from fermented fish and mainly from coastal areas of Pasaman Regency and Padang Pariaman in West Sumatera, Indonesia. In Indonesia, it is found only in West Sumatra. This fermented food product is made from large sea fish and has white meat. Example of fishes used for making budu are mackerel (*Scomberomorus guttatus*) and chamfer (*Chorinemus lyson* L). Budu fish has a distinctive aroma, not rotten, non-rancid and soft-textured. Yusra et al. (2014) found that *Bacillus* and *Micrococcus* species were involved in ikan budu fermentation. Malaysia also produce budu but the fermentation process and type of fish used is different from that of West Sumatra. Budu from Malaysia appear as thick brown, and it is prepared from salted raw anchovy (*Stolephorus* spp.) and stored in a large concrete tank at 30-40°C for 6-12 months (Sim et al., 2015). Sim et al. (2015) identified *Pediococcus* (9.4%), *Candida* (8.0%), *Micrococcus* species (28.7%), *Staphylococcus* (26.7%), [Lactobacillus \(6.7%\)](#), [Saccharomyces](#)

[\(4.0%\)](#), and [Lactococcus \(2.7%\)](#) from Malaysian budu. Fujimura et al. (2012) also isolated LABs from Malaysian budu. Tapai Tapai is one of Indonesia's traditional foods produced from the fermentation of carbohydrate foods such as cassava and sticky rice. Tapai fermentation process takes place because of the activities of several types of microorganisms such as bacteria, yeast and fungi. Tapai is produced by fermenting cassava with yeast, LAB, and amylolytic bacteria (Sujana, 2001). Different from other places, glutinous tapai from Indonesia, especially from West Sumatra, uses black glutinous rice stored in containers coated with banana leaves, while in other areas guava leaves (*Syzygium*) or rubber (*Hevea brasiliensis*)

are used. Suhartatik et al. (2014) isolated LAB [such as L.](#) *mesenteroides*, *L.* *pentosus*, *L. brevis*, and

*L.* *plantarum*, from Tapai. Also, Maslami et al. (2017) isolated *Lactobacillus* sp from tapai of West Sumatera origin. In Malaysia, different ingredients and fermentation processes are used in tapai preparation. Tapai from Sabah, Malaysia is an alcoholic drink but not food (Campbell-Platt, 2000). In Sabah, the basic ingredients for making tapai are rice, cassava, pineapple and corn which are fermented using cultural starter from Sasak (Chiang et al., 2006). Tapai is also produced at Peninsula, Malaysia. The processing method employed in Peninsula, Malaysia is similar to how tapai is prepared in Singapore and Brunei, that is, they all carry out fermentation in banana leaves (Campbell-Platt, 2000). LAB found in tapai of Sabah

origin include *L. brevis*, *L. plantarum*, [P. pentosaceus](#), *L.* *lactis* and *L.* *s paracasei* (Chiang et al.,

2006). [L. casei](#) is also involved in the fermentation process of tapai of Malaysia origin (Adnan and Tan, 2007). LAB Producing Glutamate Research on glutamate producing microbes has been carried out by several researchers. Glutamate producing microbes isolated from various sources can be seen in table 1. Effects of glutamate on performance and quality of

broiler carcasses Glutamate [plays an essential role in](#) the [metabolism and](#) increases feed consumption.

[The](#) purposes and dosages for the administration glutamate in broiler is presented in table 2. Table 1. Glutamate producing lactic acid bacteria and their source of isolation Researcher Source Bacteria Production Tarek and Mustafa (2009) Egyptian infants Lactobacillus paracasei 68.78 mg/l. Zareian et al. (2012) Fermented soybean Fermented durian Fermented tapioca Fermented rice Fermented shrimp Fermented fish Lactobacillus plantarum 489 Mmol/ L 20 Mmol/ L 59 Mmol/ L 65

Mmol/ L 11 Mmol/ L 106 Mmol/ L [Fudou et al. \(2002\)](#) Soils [and](#) vegetable [Corynebacterium.](#) - Lawal [et](#)

[al.](#) (2011) Vegetable protein Bacillus subtilis Bacillus pumilus Bacillus licheniformis Bacillus polymyxa 8.4 mg/ml 8.2 mg/ml 6. 4 mg/ml 6.2 mg/ml Table 2. Purpose and dosage of glutamate administration Researcher Purpose Glutamate dose Stilborn and Moran (2010) Increase performance and anticoccidial 1.75 % Zulkifli et al. (2016) Growth and physiological anti-stress on broiler chicks 1% Joshua et al. (2015) Anti-stress due to heat stress in the tropics Stater (0.50% - 1 %) Finisher (0.50% - 1 %) Ribeiro et al. (2015) Improve performance and immune system 0.4 % Shakeri et al., (2014) Anti-stress on broiler chicks which are affected by heat stress 0.50 % Berres et al. (2010) Improve feed efficiency Age 1-7 days (0.13% and 0.26%) Age 8-21 days (1.46%, 1.66% and 3.13%) Age 22-35 days (1.24%, 1.40% and 3.50%) Age 36-42 days (1.04%, 1.18% and 3.22%) Khadiga et al. (2009) Increase performance 1% Bezerra et al. (2015) Efficient use of feed protein Age 1-7 days (crude protein 20.29% and glutamate 3.41) Age 8-21 days (crude protein 18.99 and glutamate (3.32) Age 22-35 (crude protein 16.24 and glutamate 2.64 Age 36-45 (crude protein 17.04 and glutamate 2.96) Improve performance Glutamate is known as a conditional or non- essential amino acid. Glutamate [binds with other amino acids to](#)

[form protein](#) structures (Berres et al., 2010). Weight gain in 21 day broilers given an additional 1% of glutamate supplements was significantly higher (11%) than those fed without 1% glutamate addition (Bartell and Batal, 2007). The positive effect of glutamate on broiler chickens have been reported by other researchers. According to Bazzera et al. (2014), broilers fed on glutamate containing feed (1.66% up to 3.13%) performed better than those fed without glutamate supplemented feed during heat stress. Feeding broilers on glutamate increases broiler body weight and the deposition of connective tissues at growth phase (Zhang et al., 2008). According to Zulkifli et al. (2016) administration of 1% glutamate in rations can increase body weight, reduce feed consumption, improve feed conversion, and reduce death. Administration of 2% glutamic acid reduces crude protein of ration and can improve performance and reduce faecal ammonia (Ribeiro et al., 2015; Bezerra et al., 2016). Glutamate is essential for the maintenance of intestinal mucosa and seen as an efficient source of a non-specific nitrogen source. This is because its energy source is for rotating the mucosa, through ATP which results from the citric acid cycle (Berres et al., 2010). The synthesis of nitric oxide in some tissues (e.g. brain tissue) are regulated by glutamate (Li et al., 2007). Glutamate also act as precursor for glutathione. Glutathione is an important compound

responsible for elimination of oxidants and immune system modulation (Li [et al.](#), 2007). [According to Silva et al.](#)

[\(2001\)](#), adding L- glutamate (5, 10 and 15%) to broiler feed takes care of protein deficiencies in rations with efficiencies greater than 10%. Carcass quality Glutamate is a vital amino acid responsible for flavor enhancement and as feed additive (Fujimura et al., 2001). In its free form or unbound to other amino acids in protein, glutamate has a flavor enhancing effect (Yamaguchi and Ninomiya, 2000). This free glutamate functions effectively as a flavor generator and plays a role in improving delicacy. Free glutamate also known as an important taste component of meat and contributes to meat tastes (umami, brothy taste and delicious). The chicks of poultry contains more free glutamate than pork and beef (Kato and Nishimura, 1987). Pork also contain more free glutamate than beef (Kato and Nishimura 1987). The hydrolysis of protein by the supplementation of heat during cooking causes glutamate to release protein molecule that can causes umami taste (Khropychheva et al., 2009). However, when protein molecules bind with glutamate, umami taste is not released and make the protein source tasteless (Khropychheva et al., 2009). Glutamate supplementation has a significant effect on broiler carcass quality. Low glutamate content in broiler meat will result in less glutamate being stored in muscle fibers. Glutamate level can become low because it is widely used for metabolism and as an energy source by intestinal cells. When free glutamate is present in broiler meat its umami taste reduces (Tang et al., 2009). Glutamate content in broiler meat can be increased by adding glutamate to broiler feeds. Glutamate can also improve the quality of broiler carcasses. According to Ajinomoto (2007) and Bezerra et al. (2015) glutamate can increase meat protein, reduce meat fat, reduce meat bruises and improve meat flavor. Meat protein is increased because glutamate is a precursor of several amino acids (Berres et al., 2010). In addition, a decrease in meat fat is caused by foods that contain good source of amino acids and less energy for fat accumulation. Moran and Stilborn (1996) reported that, there was a decrease in bruises and thigh deformation and an increase in connective tissue deposition at growth phase of broilers fed by glutamate supplemented feed. CONCLUSION Glutamate can improve carcass traits, performance and quality of meat. One of the microbes that can produce glutamate is LAB. A potential and unexplored LAB source, that produce the glutamate found in fermented foods in West Sumatera,

Indonesia. DECLARATIONS Acknowledgment [We are very grateful to the](#) Minister [of Research, Technology and](#)

[Higher Education of the Republic of Indonesia](#) for support [and](#) finance through PMDSU No: No: 1387 / E4.

2015 and support through the B-Class [World Class Professor Program](#) Scheme [123.](#) 57 / [D.2.3 / KP /](#)

[2018.](#) [Competing interest](#) The authors declare that they have no competing interests. Author's contribution All

the authors have made contribution to the study. Consent to publish All the authors gave their informed consent prior to their inclusion in the study. REFERENCES Adnan AFM and Tan IK (2007). Isolation of lactic acid bacteria from Malaysian foods and assessment of the isolates for industrial potential. *Bioresource Technology*, 98: 1380– 1385. DOI: 10.1016/j.biortech.2006.05.034. Anggraini L and Widawati L (2014). Pengaruh waktu fermentasi tempoyak terhadap sifat organoleptik sambal tempoyak. *Agritepa*, 1(2): 118-127. Bartell SM and Batal AB (2007). The Effect of Supplemental Glutamine on Growth Performance, Development of the Gastrointestinal Tract, and Humoral Immune Response of Broilers. *Poultry Science*, 86:1940-1947. DOI: 10.1093/ps/86.9.1940. Berres J, Vieira SI, Dozier III, Cortes MEM, De Barros R, Nogueira ET and Kutschenko M (2010). Broiler responses to reduced-protein diets supplemented with valine, isoleucine, Glycine, and glutamic acid. *The Journal of Applied Poultry Research*, 19: 68–79. DOI: 10.3382/japr.2009-00085. Bezerra RM, Costa FGP, Givisiez PEN, Freitas ER, Goulart CC, Santos RA, Souza JG, Brand~ao PA, Lima MR, Melo ML, Rodrigues VP, Nogueira ET and Vieir DVG (2016). Effect of L-glutamate acid supplementation on performance and nitrogen balance of broilers fed low protein diets. *Journal of animal physiology and animal nutrition*, 100: 590–600. DOI: 10.1111/jpn.12405. Campbell-Platt G (2000). *Fermented Foods in Encyclopedia of Food Microbiology*. Robinson RK, Batt CA and Patel, PD (second edition). London: Academic Press 736-773. Chiang YW, Chye FY and Mohd Ismail A (2006). Microbial diversity and proximate composition of tapai, a sabah's fermented Beverage. *Malaysian Journal of Microbiology*, 2(1): 1-6. Chuah L, Shamila-Syuhada AK, Liong MT, Rosma A, Thong KL and Rusul G (2016). Physio-chemical, microbiological properties of tempoyak and molecular characterisation of lactic acid bacteria isolated from tempoyak. *Food Microbiology*, 58: 95-104. DOI: 10.1016/j.fm.2016.04.002. Fudou R, Jojima Y, Seto A, Yamada K, Kimura E, Nakamatsu T, Hiraisi A and Yamanaka S (2002). *Corynebacterium efficiens* sp. nov, a glutamic acid- producing species from soil and vegetables. *International Journal of Systematic and Evolutionary Microbiology*, 52: 1127–1131. DOI: 10.1099/00207713-52-4-1127. Fujimura S, Sakai F and Kadowaki M (2001). Effect of restricted feeding before marketing on taste active components of broiler chickens. *Journal of Animal Science*, 72: 223–229. DOI: 10.2508/chikusan.72.223. Hasim, Mustopa AZ, Andrianto N, Fatimah and Faridah DN (2017). Antioxidant Production of Lactic Acid Bacteria Isolated from Indonesian Traditional Fermented Buffalo Milk (Dadih). *IOSR Journal of Pharmacy and Biological Sciences*, 12 (5): 76-82. DOI: 10.9790/3008-1205057682. Imanari M, Kadowaki and Fujimura (2008). Regulation of taste-active components of meat by dietary branched- chain amino acids; effects of branched-chain amino acid antagonism. *British Poultry Science*, 49(3): 299-307. DOI: 10.1080/00071660802155080. Ishak MSMd, Ibrahim MZ and Ishak Hmd (2017). Screening of Lactic Acid Bacteria from "TapaiPulut" for Biosynthesis of Glutamic Acid. *Modern Agricultural Science and Technology*, 3: 37-41. DOI: 10.15341/mast (2375- 9402)/03.03.2017/007. Joshua O, Olubodun, Zulkifli I, Farjam AS, Hair- Bejo M and Kasim A (2016). Glutamine and glutamic acid supplementation enhances performance of broiler chickens under the hot and humid tropical condition. *Italian Journal of Animal Science*, 14 (1): 25-29. DOI: 10.4081/ijas.2015.3263/ Kato H and Nishimura T (1987). Taste components and condition of beef, pork and chicken. In: Kawamura Y, Kare RM (eds), *Umami, A Basic Taste*, pp. 289–306. Marcel Dekker, New York. Kong SE, Hall JC. Khadiga AAA, Mohammed S, Saad AM and Mohamed HE (2009). Response of broiler chicks to dietary monosodium glutamate. *Pakistan Veterinary Journal*, 29(4): 165-168. ISSN (online): 2074-7764. Khropycheva R, Zolotarev V, Uneyama H and Torii K (2009). Effect of free dietary glutamate on gastric secretion in dogs. *Annals of the New York Academy of Sciences*, 1170: 87–90. DOI: .1111/j.1749-6632.2009.03900. x. Lawal A K, Oso BA, Sanni AI and Olatunji OO (2011). L- Glutamic acid production by *Bacillus* Spp. Isolated from vegetable proteins. *African Journal of Biotechnology*, 10(27): 5337-5345. DOI: 10.5897/AJB10.1000. Lesson S and Summers JD (2001). *Broiler Breeder Production*. University Books. Guelph, Ontario, Canada. ISBN- 13: 978-1899043392. Li P, Yin YL and Li D (2007). Amino acids and immune function. *British Journal of Nutrition*, 98: 237- 252. DOI: 10.1017/S000711450769936X. Lucke FK (2000). Utilization of microbes to process and preserve meat. *Meat Science*, 56: 105–115. DOI: 10.1016/S0309-1740(00)00029-2. Maskiyah and Broto B (2011). Pengaruh kemasan terhadap kualitas dadih susu sapi. *Buletin Peternakan*, 35(2): 96- 106. DOI: 10.21059/buletinpeternak.v35i2.596. Maslami V, Marlida Y, Mirnawati M, Jamsari J, and Nur YS (2018). Isolasi bakteri asam laktat (BAL) penghasil asam glutamat dari ikan budu sebagai feed suplemen ayam broiler. *Jurnal Peternakan Indonesia*, 20(1): 45-51. ISSN: 1907-1760. Moran ET Jr and Stilborn HL (1996). Effect of glutamic acid on broilers given submarginal crude protein with adequate essential amino acids using feeds high and low in potassium. *Poultry Science*, 1(1) 120–129. DOI: 10.3382/ps.0750120. Mustopa AZ and Fatimah (2014). Diversity of lactic acid bacteria isolated from Indonesian traditional fermented foods. *Microbiology*, 8(2): 48-57. DOI: 10.5454/mi.8.2.2. Nuraida L (2015). A review: Health promoting lactic acid bacteria in traditional Indonesian fermented foods. *Food Science and Human Wellness*, 4(2): 47-55. DOI: 10.1016/j.fshw.2015.06.001. Nuraida L (2015). A review: Health promoting lactic acid bacteria in traditional Indonesian fermented foods. *Food Science and Human Wellness*, 4: 47–55. DOI: 10.1016/j.fshw.2015.06.001. Pato UMA and Parlindungan AK (2005). Taurocholate deconjugation and cholesterol binding by indigenous dadih lactic acid bacteria. *Hayati Journal of Biosciences*, 12(3): 103–107. DOI: 10.1016/S1978-3019(16)30334-5. Purwandhani SN, Utami T, Millati R and Rahayu ES (2018). Isolation, characterization and screening of folate- producing bacteria from producing fermented food (dadih). *International Food Research Journal*, 25(2): 566- 572. ISSN (Online): 2231 754. Ribeiro JV, Albino LFT, Rostagno HS, Hannas N, Ribeiro CLN, Vieira RA, Araújo WAG, Pessoa GBS, Vessias RKG and Silva DL (2015). Effects of dietary l-glutamine or l-glutamine plus l-glutamic acid supplementation programs on the performance and breast meat yield uniformity of 42-d-old broilers. *Brazilian Journal of Poultry Science*, 93-98. DOI: 10.1590/1516-635X/SPECIALISSUENutrition-PoultryFeedingAdditives093- 098. Sano C (2009). History of glutamate production. *The American Journal of Clinical Nutrition*, 90: 738-732. DOI: 10.3945/ajcn.2009.27462F. Shakeri M, Zulkifli I, Soleimani 1 AF, O'Reilly E L, Eckersall PD, Anna AA, Kumari S and Abdullah FFJ (2014). Response to dietary supplementation of L-glutamine and L-glutamate in broiler chickens reared at different stocking densities under hot, humid tropical conditions. *Poultry Science*, 93: 1–9. DOI: 10.3382/ps.2014-03910. Silva FA, Moraes GHK, Rodrigues ACP, Oliveira MGA, Rostagno HS, Albino LFT, Fonseca CC and Minafra CS (2001). Efeitos do Ácido L-Glutâmico e da Vitamina D3 no Desempenho e nas Anomalias Ósseas de Pintos de Corte. *Revista Brasileira de Zootecnia*, 30(6): 2059-2066. DOI: 10.1590/S1516-35982001000800015. Sim KY, Chye FY and Anton A (2015) Chemical composition and microbial dynamics of budu fermentation, a traditional malaysian fish sauce. *Acta Alimentaria*, 44 (2):185–194. DOI: 10.1556/AAlim.2014.0003. Slyamova AY, Sarsembayeva NB, Ussenbayev AE and Paritova AY (2016). Influence of functional feed additive at the basis of the chankanay deposit's zeolite to the intestinal microbiocenosis of broiler chickens. *International Journal of Advances in Chemical Engineering and Biological Sciences*, 3 (1): 85-87. DOI: 10.15242/IJACEBS. AE0416119. Steinkraus KH (2002). *Fermentations in world food processing. Comprehensive Reviews in Food Science and Food Safety*, 24–32. DOI: 10.1111/j.1541-4337.2002.tb00004.x. Stilborn HL and Moran Jr E T (2010). Effect of added l- glutamic acid on male broiler performance when using wheat- or corn-based diets and 2 different anticoccidials. *Poultry Science Association*, 93: 401–414. DOI: 10.3382/japr.2009-00116. Suhartatik N, Cahyanto MN, Rahardjo S, Miyashita M and Rahayu ES (2014). Isolation and identification of lactic acid bacteria producing  $\beta$  glucosidase from Indonesian fermented foods. *International Food Research Journal*, 21(3): 973-978. ISSN (Online): 2231 754. Sujaya IN, Amachi S, Yokota A, Asano K and Tomita F (2001). Identification and characterization of lactic acid bacteria in ragi tape. *World Journal of Microbiology and Biotechnology*, 17: 349–357. DOI: 10.1023/A:1016642315022. Surono IS, Pato U, Koesnandar and Hosono A (2009) In vivo antimutagenicity of dadihprobiotic bacteria towards Trp- P1. *Asian-Australasian Journal of Animal Sciences*, 22 (1): 119–123. DOI: 10.5713/ajas.2009.80122. Tang H, Gong YZ, Wu CX, Jiang J, Wang Y and Li K (2009). Variation of meat quality traits among five genotypes of chicken. *Poultry Science*, 88: 2212-2218. DOI: 10.3382/ps.2008-00036. Tarek M and Mostafa HE (2010). Screening of potential infants' lactobacilli isolates for amino acids production. *African Journal of Microbiology Research*, 4: 226–232. ISSN 1996-

0808. Wijayasekara K and Wansapala J (2017). Uses, effects and properties of monosodium glutamate (MSG) on food and nutrition. *International Journal of Food Science and Nutrition*, 2(3): 132-143. DOI: 10.22271/food. Ymanagguchi S and Ninomia K (2000). The use and utility of glutamate as flavoring agents in food. *Journal of Nutrition*, 130:921-926. Young VR and Ajami AM (2000). Glutamate: an amino acid of particular distinction. *Journal of Nutrition*, 130: 892-900. DOI: 10.1093/jn/130.4.892S. Yuliana N and Dizon EI (2011). Phenotypic identification of lactic acid bacteria isolated from Tempoyak (fermented durian) made in the Philippines. *International Journal of Biology*, 3(2): 145-152. DOI: 10.5539/ijb.v3n2p145. Yusra Y, Azima F, Novelina N and Periadnadi P (2014). Isolasi dan identifikasi mikroflora indigenous dalam budu. *Agritech*, 34(3): 316-321. DOI: 10.22146/agritech.9460. Zalan Z, Hudacek J, Stetina J, Chumchalava J and Halasz A (2010). Production of organic acids by *Lactobacillus* strains in three different media. *European Food Research and Technology*, 230: 395-404. DOI: 10.1007/s00217-009-1179-9. Zareian M, Ebrahimpour A, Bakar FA, Mohamed AKS, Forghani B, Ab-Kadir MSB and Saari N (2012). A glutamic acid-producing lactic acid bacteria isolated from Malaysian Fermented foods. *International Journal of Molecular Sciences*, 13: 5482-5497. DOI: 10.3390/ijms1305548. Zhang GQ, Ma QG and Ji C (2008). Effects of dietary inosinic acid on carcass characteristics, meat quality, and deposition of inosinic acid in broilers. *Poultry Science*, 87(7): 1364-1369. DOI: 10.3382/ps.2007-00193. Zulkifli M, Shakeri and Soleimani AF (2016). Dietary supplementation of L-glutamine and L-glutamate in broiler chicks subjected to delayed placement. *Poultry Science*,

1: 1-7. DOI: 10.3382/ps/pew267 Maslami et al., 2018 [J. World's Poult. Res.](#), 8 (4): 120-126, 2018 Maslami

et al., 2018 [J. World's Poult. Res.](#), 8 (4): 120-126, 2018 Maslami et al., 2018 [J. World's Poult. Res.](#), 8 (4): 120-126, 2018 121 122 123 124 125 126