

Turnitin Originality Report

- Processed on: 23-Feb-2020 8:35 PM +08
- ID: 1262260532
- Word Count: 4838
- Submitted: 1

Isolation and Characterization of Lactic Acid Bacteria Producing GABA from Indigenous West Sumatera Fermented Food

Similarity Index

14%

Similarity by Source

Internet Sources:
N/A

Publications:
14%

Student Papers:
N/A

3% match (publications)

[Lili Anggraini, Yetti Marlida, Wizna Wizna, Jamsari Jamsari, Mirzah Mirzah, Frederick Adzitey, Nurul Huda. "Molecular identification and phylogenetic analysis of GABA-producing lactic acid bacteria isolated from indigenous dadih of West Sumatera, Indonesia", F1000Research, 2019](#)

2% match (publications)

[YS Ismail, C Yulvizar, B Mazhitov. "Characterization of lactic acid bacteria from local cow 's milk kefir", IOP Conference Series: Earth and Environmental Science, 2018](#)

1% match (publications)

[Haixing Li. "Production of gamma-aminobutyric acid by Lactobacillus brevis NCL912 using fed-batch fermentation", Microbial Cell Factories, 2010](#)

1% match (publications)

[Tsvetanka Teneva-Angelova, Ivelina Hristova, Atanas Pavlov, Dora Beshkova. "Lactic Acid Bacteria—From Nature Through Food to Health", Elsevier BV, 2018](#)

1% match (publications)

[Anupama Gupta, Nivedita Sharma. "IN VITRO CHARACTERIZATION OF LACTIC ACID BACTERIA ISOLATED FROM LASODA BARI- A RARE FERMENTED FOOD OF HIMACHAL PRADESH-INDIA FOR POTENTIAL PROBIOTIC ATTRIBUTES", Journal of Microbiology, Biotechnology and Food Sciences, 2017](#)

1% match (publications)

[Joseph P. Brown, Carl Lamanna. "Role of gut bacterial flora in nutrition and health: A review of recent advances in bacteriological techniques, metabolism, and factors affecting flora composition", C R C Critical Reviews in Food Science and Nutrition, 2009](#)

1% match (publications)

[Lilis Nuraida. "A review: Health promoting lactic acid bacteria in traditional Indonesian fermented foods", Food Science and Human Wellness, 2015](#)

< 1% match (publications)

[Al-Haik, Wedad Mohammed Saeed. "Inhibition of Toxicogenic Microorganisms by Lactic Acid Bacteria, Isolated from Milk and Fermented Food = المخمر والغذاء الحليب من المعزولة اللاكتيك حمض بكتيريوبا بواسطة ل لسعوم المتجة الميكروبيات ت تيط", King Abdulaziz University : Scientific Publishing Centre, 2012](#)

< 1% match (publications)

[Mardijah, Guisheng Zhai, Dieky Adzkiya, Lutfi Mardianto, Muhammad Ikhwan. "Modified T2FSMC approach for solar panel systems", Systems Science & Control Engineering, 2019](#)

< 1% match (publications)

[Lal Krishna. "THE IDENTIFICATION, PRODUCTION AND CHARACTERIZATION OF NATTOKINASE, BACTERIOCIN FROM BACTERIAL SPECIES", Bacterial Empire, 2019](#)

< 1% match (publications)

[Radhika Dhakal, Vivek K. Bajpai, Kwang-Hyun Baek. "Production of gaba \(γ - aminobutyric acid\) by microorganisms: a review", Brazilian Journal of Microbiology, 2012](#)

< 1% match (publications)

[Lili Anggraini, Yetti Marlida, Wizna Wizna, Jamsari Jamsari, Mirzah Mirzah, Frederick Adzitey, Nurul Huda. "Molecular identification and phylogenetic analysis of GABA-producing lactic acid bacteria isolated from indigenous dadih of West Sumatera, Indonesia", F1000Research, 2019](#)

< 1% match (publications)

[Chang-jiang Lyu, Wei-rui Zhao, Sheng Hu, Jun Huang, Tao Lu, Zhi-hua Jin, Le-he Mei, Shan-jing Yao. " Physiology-Oriented Engineering Strategy to Improve Gamma-Aminobutyrate Production in ", Journal of Agricultural and Food Chemistry, 2017](#)

< 1% match (publications)

[Kanintra Suwanmanon, Pao-Chuan Hsieh. " Isolating and optimizing its fermentative medium for GABA and nattokinase production ", CyTA - Journal of Food, 2013](#)

< 1% match (publications)

[Mardiati Zain, Rusmana W S Ningrat, Erpomen, Ezi Masdia Putri, Malik Makmur. " The effects of leguminous supplementation on ammoniated rice straw based completed feed on nutrient digestibility on microbial protein synthesis ", IOP Conference Series: Earth and Environmental Science, 2019](#)

< 1% match (publications)

[Huda, Nurul. "Indonesian Fermented Fish Products", Handbook of Animal-Based Fermented Food and Beverage Technology Second Edition, 2012.](#)

< 1% match (publications)

[Indri Juliyarsi, Arief, Akmal Djamaan, Endang Purwati. "Characteristics based of edible film made from whey with isolated lactic acid bacteria from tempoyak as probiotics packaging", IOP Conference Series: Earth and Environmental Science, 2019](#)

< 1% match (publications)

Min Kim, Seockmo Ku, Sun Kim, Hyun Lee, Hui Jin, Sini Kang, Rui Li, Tony Johnston, Myeong Park, Geun Ji. "Safety Evaluations of *Bifidobacterium bifidum* BGN4 and *Bifidobacterium longum* BORI", *International Journal of Molecular Sciences*, 2018
 < 1% match (publications)
 Fauzia Agustín, Elihasridas, Indri Juliyarsi. "The effect of urea supplementation and incubation time in fermentation process of bagasse by using on the growth of and the nutritive value of bagasse", *IOP Conference Series: Earth and Environmental Science*, 2019
 < 1% match (publications)
 Milena Brasca, Johannes A. Hogenboom, Stefano Morandi, Veronica Rosi, Paolo D'Incecco, Tiziana Silvetti, Luisa Pellegrino. "Proteolytic Activity and Production of γ -Aminobutyric Acid by Cultivated in Microfiltered Pasteurized Milk", *Journal of Agricultural and Food Chemistry*, 2016
 < 1% match (publications)
 R Jawan, ME Kasimin, SN Jalal, AA Mohd. Faik, S Abbasliasi, A Ariff. "Isolation, characterisation and evaluation of bacteriocins-producing lactic acid bacteria from fermented products of Northern Borneo for their beneficial roles in food industry", *Journal of Physics: Conference Series*, 2019
 < 1% match (publications)
 Woong-Kyu Yoon, Coralia V. Garcia, Chan-Schick Kim, Sam-Pin Lee. "Fortification of Mucilage and GABA in *Hovenia dulcis* Extract by Co-fermentation with *Bacillus subtilis* HA and *Lactobacillus plantarum* EJ2014", *Food Science and Technology Research*, 2018
 < 1% match (publications)
 Gopal Reddy, Md. Altaf, B.J. Naveena, M. Venkateshwar, E. Vijay Kumar. "Amylolytic bacterial lactic acid fermentation — A review", *Biotechnology Advances*, 2008
 < 1% match (publications)
 Manuel Venturi, Viola Galli, Niccolò Pini, Simona Guerrini, Lisa Granchi. "Use of Selected Lactobacilli to Increase γ -Aminobutyric Acid (GABA) Content in Sourdough Bread Enriched with Amaranth Flour", *Foods*, 2019
 < 1% match (publications)
 Eman Zakaria Gomaa. "Enhancement of γ -Aminobutyric Acid Production by Co-Culturing of Two Lactobacilli Strains", *Asian Journal of Biotechnology*, 2015
 < 1% match (publications)
 Qiuhua Bao, Jie Yu, Wenjun Liu, Manjun Qing, Weihong Wang, Xia Chen, Fang Wang, Meihua Li, Hongmei Wang, Qiang Lv, Heping Zhang. "中国四川地区传统发酵牦牛乳制品中优势乳酸菌的分离和鉴定", *Dairy Science & Technology*, 2012
 < 1% match (publications)
 Nant Kay Thwe Moe, Su Myo Thwe, Takaaki Shirai, Takeshi Terahara, Chiaki Imada, Takeshi Kobayashi. "Characterization of lactic acid bacteria distributed in small fish fermented with boiled rice in Myanmar", *Fisheries Science*, 2015
 < 1% match (publications)
 Lactic Acid Bacteria Genetics Metabolism and Applications, 1999.
 < 1% match (publications)
 Weiguo Lu. "Properties of Glutamate Decarboxylase of *Aspergillus oryzae* and Its Application for Biotransformation of Glutamate into γ -Aminobutyric Acid", 2010 4th International Conference on Bioinformatics and Biomedical Engineering, 06/2010
 < 1% match (publications)
 George K. Auer, Douglas B. Weibel. "Bacterial Cell Mechanics", *Biochemistry*, 2017

Vol.9 (2019) No. 3 ISSN: 2088-5334 Isolation and Characterization of

Lactic Acid Bacteria Producing GABA from Indigenous West Sumatera Fermented Food ^{Lili}

Angraini#, Yetti Marlida#*, Mirzah#, Wizna#, Jamsari+, Nurul Huda# #Department [Animal Nutrition and Feed Technology](#),

[Faculty of Animal Science, Andalas](#) University, [Padang](#), 25163, [Indonesia](#) E-mail: *yettimarlida

@ansci.unand.ac.id +Department of Crop Science, [Faculty of](#) Agriculture, Universitas [Andalas](#), 25163,

[Indonesia](#) [Faculty of Food Science and Nutrition, Universiti Malaysia Sabah](#), 88400 [Kota Kinabalu, Sabah](#),

[Malaysia](#) Abstract— The purpose of this study was to isolate and characterize [lactic acid](#)

[bacteria](#) (LAB) from indigenous West Sumatera fermented foods. The aim was to obtain LAB isolates that

have the ability to produce γ -aminobutyric acid (GABA) and to characterize the highest GABA producing LAB. The indigenous West Sumatera fermented foods sampled were dadih (fermented buffalo milk), asam durian (fermented durian), tape singkong (fermented cassava) and ikan budu (fermented fish). The conventional method was used to isolate LAB, after which LAB were screened for γ -amino butyric acid (GABA) and GABA producing LAB characterized by biochemical tests. A total of 704 isolates were successfully isolated. Five hundred fifty two (552) isolates were identified as LAB and 103 isolates were confirmed as GABA producing LAB. The highest GABA producing LAB was DS15 and produced 49.365 mg/ml of GABA. DS18 produced the least GABA of 3.415 mg/ml. Biochemical assay of DS15 showed that the isolate could grow on the MRSA, undergoes aerobic respiration, do not use lactose, glucose, mannitol, arginine, aesculin, arabinose, raffinose, sorbitol, trehalase and xylose/melezitose. The isolate DS15 was also negative for arginine, nitrate reduction, catalase and oxidase tests. Based on the gram stain reaction, the isolate DS15 was a gram-positive bacterium and could be classified as

Lactobacillus sp. Keywords— [lactic acid bacteria](#); [\$\gamma\$ -aminobutyric acid](#); indigenous fermented [food](#). I.

INTRODUCTION [\$\gamma\$ -Aminobutyric acid \(GABA\) is the](#) major [inhibitory neurotransmitter in the](#) mammalian

[central nervous system](#) (CNS) [1]. [GABA](#) is formed by decarboxylation of L- glutamate, a reaction catalyzed by

an enzyme that depends on the peridoxal phosphate of decarboxylated L-glutamate. This decarboxylase enzyme is

present in the central nervous system [2]. Dhakal et al., [3] reported that GABA can improve plasma

concentration, growth hormones and protein synthesis in the brain, and reduce blood pressure in animals and

humans. Dysfunction of the gamma-aminobutyric (GABA) in central nervous system has long been associated with anxiety disorders, such as epilepsy, anxiety, alcoholism, angelman's syndrome, autism, depression, premenstrual syndrome, sleep disorders, and alzheimer's disease [4]. For that reason, GABA is regarded as very important for physiological function in the mammalian central nervous system, induction of hypotensive, diuretic effects, and

tranquilizer effects for human [5]. GABA was reported to take part in preventing heat-induced stress in broilers [6] and increasing body weight gain in pigs [7]. GABA is naturally present in animals, plants and microorganisms (bacteria, yeast and fungi) [3]. Thus GABA can be produced from a variety of microorganisms. Lactic acid bacteria (LAB) is generally

regarded as safe, therefore, LAB -producing GABA can be directly used in functional foods [8]. The LAB-producing GABA has become a focus of research in recent years due to their special physiological activity and safety. Some strains of LAB are able to change glutamate and produce GABA and CO₂, such reaction will produce energy

that can be used for ATP synthesis [9]. Many fermented food products can be used as sources for the isolation of

LAB. A number of works have reported on the isolation of LABs from different types of fermented food

products such as Cheese [10], Kimchi [11], and fish [12] that could produce GABA. West Sumatera Province has many indigenous fermented food products such as dadih, asam durian, tape singkong and ikan budu that can be used as

source of LAB. Isolation and screening of such LAB from naturally fermented foods of West Sumatera Province

origin is rare, and novel food products often have present means for obtaining useful cultures for scientific

and commercial purposes [13]. Therefore, this study was conducted to isolate and characterize

LAB GABA from indigenous West Sumatera fermented foods. II. MATERIALS AND METHODS A. Sample

Collection Samples from indigenous fermented food products (dadih, asam durian, tape singkong and ikan budu) were collected from different traditional markets in West Sumatera, Indonesia. Dadih was collected from Aiadingin, Sijunjung and Solok, Asam durian was collected from Padang and Solok, tape singkong was collected from Bukittinggi and ikan budu was collected from Tiku. The samples were collected between June-August 2017 and analyzed at the Faculty of Animal

Science Andalas University, West Sumatera, Indonesia. B. Isolation of lactic acid bacteria Isolation of LAB was done according to the protocol described by [14] with some modifications. Ten (10) g of sample (dadih, asam durian, tape singkong or ikan budu) was mixed with 90 ml MRS Broth (Merck, Darmstadt, Germany) in a 250 ml Erlenmeyer flask prior to adding 2 g of glucose (Merck, Darmstadt, Germany). The mixture was incubated at 30°C at 100 rpm for 7 days (Incubator shaker series I26, New Brunswick Scientific, United States). A dilution of 10⁻¹² was done after incubation using aquabides. One (1) ml of diluted mixture was spread evenly in the three petri dishes containing selective media of MRS Agar (Merck, Darmstadt, Germany). All colonies were tested for the formation of clear zone by adding 2% of CaCO₃ (Merck, Darmstadt, Germany) to the MRS agar and incubated at 30°C for 24 hours. Isolates that could produce clear zone in the medium were considered as LAB. C. Qualitative screening of GABA producing LAB Each selected LAB isolate was grown in a production medium consisting of MRS broth with the addition of 50 mM of L-glutamate (Merck, Darmstadt, Germany) as an inducer. Incubation was carried out at a temperature of 30°C under anaerobic for three days. After incubation, the media was

centrifuged at 10,000 rpm for 20 minutes at 4°C (Refrigerated Centrifuge TGL-20M, China). The supernatant was collected and analyzed using TLC (Thin Layer Chromatography) (Merck, Germany) to check for γ-Aminobutyric acid (GABA) production. Commercial GABA (Sigma-Aldrich, China) was used as standard and assayed on a TLC plate using n-Butanol: acetic acid: H₂O (v/v/v) as mobile phase. The TLC plate was subjected to a ninhydrin reagent and heated (60°C) for 3-5 minutes. Spots and colors formed in this assay was compared to the standard [15]. D. Quantitative screening of GABA producing LAB Quantitative screening was performed to determine the GABA concentrations produced during incubation. The quantification was done using the method described by Spies [16]. Aliquot from each sample was taken from 1 ml supernatant and added to 1 ml of 0.1% ninhydrin reagent. After which it was heated in water bath for 5 minutes, cooled under running water and the absorbance measured at 570 *Hovenia dulcis* Extract by Co-fermentation with *Bacillus subtilis* HA and *Lactobacillus plantarum* EJ2014", *Food Science and Technology Research*, 2018"> nm.

The GABA *Hovenia dulcis* Extract by Co-fermentation with *Bacillus subtilis* HA and *Lactobacillus plantarum* EJ2014",

Food Science and Technology Research, 2018">[content in the](#) sample Hovenia dulcis Extract by Co-fermentation with Bacillus subtilis HA and Lactobacillus plantarum EJ2014", Food Science and Technology Research, 2018">[was](#)

[determined using](#) GABA Hovenia dulcis Extract by Co-fermentation with Bacillus subtilis HA and Lactobacillus

plantarum EJ2014", Food Science and Technology Research, 2018">[standard](#) curve. E. Characterization of GABA producing LAB Characterization of LAB was performed according to the protocol described Gad et al., [17]. The biochemical test was conducted according to the Systematic Bacteriological Bergey's Manual [18]. The biochemical test covered: TSIA, catalase, oxidase, lactose, glucose, mannitol, VP, OF, arginine, aesculin, arabinose, raffinose, sorbitol, trehalase, maltose, melezitose, and nitrate tests. III. RESULTS AND DISCUSSION A. Isolation [of LAB from indigenous West Sumatera fermented](#)

[food](#) Fermentation is [one of the oldest forms of food preservation in the](#) word. [Fermentation can](#)

improve [the](#) storability and taste [of](#) food [and](#) food products. West Sumatera has a variety of

[fermented foods, such as dadih,](#) asam durian, [tape](#) singkong [and](#) ikan budu. In West Sumatera, fermented foods are produced on small scales as household business, and fermentation is done using traditional processes that are influenced by local culture. The fermentation process can occur naturally because of the microbes that are already present in these foods when they grow and/or the addition of microbial cultures which results in a more uniform product. A total of 704 isolates were successfully identified from all the fermented food samples examined. Distribution of the isolates and the number of GABA producing LAB is presented in Table 1. Fermented milk (Dadiah) obtained from Aiadingin produced the highest number of isolates (131 isolates) while tape singkong produced the least number of isolates (35 isolates). From the 704 isolates, only 552 isolates could be classified as LAB (based on the clear zone). The clear zone is formed from the reaction between lactic acid produced by LAB cell and calcium carbonate (CaCO₃) added to the medium. The Ca-lactic acid is seen as clear zones around the colony on the medium [19]. The present of CaCO₃ in the growing medium is an early step of LAB detection, where LAB show clear zones after the incubation period [20]. LAB needs complex nutrition such as [amino](#)

[acids, peptides, nucleotide bases, vitamins, fatty](#) acid, minerals [and carbohydrates](#) for their [growth](#)

[21]. Variation [of](#) LAB number [on](#) each food sample is due to the differences in the characteristics of the food material used for the fermentation process. Each animal and vegetable products produce different materials, so that there are differences in nutritional content available for the bacterial growth. Dadiah is fermented buffalo milk, and it is a [popular](#)

[dairy product in Bukittinggi, Padangpanjang, Solok, Lima Puluh Kota and Tanah Datar](#) [22]. Dadiah obtained from Bukittinggi have a nutritional content of 7.57% protein, 6.48% fat, 3.79% carbohydrates and 81.03% moisture. Ikan budu is fermented fish product from Pasaman and has been reported to have a proximate composition of [51% moisture content,](#)

[33% protein, 0.5% fat and 14% ash](#) [23]. Fermented food products derived from vegetables or plants have higher carbohydrate content. Tape singkong is made from fermented cassava, and has 32-35% carbohydrate, 0.1-0.3% fat [24] and 6.8% protein. Asam durian is made from fermented durian and contains carbohydrate 6.3%, protein 8.1%, and fat

6.6% [25]. TABLE I [DISTRIBUTION OF LAB ISOLATES FROM THE](#) INDIGENOUSWEST [SUMATERA FERMENED](#)

[FOOD](#) No Samples [Origin\(](#) s) isolates LAB Producing GABA 1 Dadih Aia dingin 131 125 24 Sijunjung 166 93 19 Solok 100 96 19 2 Ikan budu Pariaman 135 17 9 3 Asam durian Padang 132 130 13 Solok 75 59 9 4 Tape Bukit tinggi 35 32 10 B. Qualitative screening of GABA producing LAB Screening was performed to detect isolates that were capable of producing extracellular γ -Aminobutyric acid (GABA). Of the 552 LAB isolates obtained, only 103 isolates can produce extracellular GABA (data not shown). The highest number of GABA-producing LAB isolates (24 isolates), was found in dadiah obtained from Aiadingin, while the lowest number (9 isolates) was identified in asam durian from Solok and ikan budu from Pariaman. This can occur because of differences in the types of each isolate obtained, so that they have different GABA. Li et al., [26] explains that the [GABA-producing ability varies widely among the strains of LAB, and some GABA-producing LAB](#)

[strains have shown a great](#) promise [potential in large-scale fermentation for the](#) production [GABA.](#) C.

Quantitative [screening of GABA producing](#) LAB [In](#) order [to](#) identify [the](#) capability [of](#) LAB to produce GABA, putative promising isolates were further analyzed quantitatively (Table 2). The highest GABA production was shown by isolate DS15 (49.365 mg/ml), while the lowest GABA production was shown by isolate DS18 (3.415 mg/ml). Both isolates were collected from Dadiah. In general, isolates collected from Dadiah produced GABA in high concentration (16.123 mg/ml) compared to the other fermented foods (ikan budu, asam durian and tapai singkong) which produced 10.871mg/ml, 11.035 mg/ml and 12.110 mg/ml, respectively. The GABA production capability shown by DS15 was higher than that described by [27] from Lactobacillus brevis NCL912 (35.662 mg/ml). GABA [is formed by decarboxylation of L-](#)

[glutamate, a reaction catalyzed by an enzyme](#) dependent [on](#) pyridoxal [phosphate](#) [2]. Glutamate decarboxylase (GAD) is also produced by several microorganisms such as LAB. Research conducted by Marlida et al., [28] found a LAB that produced glutamate from dadiah with GABA as the precursor. GABA biosynthesis in microorganisms is regulated by several factors that will affect the fermentation process. These [factors are pH, temperature, time of the](#)

fermentation process, the nutrient content such as carbon and nitrogen, and inducer. The optimum conditions of fermentation from each bacterium could be different, this is due to differences in GAD enzymes catalyzing glutamic acid to GABA. Therefore, the characteristics of microorganisms especially LAB in producing GABA need to be elucidated in order to help achieve high GABA production. TABLE II GABA PRODUCTION OF ISOLATES COLLECTED FROM DIFFERENT FERMENTED FOOD SOURCES No Name of Isolates Gaba yield (mg/ml) 1 Da .9 29.12 2 Da.16 22.91 3 Da.20 29.82 4 Da.21 44.15 5 Da.24 35.02 6 Ds.2 42.34 7 Ds.3 31.52 8 Ds.15 49.40 9 Ap.8 22.62 10 Ap.5 16.12 D. Characteristic of GABA producing LAB Since DS15 showed the highest GABA production (Table 2), further analysis was undertaken on this isolate (Table 3). The isolate DS15 was tested for its oxygen requirement, and the isolate was found to be aerobe or grow under oxygen

conditions. LAB is a type of microorganism that prefers to grow under anaerobic conditions, but LAB can also

grow under oxygenated conditions, also called anaerobic aerotrophic bacteria [21]. So, it can be concluded that the whole bacterial isolate is anaerobic aerotolerant. TSIA Test (Triple Sugar-Iron Agar) is used to determine the ability of bacteria to ferment lactose, glucose and to produce acids. This ability is based on the carbohydrate fermentation pathway. In the TSIA test, when the color of the media turns red at the top and bottom (M/M), it indicates the absence of carbohydrate fermentation, when it turns red at the top and yellow at the bottom (M/K), it indicates the bacteria can decompose glucose, and when the media turns yellow at the top and bottom (K/M), the bacteria is said to ferment lactose and/or sucrose [29]. TSIA test for DS15 isolate showed red at the top and bottom (M/M), which means that, DS15 does not use glucose, lactose and sucrose as a source of energy. This can happen because of the many other carbon sources that can be utilized by bacteria as a source of energy. In addition, bacteria also use nitrogen such as amino acids and peptides as an energy source. The catalase test was performed to distinguish microorganisms that have catalase enzymes used to dexter hydrogen

peroxide which is toxic [30]. Catalase reaction is positive when there is formation of air bubbles, which

indicates the formation of O2 gas, and negative if it does not indicate the presence of gas bubbles. DS15 was not able to produce the catalase enzyme to convert hydrogen peroxide/H2O2 into water and oxygen, which is one of the

characteristics of lactic acid bacteria [31]. Therefore, DS15 it's a catalase negative bacterium. Similarly with Guessas

and Kihal [32] found that nine lactic acid bacteria isolated from fermented goat milk were catalase negative. Adam and Nout [33] also indicated that there are some lactic acid bacteria that show negative results for catalase. The oxidation test was performed to test the activity of the bacterial cytochrome oxidase [34]. Aerobic and facultative anaerobic bacteria will exhibit oxidase activity. The oxidase enzyme plays a role in the electron transport system during aerobic respiration. Cytochrome oxidase will catalyze the oxidation of the reduced cytochrome by oxygen producing H2O and H2O2 [35]. DS15 was negative for oxidase test (Table 3), that is, no color change occurred when p- aminodimethylaniline oxalate reagents were added. The reagent acts as an electron donor and will oxidize to a black compound if there is free oxidase and oxygen. Starch hydrolysis test (mannitol, aesculin, arabinose, raffinose, sorbitol, trehalase, maltose and melezitose) is used to determine the bacterial ability to excrete extracellular enzymes. Agar starch composed of nutrient agar with starch substrate and an indicator is used to test starch hydrolysis. Hydrolysis of starch is shown by differences in color change [34], [35]. For instance, aesculin media will turn black if bacteria are able to ferment it. From Table 3, D15 was negative for lactose, glucose, arginine, aesculin, arabinose, raffinose, sorbitol, trehalase, xylose and melezitose. Generally, bacteria use simple sugars as carbon sources, but if simple sugars are not available then bacteria will use complex sugars as a carbon source in

fermentation [35]. The VP test (Voges-Proskauer) is used to determine the ability of a bacterium to produce non-acidic substances or neutral end products, such as acetylmethylcarbinol from organic acids because of glucose metabolism [30], [34]. DS15 showed negative result for VP test. This is because there was no color change of the medium to red caused by the catalyst reaction of α -naphthol and guanidine group present in peptone from test medium (MR-VP) with

reagent Barritt's [29]. Amino acid test is perform to determine the ability of bacteria to degrade amino acid substrates. Alpha amino acid is composed of carbon (-C-), amino group (-NH₂), a carboxylic group (-COOH) and hydrogen atom (-H). Amino acid substrate undergoes decarboxylation (catalyze by decarboxylase enzyme with the elimination of the carboxylic group) to produce an amine or diamine and carbon dioxide. Amines are then used to synthesize the various molecules required by the cell. The final production of the amine will result in a change in pH. Tests using the amino acid arginine can be seen in Table 3. DS15 showed a negative result for this test envisage by the absence of discoloration of the medium to purple. The color change occurs in the medium containing bromocresol purple as an indicator when all oxygen is used up, and anaerobic respiration takes over leading to the production of CO₂ and altering the pH of the

medium. The presence of acid will change the indicator color to yellow which means the enzyme decarboxylase has been activated and the reverse will occur when alkaline is produced. Sunatmo, [35] indicated that when the active enzyme produces alkaline diamine (cadaverine) and CO₂, it causes the indicator to retain its purple color, which was the same as what was observed for DS15. Nitrate reduction test was performed to test the ability of DS15 to reduce nitrate (NO₃) to nitrite (NO₂). After incubation, organisms that reduce nitrate to nitrite will turn the medium to red color when reagent A which contains sulfanilate and alfa naphthylamine is added. If there is no color change, the bacteria may have nitrate reductase which can reduce nitrite to ammonia or molecular nitrogen. It is therefore necessary to add a bit of Zn powder to detect whether the nitrate is reduced after the formation of nitrite [35]. If the color turns red, then the bacteria does not reduce nitrate to nitrite. From Table 3, the DS15 isolate showed a negative reaction for the nitrate test. Generally, LAB has limited biosynthetic capacity [35], but requires complex nutrients such as amino acids, peptides,

nucleotide bases, vitamins, minerals and fatty acids [21]. The absence of carbohydrate fermentation during incubation does not mean bacterial isolates do not grow because bacteria can use other nutrients

contained in the media as an energy source. As peptone can be converted to amino acids which will be converted by oxidative deamination into ketoamino acids, the substrate will be further metabolized through the Krebs cycle for energy production [35]. The last test conducted to identify the identity of DS15 was gram stain. Gram stain is a differential staining performed to distinguish between gram-positive and gram-negative bacteria. Gram staining of DS15 showed a purplish-blue color, indicating that the isolate was a gram-positive bacterium (Fig. 1). For bacteria classified as gram negative, a pink

color will be produced. [Gram positive bacterial cell wall](#) is composed of [peptidoglycan](#) layer (90%) and other small molecules such as ketoic acid, teicuronic acid, polysaccharide, lipoteichoic acid, glycolipid and micolate acid [36]. While the gram-negative bacteria have more complex cell wall in the form of peptidoglycan layer (10%) and the outer membrane that protects peptidoglycan in the form of phospholipid and lipopolysaccharide [29], [36]. In gram-positive cells, the thickness of the peptidoglycan layer can withstand the purple Iodine crystals (CV-I) complex and the pore becomes smaller due to the dehydration properties of the alcohol so that the cells remain purple [29], [35].

TABLE III BIOCHEMICAL CHARACTERISTICS OF ISOLATE DS15

No	Characteristic Results
1	MRSA +
2	Arginine -
3	Aerobe/ Anaerobe A
4	Aesculine -
5	TSI M/M
6	Arabinose -
7	Catalase -
8	Raffinose -
9	Oxidase -
10	Sorbitol -
11	Motilities -
12	Trehalase -
13	Lactose -
14	Xylose/Maltose -
15	Glucose -
16	Dulcitol/ Melezitose -
17	Mannitol -
18	Nitrate -
19	VP -
20	Gram +

Fig. 1 Gram assay of isolate DS15 under a 1000 x magnification of light microscope. IV. CONCLUSION The research successfully identified 704 isolates from four different West Sumatera indigenous fermented foods. A total of 552 isolates were identified as LAB and 103 among them were able to produce GABA. The isolate DS15 could produce GABA of 49.365 mg/ml, while DS18 produced only 3.415 mg/ml. The isolate DS15 is characterized by its ability to grow on the MRSA, aerobic, do not use lactose, glucose, mannitol, arginine, aesculin, arabinose, raffinose, sorbitol, trehalase, and xylose/melezitose. Moreover, the isolate DS15 do not produce acidic substances, do not reduce nitrate, negative for catalase and oxidase. Based on gram stain reaction, the isolate DS15 was a gram-positive bacterium and classified as *Lactobacillus* sp. This study aims to obtain LAB isolates from West Sumatra fermented foods that can produce GABA as anti-stress. This

study helps researchers and industry to produce GABA production. Thus, new [lactic acid bacteria isolated from](#) West

Sumatra [fermented food](#) can be used as a GABA producer which is used as a supplement for feed or additional feed

for livestock as an anti-stress. ACKNOWLEDGMENT We are very grateful to the Minister of Research,

[Technology and Higher Education of the Republic of Indonesia for](#) funding this research through PMDSU No: No:

1387/E4./2015 [and the support through World Class Professor Program Scheme-B No. 123.57/D2.3/KP/2018.](#)

REFERENCES [1] Reis H.J., Paquet M., Guatimosim C., Palotas A. Neurotransmitters in the central nervous system & their implication in learning and memory processes. *Curr. Medic. Chemist.* 16: 796-840. 2009 <http://dx.doi.org/10.2174/092986709787549271> [2] Burton A.R., Baquet Z., Eisenbarth G.S., Tisch R., Smeyne R., Workman C.J., Vignali DAA. Central nervous system destruction mediated by glutamic acid decarboxylase-specific cd4+ t cells. *J Immunol.* 184: 4863-4870. 2010 <https://doi.org/10.4049/jimmunol.0903728> [3] Dhakal R., Baipai V.K., Baek K.H. Production of GABA (γ - Aminobutyric Acid) by microorganisms: a Review. *Braz. J Microbiol.* 43: 1230-1241. 2012. <http://dx.doi.org/10.1590/S1517-83822012000400001> [4] Esmaeili A., Ghaed K. GABAA receptors as novel drug targets for treatment of mental disorders. *J Paramedi. Sci.* 1: 50-61. 2010. ISSN 2008-496X [5] Suwanmanona K, Chuan HP. Isolating *Bacillus subtilis* and optimizing its fermentative medium for GABA and nattokinase production. *Cyta J Food.* 12: 282-290. 2014. <http://dx.doi.org/10.1080/19476337.2013.848472> [6] Dai S.F., Gao F., Zhang W.H., Song S.X., Xu X.L., Zhou G.H. Effects of dietary glutamine and gamma-aminobutyric acid on meat colour, pH, composition, and water-holding characteristic in broilers under cyclic heat stress. *British Poultry Sci.* 45: 471-481. 2012. <https://doi.org/10.1080/00071668.2012.719148> [7] Li Y.H., Li F., Liu M., Yin J.J., Cheng B.J., Shi B.M., Shan A.S. Effect of γ -aminobutyric acid on growth performance, behavior and plasma hormones in weaned pigs. *Canadian J Anim. Sci.* 95: 165-171. 2015. <https://doi.org/10.4141/cjas2013-148> [8] Zhao A., Hu Z., Wang X.. Metabolic engineering of *Escherichia coli* to produce gamma-aminobutyric acid using xylose. *Appl. Microbiol. Biotechnol.* 101: 3587-3603. 2017 <https://doi.org/10.1007/s00253-017-8162-3> [9] Higuchi T., Hayashi H., Abe K. Exchange of glutamate and gamma-aminobutyrate in a *Lactobacillus* strain. *J Bacteriol.* 179: 3362-3364. 1997. ISSN: 1098-5530 [10] Siragusa S., Angelis M.D., Cagno R.D., Rizzello C.G., Coda R., Gobbetti M.. Synthesis of γ -aminobutyric acid by lactic acid bacteria isolated from a variety of Italian cheeses. *Appl. Environ. Microbiol.* 73: 7283-7290. 2007. <http://doi.org/10.1128/AEM.01064-07> [11] Xue L.X., Yan X.C., Xin G.Z. Optimisation of fermentative parameters for GABA enrichment by *Lactococcus lactis*. *Czech J Food Sci.* 27: 433-442. 2009. ISSN 1750-3841 [12] Thwe S.M., Kobayashi T., Luan T., Shirai T., Onodera M., Sato N.H., Imada C. Isolation, characterization, and utilization of γ -aminobutyric acid (GABA)-producing lactic acid bacteria from Myanmar fishery products fermented with boiled rice. *Food Sci. Technol.* 77: 279-288. 2011. <https://doi.org/10.1007/s12562-011-0328-9> [13] Gupta A., Sharma N. In vitro characterization of lactic acid bacteria isolated from lasoda bari- a rare fermented food of Himachal Pradesh-India for potential probiotic attributes. *J Microbiol. Biotechnol. Food Sci.* 2017. <https://doi.org/10.15414/jmbfs.2017.6.6.1323-1328> [14] Ali F.W.O., Abdulmir A.S., Mohammed A.S., Bakar F.A., Manap Y.A., Zulkifli A.H., Saari N. Novel, practical and cheap source for isolating γ -aminobutyric acid-producing *Leuconostoc* NC5 bacteria. *Res. J Med. Sci.* 3: 146-153. 2009. ISSN 1815-9346 [15] Kook M.C., Cho S.C. Production of GABA (gamma amino butyric acid) by lactic acid bacteria. *Korean J Food Sci.* 33: 377-389. 2013. <https://doi.org/10.5851/kosfa.2013.33.3.377> [16] Spies J.R. Colorimetric procedures for amino acid methods in enzymology. *Meth. Enzym.* 76: 467-477. 1957. [https://doi.org/10.1016/S0076-6879\(57\)03417-5](https://doi.org/10.1016/S0076-6879(57)03417-5) [17] Gad G.F.M., Hamid A.M.A., Farag Z.S.H. Antibiotic resistance in lactic acid bacteria isolated from some pharmaceutical and dairy product. *Braz. J Microbiol.* 45: 25-33. 2014. ISSN 1678-4405 [18] Islam K.N., Akbar T., Akther F., Islam N.N. Characterization and confirmation of *Lactobacillus* spp. from selective regional yoghurts for probiotic and interference with pathogenic bacterial growth. *Asian J Biol Sci.* 9: 1-9. 2016. <https://doi.org/10.3923/ajbs.2016.1.9> [19] Harimurti S., Rahayu E.S., Nasroedin, Kurniasih. Lactic acid bacteria isolated from the Gastro-intestinal tract of chicken: Potential use as probiotic. *Anim Prod.* 9: 82-91. 2005. ISSN 1411-2017 [20] Julendra H., Suryania A.E., Istiqomaha L., Damayantia E., Anwara M., Fitriani N. Isolation of lactic acid bacteria with cholesterol- lowering activity from digestive tracts of Indonesia native chickens. *Media Peternakan.* 40: 35-41. 2017. <https://doi.org/10.5398/medpet.2017.40.1.35> [21] Khalid K. An overview of lactic acid bacteria. *Int. J Biosci.* 1: 1-13. 2011. ISSN 2222-5234 [22] Surono I.S., Hosono A. Antimutagenicity of milk cultured with lactic acid bacteria from dadih against mutagenic Teerasi. *Milchwissenschaft.* 51: 478-487. 1996 [23] Huda N., Ahmad R. Budu and tukai-endemic fermented fish products from West Sumatera. *INFOFISH Int.* 3. 2006. ISSN 15115976 [24] Montagnac J.A., Davis C.R.. Tanumihardjo S.A.

Nutritional value of Cassava for use as a staple food and recent advance for improvement, *Compr. Rev. Food Sci. Food Saf.* 8: 181-194. 2009. <https://doi.org/10.1111/j.1541-4337.2009.00077.x85> [25] Pato U., Surona I.S. Bile and acid tolerance of lactic acid bacteria isolated from tempoyak and their probiotic potential. *J Agricultur. Technol.* 9: 1849-1862. 2013. ISSN 1686-914 [26] Li H., Ting Q., Guidong H., Yusheng C. Production of gamma-aminobutyric acid by *Lactobacillus brevis* NCL912 using fed-batch fermentation. *Microbila Cell Factories.* 9: 85. 2010. <https://doi.org/10.1186/1475-2859-9-85> [27] Lee B.J., Kim J.S., Kang Y.M., Lim J.H., Lim Y.M., Kim Y.M., Lee M.S., Jeong M.H., Ahn C.B., Je J.Y. Antioxidant activity and gamma-aminobutyric acid (GABA) content in sea tangle fermented by *Lactobacillus brevis* BJ20 isolated from traditional fermented foods. *Food Chem.* 122: 271-276. 2010. <https://doi.org/10.1016/j.foodchem.2010.02.071> [28] Marlida Y., Harnentis, Nurmiati. Isolation and screening of lactic acid bacteria from dadih for glutamic Acid production as precursor of gamma-Amino Butyric Acid (GABA) induced heat stress in broiler. *Int. J Chem. Tech. Research.* 9: 534-540. 2016. ISSN 2455-9555 [29] Sunatmo T.I. *Mikrobiologi esensial Jilid 1.* Ardy Agency, Jakarta. 2009a. ISBN 979-9527-95-3 [30] Ismail Y.S., Yulvizar C., Mazhitov B. Characterization of lactic acid bacteria from local cow's milk kefir, *IOP Conf Ser: Earth Environ. Sci.* 130: 012019. 2018. <http://dx.doi.org/10.1088/1755-1315/130/1/012019> [31] Nur F., Hafsan, Paramitasari D. Isolasi dan identifikasi bakteri asam laktat berpotensi probiotik dari dangke susu sapi di Kabupaten Enrekang. *Jurnal Biotek.* 3: 52-66. 2015. ISSN 2354-9106 [32] Guessas B., Kihal M. Characterization of lactic acid bacteria isolated from Algerian arid zone raw goats' milk. *African J Biotechnol.* 3: 339-342. 2004. <http://dx.doi.org/10.5897/AJB2004.000-2062> [33] Adam M.R., Nout R. *Fermentation and Food Safety.* Aspen Publisher, Washington. 2001. ISBN 0-8342-1843-7 [34] Hemraj V., Diksha S., Avneet G. A review on commonly used biochemical test for bacteria. *Innovare J. Life Sci.* 1: 1-7. 2013 [35] Sunatmo T.I. *Eksperimen mikrobiologi dalam laboratorium.* Ardy Agency, Jakarta. 2009b ISBN 979-9527-63-5 [36] Kim S.J., Chang J., Singh M. Peptidoglycan architecture of gram-positive bacteria by solid-state NMR. *Biochimica Biophysica Acta (BBA)- Biomembranes.* 1848: 350-362. 2015. <https://doi.org/10.1016/j.bbamem.2014.05.031> 855 856 857 858 859 860