

Turnitin Originality Report

Processed on: 04-Feb-2020 11:06 PM +08

ID: 1251362397

Word Count: 5042

Submitted: 1

MZ 3 By Mardiaty Zain

8% match (publications)

[Novirman Jamarun, Mardiaty Zain, Arief ., Roni Pazla. "Effects of Calcium, Phosphorus and Manganese Supplementation During Oil Palm Frond Fermentation by Phanerochaete chrysosporium on Laccase Activity and in vitro Digestibility", Pakistan Journal of Nutrition, 2017](#)

Similarity Index	Similarity by Source	
23%	Internet Sources:	N/A
	Publications:	23%
	Student Papers:	N/A

8% match (publications)

[Dewi Febrina, Novirman Jamarun, Mardiaty Zain, Khasrad . "Effects of Calcium \(Ca\) and Manganese \(Mn\) Supplementation During Oil Palm Frond Fermentation by Phanerochaete chrysosporium on In vitro Digestibility and Rumen Fluid Characteristics", Pakistan Journal of Nutrition, 2016](#)

6% match (publications)

[Dewi Febrina, Novirman Jamarun, Mardiaty Zain, Khasrad . "Effects of Using Different Levels of Oil Palm Fronds \(FOPFS\) Fermented with Phanerochaete chrysosporium plus Minerals \(P, S and Mg\) Instead of Napier Grass on Nutrient Consumption and the Growth Performance of Goats", Pakistan Journal of Nutrition, 2017](#)

Pakistan Journal of Nutrition 15 (4): 352-358, 2016 ISSN 1680-5194 © Asian Network for Scientific Information, 2016 [Effects of Calcium \(Ca\) and Manganese \(Mn\) Supplementation During Oil Palm Frond Fermentation by Phanerochaete chrysosporium on In vitro Digestibility and Rumen Fluid Characteristics](#) Dewi Febrina¹, [Novirman Jamarun](#)², [Mardiaty Zain](#)² and [Khasrad](#)² ¹Faculty of Agriculture and Animal Science, State Islamic University of Sultan Syarif Kasim Riau, Indonesia ²Faculty of Animal Science, Andalas University, Padang, Indonesia **Abstract:** [The objectives of the study were to evaluate the effects of calcium and manganese supplementation during oil palm frond fermentation by Phanerochaete chrysosporium on in vitro digestibility and rumen fluid characteristics. This research used a Completely Randomized Design comprising two factors as treatments and each treatment was repeated three times. The factors were Ca dose \(1.000, 2.000 and 3.000 ppm\) and Mn dose \(50, 100 and 150 ppm\). The data were analyzed using analysis of variance to measure the differences among treatments; Duncan's Multiple Range Test was used to determine the significance of differences. The parameters measured were as follows: digestibility of dry matter \(%\), organic matter \(%\), Neutral Detergent Fibre \(NDF\) \(%\), Acid Detergent Fibre \(ADF\) \(%\), cellulose \(%\) and hemicellulose \(%\). The rumen fluid characteristics considered included the following: pH and the concentrations of NH₃ \(mM\) and Volatile Fatty Acid \(VFA\) \(mM\). The results indicated interactions between Ca and Mn in relation to the digestibility of dry matter, organic matter, NDF, ADF, cellulose, pH, NH₃ and VFA concentration of rumen fluid. The optimal combination of the minerals used were 2.000 ppm of Mn and 150 ppm of Ca; these values yielded the highest values of digestibility of dry matter \(41.914%\), organic matter \(40.990%\), NDF \(66.429%\), ADF \(64.396%\) and cellulose \(68.524%\) and the highest concentration of VFA \(117.302 mM\).](#) Key words: Oil palm frond, Phanerochaete chrysosporium, Ca, Mn **INTRODUCTION** Indonesia is the largest palm oil producer in the world and 10.956.231 hectares were planted with oil palms in 2014 (Ministry of Agriculture, 2014). Palm trees produce various agricultural by-products, such as the leaves and petiole, which are also termed fronds. Use of oil palm frond as an animal feed is very limited due to its high lignin content (30.18%) (Febrina et al., 2014; Zain et al., 2014), low dry matter digestibility (40%) (Kawamoto et al., 2001) and low energy value (4.9 and 5.6 MJ metabolizable energy (ME)/kg DM (Alimon, 2005; Zahari and Alimon, 2005). Lignocellulose is a macromolecular complex comprising [lignin, cellulose and hemicellulose](#). [The degradation of lignin is the key step in lignocellulose](#) transformation (Zeng et al., 2013). [Phanerochaete chrysosporium is white-rot fungus that can degrade all major components of plant cell walls, including cellulose, hemicellulose and](#) lignin (Aitken et al., 1989; Gusse et al., 2006) and is the most active ligninolytic organism described to date. Its ability to degrade lignin and a wide variety of aromatic compounds is due to a non-specific extracellular enzyme system including manganese peroxidase (MnP), lignin peroxidase (LiP) and laccases (Lac) (Lopez et al., 2007; Tang et al., 2006). Phanerochaete chrysosporium is the most intensively studied white-rot fungus due to its ability to degrade a wide range of organic substrates (Fragoieiro and Magan, 2005; Wen et al., 2009; Huang et al., 2010). Ligninolytic enzyme production can be increased by adding Ca and Mn salts (Brown et al., 1990). Extracellular enzymes can be added directly to the media or extracellular enzymes can be produced by altering the levels of transcription and translation (Baldrin, 2003). Wuyep et al. (2003) reported that

Mn²⁺ and Ca²⁺ ions can support the growth and extension of 2 Basidiomycetes mycelia: *Lentinus squarrolus* and *Psathyrella atroumbonata*; the addition of 2.400-3.200 ppm of Ca stimulated growth and the optimal activity was obtained using *Lentinus squarrolus* dan *Psathyrella atroumbonata*; addition of Mn salts into the substrate improved the degradation of lignin and dry matter digestibility. Supplementation with Ca salts (Febrina, 2014) during palm frond biodelignification using *Phanerochaete chrysosporium* decreased lignin content by up to 27.34% and by 29.89% when Mn salts were added (Febrina et al., 2014). [The addition of 2.000 ppm Ca and 150 Mn salts to palm frond fermentation using Phanerochaete chrysosporium](#) resulted in low lignin contents of 22.4%. Corresponding Author: Dewi Febrina, Faculty of Agriculture and Animal Science, State Islamic University of Sultan Syarif Kasim Riau, Indonesia 352 and 25.77%, respectively (a decrease in lignin content from 30.18 to 22.4%) (Febrina, 2015). The purpose of [this study was to evaluate the effects of calcium and manganese supplementation during oil palm frond fermentation using Phanerochaete chrysosporium](#) in vitro. MATERIALS AND METHODS [Oil palm frond \(OPF\) fermentation: The OPF substrate was cut, dried and finely milled. Ca was used as CaCl₂ and Mn was used as MnSO₄.H₂O. Phanerochaete chrysosporium was maintained on Potato Dextrose Agar \(PDA\), slanted at 4°C and transferred to PDA plates at 37°C for 6 days; the fungus was subsequently grown on rice bran. The fermentation process was initiated by adding water to the OPF to a water content of 70%; Ca and Mn were then added according to the various treatments. After 10 days, all oil palm fronds were covered by white rot fungus and samples were taken for proximate analysis and fibre fraction determination. The proximate components were determined as described by AOAC \(1995\) using Foss equipment. The predominantly fibre residues \(hemicellulose, cellulose and lignin\) were determined according to the method of Van Soest et al. \(1991\) using Foss Fibretec. In vitro digestibility assay: In vitro digestibility was analyzed as described by Tilley and Terry \(1969\). Buffalo rumen fluid was diluted using McDougal buffer \(1: 2\) and dispensed into a 1 g substrate-prepared incubation tube, which was purged with CO₂ to maintain anaerobic conditions. The tubes were incubated in a water bath at 39°C for 24 h. After fermentation, the Erlenmeyer tube containing the sample was inserted into ice water to stop the fermentation. All samples were then centrifuged at 1.200 rpm for 15 min. The pH, NH₃-N and total VFA of the supernatants were then recorded. NH₃-N concentration was determined using microdiffusion Conway method and the total VFA concentration was measured using the steam distillation method. The previously incubated samples were vacuum-filtered \(Whatman no. 41\) and dried at 60°C in an oven. The dried samples were used to analyze the dry matter, organic matter, NDF, ADF, hemicellulose and cellulose in an in vitro digestibility assay. Experimental design and statistical analysis: The study implemented a Completely Randomized Design using a 3 x 3 factorial pattern and 3 repetitions. The first factor was the level of Ca \(1.000, 2.000 and 3.000 ppm\) and the second factor was the level of Mn \(50, 100 and 150 ppm\). The significance of differences among the treatments were then analyzed using Duncan's Multiple Range Test. RESULTS AND DISCUSSION In this study, the interaction between Ca and Mn affected \(p<0.05\) the in vitro digestibility of dry matter, organic matter, NDF, ADF and cellulose. Supplementation with Mn significantly \(p<0.05\) affected the digestibility of hemicellulose \(Table 1\). The lowest dry matter digestibility \(34.038%\) appeared in experiment A1B2 \(1.000 ppm Ca and 100 ppm Mn\). This indicated that the combination of Ca and Mn was not optimal for supporting the growth and mycelial increase of the fungus. The fungus did not produce optimal amounts of enzyme to degrade lignin; therefore, large amounts of cellulose and hemicellulose remained attached to the lignin. Lignin degrades with difficulty into fractions \(Lynch, 1993\); lignin bonds to cellulose and hemicellulose, forming lignocellulose and lignohemicellulose, which rumen microbes find difficult to degrade. Cleavage of the lignocellulosic bonds facilitates access to the substrate by hydrolytic enzymes such that microbes can degrade and thus decrease lignin contents; according to Sun and Cheng \(2002\), breakdown of the crystal structure of cellulose facilitates access to the substrate by hydrolytic enzymes. Increasing the concentration of Ca from 1.000 ppm to 2.000 ppm and the concentration of Mn from 100 ppm to 150 ppm \(A2B3\) stimulated the growth of fungal mycelia, indicating that the production of ligninase was increased. Ligninase belongs to a group of isoenzymes, oxidases and peroxidases that depolymerize lignin \(Chahal and Chahal, 1998\). When more enzymes are produced, lignocellulose and lignohemicellulose become more stretchable. The cellulose and hemicellulose can then be separated and utilized by rumen microbes as a source of energy. Optimal microbe growth, as achieved through appropriate nutrient supplementation, will produce the highest dry matter digestibility. The highest dry matter digestibility was found in treatment A2B3 and was associated with the optimal degradation of lignin by *Phanerochaete chrysosporium* \(this treatment resulted in the lowest lignin content of 22.40%\). Low lignin content \(indicating greater lignin degradation\) leads to the digestion of more nutrients and produces the highest dry matter digestibility. Lignin degradation causes the stretching of lignocellulose and lignohemicellulose such that the cellulose and hemicellulose can be utilized by fungi mycelia for growth. Suparjo \(2010\) suggested that mycelial multiplication is an indicator of growth and shows an increase in the of dry matter content. Ballgees et al. \(2009\) also described the increase of dry matter digestibility of bagasse by fermentation using white rot fungi. The addition of Mn to the substrate increases the degradation of lignin \(Kerem and Hadar, 1997\) and the dry matter digestibility of substrates \(Kerem and Hadar, 1995\). Dry matter with high digestibility is considered appropriate for feeding to livestock \(Alwi et al., 2013\). Decreased lignin content was found to be positively correlated with the in vitro dry matter digestibility of bagasse when fermented using *L. edodes* and *C. subvermispora* \(Okano et al., 2006\). The organic matter digestibility of oil palm frond was 24.34% before fermentation and increased to 24.81- 40.61% after fermentation, ranging from 34.396- 40.990%. This result indicates that the addition of Ca and Mn affected the organic matter digestibility of the fermented oil palm fronds. Muhktarudin and Liman \(2006\) found that supplementing with organic macro minerals in feed rations increased the organic digestibility. Organic matter digestibility follows the trends seen with dry matter because low dry matter](#)

digestibility causes low organic matter digestibility. Muhktarudin and Liman (2006) reported that increased dry matter digestibility results in increased organic matter digestibility. Elihasridas (2012) reported that increased dry matter digestibility, which was caused by increased supplementation of Zn salts in rations of ammoniated corn cobs, was associated with increased dry matter digestibility. The lowest digestibilities of NDF, ADF and cellulose were observed in treatment A1B2 (57.29, 50.86 and 50.04%, respectively). The low digestibilities of NDF, ADF and cellulose resulted from the high contents of NDF and ADF, indicating that the lignin was still bound to cellulose and hemicellulose. According to Moore and Jung (2001), the lignin in plant cell walls reduces [the digestibility of cellulose and hemicellulose](#) by rumen microbes; [lignification process limits the ability of microbes to break down cellulose and hemicellulose as an energy source for growth](#) (Tripathi et al., 2008). The increased NDF digestibility observed for treatment A2B3 indicates that the additions of Ca and Mn were able to increase the amount of fungal mycelium and the production of lignolytic enzymes, thereby cleaving the bonds within lignocellulose and lignohemicellulose. Hammel (1997) suggested that Phanerochaete chrysosporium can degrade lignin; Kerem and Hadar (1997) suggested that the addition of Mn to the substrate would increase lignin degradation. Lignin biodegradation [is a key process](#) involved in [lignocellulosic waste composting; the increased use of carbon](#) will increase [the activity of microorganisms that increase the activity of lignolytic enzymes, thus increasing lignin degradation](#) (Feng et al., 2011). Hemicellulose digestibility increased as the Mn concentration increased from 100 to 150 ppm. Hemicellulose digestibility increased as the addition of Mn increased, indicating that Mn affected lignin degradation by affecting the production of ligninolytic enzymes. Baldrin (2003) suggested that Mn plays a role in the Manganese Peroxidase (MnP) reaction cycle. The fragmentation of lignocellulosic bonds and lignohemicellulose caused hemicellulose to become an energy source; therefore, the digestibility of hemicellulose by rumen microbes was increased. MnP activity and stability are influenced by the concentration of Mn²⁺ (Urek and Pazarlioglu, 2005); Mn²⁺ is a specific effector that encourages the production of MnP (Zhao et al., 1996). Characteristics of rumen fluid: pH affects the activity of microorganisms in the rumen and greatly affects the growth and activity of rumen microorganisms. Rumen pH is altered by the production of acid by microorganisms during fermentation in the rumen. Table 2 shows the effects of Ca and Mn on [pH and the concentrations of NH₃ \(mM\) and VFA \(mM\)](#) in fermented oil palm fronds. Rumen pH ranges from 6.667 to 6.867. This range is suitable for the growth of rumen microbes. This pH value indicates a balance between buffering capacity and the acid or alkaline fermentation products, VFA and N-NH₃ (Arora, 1995). Erdman (1988) found that the optimal pH range for cellulose digestion in the rumen is 6.4 to 6.8; when the rumen pH falls below 6.2, cellulolytic bacteria are disrupted, causing decreased fibre digestibility. The highest rumen pH (6.867) in this study was found in treatment A1B1 (1 [.000 ppm Ca and 50 ppm Mn](#)). Increased doses of [Ca and Mn significantly \(p<0.05\)](#) lowered rumen pH. Supplementation of 3 [.000 ppm Ca and 150 ppm Mn](#) (A3B3) resulted in [the](#) lowest rumen pH (6.667). Rumen pH decreased as the Ca and Mn doses increased. [Similar results were reported by](#) Suyitman [et al.](#) (2013), [who](#) noted that the effects of various treatments (steam, ammoniation and ensilage) were highly significant (p<0.01) on rumen fluid pH. The concentration of NH₃ in this study ranged from 5.523 mM to 8.365 mM. This value is sufficient for the optimal growth of rumen microbes. The concentration of ammonia required to support microbial growth is 6-21 mM (McDonald et al., 2010). The concentration of NH₃-N depends on protein intake. Higher protein intake will stimulate higher microbial activity, which increases NH₃-N concentration. The lowest NH₃, which occurred in response to [the addition of 3 .000 ppm Ca and 100 ppm Mn](#) (treatment A3B2), caused [the](#) sub optimal production of ligninases. Therefore, lignin degradation was also suboptimal. [The addition of 3 .000 ppm Ca and 100 ppm Mn](#) (treatment A3B2) resulted in lower crude protein digestibility. The low concentrations of NH₃ found in this treatment were associated with low nutrient digestibility. Low digestibility also produced the lowest concentration of NH₃ in this study. Low concentrations of NH₃ in the rumen limit the activity of rumen fermentation (Fariani et al., 2015). The highest NH₃-N concentration found in this study was observed for treatment A2B1 (2.000 ppm Ca and 50 ppm Mn) and was associated with higher protein digestibility. Proteins degraded in the rumen are used to form NH₃ and N, which are used by rumen microbes to

Table 1: Nutrient Digestibility resulting from oil palm frond fermentation by Phanerochaete chrysosporium while supplemented with Ca and Mn (%) -----
----- Mn (ppm) ----- Ca (ppm)

	B1 (50)	B2 (100)	B3 (150)	Average	Dry matter	A1 (1.000)	37.151±0.842c	34.038±0.712d
	38.643±0.191b	36.610c	A2 (2.000)	38.928±0.244b	36.016±0.631c	41.914±1.031a	38.953a	A3 (3.000)
	38.653±0.911b	36.408±0.940c	38.920±0.191b	37.994b	Average	38.244b	35.487c	39.826a
	Organic matter	A1 (1.000)	35.727±1.521cd	32.375±0.512e	37.395±0.626bc	35.166b	A2 (2.000)	37.296±0.931bc
	34.396±1.041d	40.990±1.581a	37.560a	A3 (3.000)	37.787±0.559b	34.798±1.037d	37.461±0.690bc	36.682a
	Average	36.937b	33.856c	38.616a	NDF	A1 (1.000)	58.332±0.84cd	57.293±0.55d
	64.225±1.78ab	59.946	A2 (2.000)	61.803±4.40abcd	61.018±3.98abcd	66.429±0.71a	63.083	A3 (3.000)
	60.402±4.88bcd	63.378±4.20abc	64.303±1.57ab	62.695	Average	60.176b	60.563b	64.986a
	ADF	A1 (1.000)	56.535±0.92b	50.863±5.79c	60.131±2.37ab	55.843b	A2 (2.000)	63.822±0.94a
	57.553±0.37b	64.396±0.72a	61.924a	A3 (3.000)	59.836±2.24ab	60.513±3.60ab	61.257±2.94ab	60.535a
	Average	60.065a	56.31b	61.928a	Cellulose	A1 (1.000)	56.849±0.922ab	50.044±9.046b
	66.153±3.477a	57.682	A2 (2.000)	65.780±5.003a	62.281±9.603ab	68.524±3.777a	65.528	A3 (3.000)
	56.705±1.792ab	53.152±17.832ab	63.715±6.303ab	57.857	Average	59.778ab	55.159b	66.131a
	Hemicellulose	A1 (1.000)	65.25±6.172	75.87±19.554	83.62±18.314	74.915	A2 (2.000)	55.13±16.459
	71.53±14.334	73.89±1.744	66.851	A3 (3.000)	60.06±26.956	73.51±8.158	76.74±14.317	70.102
	Average	60.148b	73.638ab	78.082a	Description:	Different superscripts in the same row and column indicate significant differences (p<0.05)		

Table 2: Characteristics of rumen fluid in response to oil palm fronds that were fermented by Phanerochaete chrysosporium in the presence

of Ca and Mn supplementation ----- Mn (ppm) -
 ----- Ca (ppm) B1 (50) B2 (100) B3 (150)
 Average pH A1 (1.000) 6.867±0.057a 6.833±0.057ab 6.866±0.057a 6.856a A2 (2.000)
 6.833±0.057ab 6.833±0.057ab 6.800±0.100ab 6.822a A3 (3.000) 6.700±0.100ab
 6.700±0.173ab 6.667±0.152b 6.689b Average 6.800 6.789 6.778 NH₃ (mM) A1 (1.000)
 7.964±0.113a 5.557±0.212c 8.342±0.178a 6.856a A2 (2.000) 8.365±0.252a 5.528±0.131c
 8.266±0.273a 6.822a A3 (3.000) 6.674±0.287b 5.348±0.253c 6.669±0.276b 6.689b Average
 6.800 6.789 6.778 VFA (mM) A1 (1.000) 113.556±16.730a 71.873±4.486b 111.971±3.041a
 99.134 A2 (2.000) 113.928±0.422a 67.119±10.230b 117.302±5.120a 99.450 A3 (3.000)
 107.064±12.990a 63.067±5.864b 111.178±6.505a 93.769 Average 111.516a 67.353b 113.484a

Description: Different superscripts in the same row and column indicate significant differences (p<0.05) support their growth. The more proteins are degraded, the more NH₃ is formed and the greater the growth of the microbes. Ali [et al. \(2015\)](#) [reported that an increase in the concentration of NH₃ was associated with the degradation of a protein-rich substrate.](#) The ability of [rumen microbes to degrade feed](#) affects [the concentration of VFA.](#) [The concentration of VFA was 63.067 and 117.302 mM.](#) According to the recommendations of Waldron et al. (2002), the total VFA concentration in the rumen was 60-120 mM. The interaction of [2.000 ppm Ca and 150 ppm Mn](#) produced [the highest concentration of VFA \(117.302 mM\).](#) This value was similar to that obtained using some other treatments [but was significantly different \(p<0.05\)](#) and higher compared to treatments A2B2 and A2B3. The highest VFA found for treatment A2B3 ([2.000 ppm Ca and 150 ppm Mn](#)) suggests that using [the appropriate combination of Ca and Mn can optimize the production of ligninolytic enzyme by Phanerochaete chrysosporium.](#) Phanerochaete chrysosporium produces haeme peroxidase, lignin peroxidase (LiP) and manganese peroxidase (MnP), which degrade the lignin that binds cellulose and hemicellulose together. LiP is a major catalyst in lignolysis by the fungus because it can degrade non-phenolic compounds that comprise 90% of the lignin structure (Srebotnik et al., 1994). The lowest VFA was found for treatment A3B2 (3.000 ppm Ca and 100 ppm Mn), which is assumed to represent the lowest nutrient digestibility. Organic matter digestibility and dry matter correlate with the production of VFA (Liu et al., 2002); high in vitro digestibility of organic matter and crude fibre increase the total production of VFA in the rumen (Sudekum et al., 2006). Conclusion: Supplementation of [2.000 ppm Ca and 150 ppm Mn](#) provided a better result than the other tested treatments. The in-vitro digestibilities of the nutrients were as follows: [dry matter, 41.914%; organic matter, 40.990%; NDF, 66.429%; ADF, 64.396% and cellulose, 68.524%.](#) [The highest value of VFA obtained was 117.302 mM.](#) [ACKNOWLEDGMENTS This study was partially supported by the DP2M Directorate General of Higher Education, Republic of Indonesia. The authors are very grateful to the Directorate General of the Department of Higher Education, National Education of the Republic of Indonesia, which funded this experiment through the "Hibah Pasca" grant program in 2015.](#)

REFERENCES Aitken, M.D., R. Venkatadri and R.L. Irvine, 1989. Oxidation of phenolic pollutants by a lignin degrading enzyme from the white-rot fungus Phanerochaete chrysosporium. Water Res., 23: 443-450. Ali, A.I.M., S. Sandi, Riswandi, A. Imsya, A. Prabowo and N. Rofiq, 2015. Evaluation of Yeast Supplementation with Urea-Molasses in Rice Straw-Based Diets on in vitro Ruminal Fermentation. Pak. J. Nutr., 14: 988-993. Alimon, A.R., 2005. The nutritive value of palm kernel cake for animal feeds. Palm Oil Developments. vol. 40. Malaysian Palm Oil Board. Kuala Lumpur, Malaysia, pp: 12-14. Alwi, M., S. Wardhana and F.M. Suhartati, 2013. Sugarcane baggase fermentation using Phanerochaete chrysosporium as effort to increase rumen fermentation products in vitro. J. Ilmiah Peternakan, 1: 479-487. AOAC, 1995. Official Methods of Analysis of the Association of Official Analytical Chemists. Association of Official Analytical Chemists. Washington, D.C. Arora, S.P., 1995. Microbial digestion in the ruminant. Translation by Retno Murwani. Gadjah Mada University Press, Yogyakarta. Baldrin, P., 2003. Interaction of heavy metal with white- rot fungi. Enzyme Microb. Technol., 32: 78-91. Ballgees, A., Atta Ellman, A.M.A. Fadal Elsseed and A.M. Salih, 2009. Effects of Albazia lebeck or wheat bran supplementation on intake, digestibility and rumen fermentation of ammoniated baggase. J. Appl. Sci. Res., 5: 1002-1006. Brown, J.A., J.K. Glend and M.H. Gold, 1990. Manganese regulate expression of manganese peroxidase by Phanerochaete chrysosporium. J. Bacteriol., 172: 3125-3130. Chahal, P.S. and D.S. Chahal, 1998. Lignocellulosic waste: biological conversion. In: Martin AM, editor. Bioconversion of Waste Materials to Industrial Products. Ed ke-2. London: Blackie Acad Prof., pp: 376-422. Elihasridas, 2012. The effect of supplementation of mineral Zinc on in vitro digestibility of amoniated corn cobs ration. J. Peternakan, 9: 9-14. Erdman, R.A., 1988. Dietary buffering requirements of lactating dairy cows. A Review. J. Dairy Sci., 71: 246- 3246. Fariani, A., A. Abrar, G. Muslim and L. Warly, 2015. Supplementation of fermented palm press fibre on digestibility of rice straw and rumen bacteria profile. Pak. J. Nutr., 14: 80-83. Febrina, D., 2014. Biological delignification by Phanerochaete chrysosporium with addition of mineral Ca and its effect on nutrient content of oil palm frond. Research Report. Institute for Research and Community Service. State Islamic University Sultan Syarif Kasim. Riau. Pekanbaru-Indonesia. (unpublished). Febrina, D., 2015. Utilization of palm fronds delignification products using Phanerochaete chrysosporium as substitute forage on ruminant feed. Research Report. Faculty of Animal Science. University of Andalas. Padang. Indonesia. (unpublished). Febrina, D., J. Novirman, Z. Mardiati, Khasrad and M. Rini, 2014. Biological delignification by Phanerochaete chrysosporium with addition of mineral Mn and its effect on nutrient content of oil palm frond. The 16th AAAP Animal Science Congress November 10-14, 2014. Yogyakarta. Indonesia, pp: 1.723-1.726. Feng, C.L., G.M. Zeng, D.L. Huang, S. Hu, M.H. Zhao, C. Lai, C. Huang, Z. Wei and N.J. Li, 2011. Effect of ligninolytic enzymes on lignin degradation and carbon utilization during. Lignocellulosic waste composting. Proc. Biochem., 46: 1515-1520. Fragoeiro, S. and N. Magan, 2005. Enzymatic activity, osmotic stress and degradation of pesticide mixtures in soil extract liquid broth inoculated with Phanerochaete chrysosporium and Trametes versicolor. Environ. Microbiol., 7: 348-355. Gusse, A.C., P.D. Miller and T.J. Volk, 2006. White-rot fungi demonstrate first biodegradation of

