

Turnitin Originality Report

Processed on: 04-Feb-2020 11:05 PM +08

ID: 1251362425

Word Count: 4344

Submitted: 1

MZ 4 By Mardiaty Zain

8% match (publications)

[Novirman Jamarun, Mardiaty Zain, Arief .., Roni Pazla. "Effects of Calcium \(Ca\), Phosphorus \(P\) and Manganese \(Mn\) Supplementation during Oil Palm Frond Fermentation by Phanerochaete chrysosporium on Rumen Fluid Characteristics and Microbial Protein Synthesis", Pakistan Journal of Nutrition, 2017](#)

Similarity by Source	
Similarity Index	
20%	
Internet Sources:	N/A
Publications:	20%
Student Papers:	N/A

7% match (publications)

[Dewi Febrina, Novirman Jamarun, Mardiaty Zain, Khasrad .. "Digestibility of Goat Rations Containing Fermented Oil Palm Fronds by Phanerochaete chrysosporium Supplemented With Phosphorus, Sulfur and Magnesium", Journal of Biological Sciences, 2017](#)

5% match (publications)

[Dewi Febrina, Novirman Jamarun, Mardiaty Zain, Khasrad .. "The Effects of P, S and Mg Supplementation of Oil Palm Fronds Fermented by Phanerochaete chrysosporium on Rumen Fluid Characteristics and Microbial Protein Synthesis", Pakistan Journal of Nutrition, 2016](#)

Pakistan Journal of Nutrition 15 (3): 299-304, 2016 ISSN 1680-5194 © Asian Network for Scientific Information, 2016 The [Effects of P, S and Mg Supplementation of Oil Palm Fronds Fermented by Phanerochaete chrysosporium on Rumen Fluid Characteristics and Microbial Protein Synthesis](#) Dewi Febrina¹, [Novirman Jamarun](#)², [Mardiaty Zain](#)² and [Khasrad](#)² ¹Faculty of [Agriculture and Animal Sciences, State Islamic University Sultan Syarif Kasim II, Riau, Indonesia](#) ²Faculty of Animal Science, Andalas University, Padang, Indonesia **Abstract: The objective of this study was to evaluate the effects of sulfur, phosphorus and magnesium supplementation of oil palm fronds fermented by Phanerochaete chrysosporium on rumen fluid characteristics and microbial protein synthesis. This research was carried out using a randomized block design with 4 treatments and 4 replications. The following treatments were performed: A = fermented oil palm fronds (FOPFs); B = FOPFs+0.4% P; C = FOPFs+0.4% P+0.3% S and D = FOPFs+0.4% P+0.3% S+0.1% Mg. The data were subjected to analysis of Variance (ANOVA) and differences between treatments were tested using Duncan's Multiple Range Test (DMRT). The observed parameters were microbial protein synthesis and rumen fluid characteristics. Supplementation of FOPFs with P, S and Mg resulted in the highest microbial protein synthesis and VFA concentration.** Key words: Oil palm frond, Phanerochaete chrysosporium, P, S, Mg **INTRODUCTION** Oil palm fronds (OPFs) are one of the main by-products of the palm oil industry and have great potential as a source of roughage or as a component of a complete feed for ruminants (Zahari et al., 2003). OPFs are limited as a feedstuff due to their high lignin content (30.18%) (Febrina et al., 2014; Zain et al., 2014). The optimization of OPFs as feed has focused on efforts to reduce their lignin content. Related physical, chemical, biological or physical-chemical-biological treatments aim to break lignocellulose-lignocellulose bonds. However, the use of chemicals pollutes the environment and reduces the ability of microorganisms to degrade lignin by inhibiting their ability to break lignocellulosic bonds. Lignin is a natural phenolic polymer that is commonly found in plant cell walls and it is a limiting factor in animal feed due to its low digestibility (Imsya et al., 2013). Lignin is degraded by Basidiomycete fungi, which are its most efficient digesters (Rabinovich et al., 2004) and the species Phanerochaete chrysosporium has been the most extensively studied due to its ability to degrade a wide range of organic substrates (Dorado et al., 1999; Fragoeiro and Magan, 2005; Wen et al., 2009; Yu et al., 2009). Phanerochaete chrysosporium strains simultaneously degrade cellulose, hemicellulose and lignin, whereas other species (such as Ceriporiopsis subvermisporea) tend to degrade lignin prior to cellulose and hemicellulose (Sanchez, 2009). Phanerochaete chrysosporium growth affects the availability of minerals in the substrate. The biodelignification of OPFs using Phanerochaete chrysosporium degrades 27.34% and 29.89% of lignin when Ca (Febrina, 2014) and Mn (Febrina et al., 2014) are added, respectively. Improving the quality of OPFs through a biodelignification process should be integrated with the optimization of rumen bioprocessing by increasing the population of rumen microbes; the digestibility of the fibre is dependent on rumen microbial activity (Zain et al., 2010a,b,c). Increasing the digestibility of the fibre should be approached in terms of providing the nutrients required for rumen microbial growth (Leng, 1990). Microbial protein synthesis can be optimized through the addition of these components. The elements P, S and Mg have low bioavailability but are essential for microbial growth; additionally, these elements are often deficient in fibrous low-quality feed (Preston and Leng, 1987; Komisarczuk and Durand, 1991;

Little, 1986). Phosphorus is important for the metabolism and maintenance of cell membranes and cell walls and is a component of nucleic acids and high-energy molecules (ATP, ADP and AMP) (Komisarczuk and Durand, 1991; Bravo et al., 2003; Rodehutsord et al., 2000; Zain et al., 2010a,b,c). Sulfur supports the formation of sulfur-containing amino acids (Slyter et al., 1986) and aids in the synthesis of microbial proteins (Zain et al., 2010a,b,c), vitamins (thiamine and biotin) and coenzyme A (CoASH) (Komisarczuk and Durand, 1991). Sulfur also supports crude fibre [degradation in the rumen](#) and [the stimulation of cellulolytic bacterial growth](#) (Bal and Ozturk, 2006). Magnesium is important Corresponding Author: Dewi Febrina, [Faculty of Agriculture and Animal Sciences, State Islamic University Sultan Syarif Kasim II, Riau, Indonesia](#) 299 for microbe growth and functions in enzyme systems (Little, 1986; Komisarczuk and Durand, 1991); however, magnesium is often deficient in low-quality forage (Leng, 1990). [The biodelignification of OPFs with Phanerochaete chrysosporium plus the elements Ca and Mn can break down lignocellulose and lignin hemicellulose bonds. P, S and Mg were added to](#) fermented oil palm fronds (FOPFs) to optimize the bioprocessing of the rumen. This study aimed to determine the effects of P, S and Mg on fermented OPFs and their effects on microbial protein synthesis and the characteristics of rumen fluid. MATERIALS AND METHODS OPF fermentation: The OPFs used in this study were cut, dried and finely milled. Ca was obtained from CaCl₂ and Mn was obtained from MnSO₄.H₂O. Phanerochaete chrysosporium was maintained on potato dextrose agar (PDA) slants at 4°C, transferred to PDA plates at 37°C for 6 days and subsequently grown on rice bran. The fermentation process was initiated by adding water to the OPFs until the water level reached 70%; either Ca or Mn was then added depending on the treatment. Rumen fluid characteristics: Rumen Fluid Characteristics (ruminal [pH](#), [NH₃-N](#) and [VFA](#) concentration). [The ruminal pH was measured using a pH meter. The](#) concentration of ammonia [-N was](#) determined via [the](#) Conway [microdiffusion method. VFA was](#) analyzed [using the steam distillation method](#). Buffalo [rumen](#) liquor was filtered through 4 layers of cheesecloth. One gram of sample was added to the 100-mL incubation tube containing 50 mL of mixed solution. Before the incubation tube was closed, it was purged with CO₂ gas for 30 sec and then incubated for 48 h. When each incubation had finished, 2 drops of HgCl₂ were added to the cultures. The incubation tubes [were then centrifuged at 4000 rpm for 10 min. The](#) supernatants were used for the determinations [of pH, NH₃-N](#) concentration [and total VFA; the residue was](#) added to 50 ml of a 0.2% pepsin HCl solution and incubated for 48 h. The residue was then filtered using Whatman no. 41 filter paper and dried at 60°C for 48 h. [Rumen protozoa population: Measurements were carried out using a counting chamber. Up to 0.5 mL of rumen solution was fixed with 0.5 mL methyl green formalin saline \(MFS\) solution in tubes and thoroughly](#) mixed (Ogimoto and Imai, 1981). [Up to 0.1 ml of this sample was](#) dropped [on the counting chamber](#) (haemocytometer) [using a pipette and a coverslip was placed over the sample. The protozoa were](#) quantified [on the counter under a microscope at 40x magnification](#). According to [the number of protozoa obtained by the above counting procedure, the number per 1 mL of rumen contents can be calculated using the following](#) formula: Protozoa population/mL = (1/0.1 x 0.065 x 5 x 16) x n x d [where n = the number of protozoa in the counting chamber and d = the sample dilution](#). Total bacterial [population](#): Brain-heart infusion (BHI) medium consisting of BHI powder, glucose, cellulobiose, starch, cysteine, hemin and resazurin was used to calculate the total bacterial population. It was heated until its colour changed from yellowish brown to reddish brown and back to yellowish brown. The substance was then cooled and the CO₂ was drained. Anaerobic BHI medium (4.9 mL) was added to Hungate tubes filled with 0.150 g of bacto. Using a pipette, 0.05 mL sample volumes of rumen fluid that had undergone treatment and incubations of 4 h were added to the diluent media. The following dilution process was used: 0.05 mL of bacterial culture was added to 4.95 mL of diluent media and then 0.05 mL of the diluted bacterial culture was added to 4.95 mL of diluent media; this process was continued until dilutions of 10⁻², 10⁻⁴, 10⁻⁶ and 10⁻⁸ were achieved. Volumes of 0.1 ml of each dilution were removed from the tube, added to the media, rotated on a roller drum. The bacteria were then incubated for 24 h. No. of colonies Bacteria/mL = 0.05 x 0.1 x 10 where x is the tube's serial dilution to 10Gx [Experimental design and statistical analysis: The](#) research [was carried out using a randomized block design](#) (4 x 4) [with four replications \(the addition of P, S and Mg as treatments\). Differences between treatments were analyzed using Duncan's Multiple Range Test](#). The following treatments were carried out: A = FOPFs without the addition of minerals (control) B = FOPFs+P (0.4% DM) C = FOPFs+P (0.4% DM)+S (0.3% DM) D = FOPFs+P (0.4% DM)+S (0.3% DM)+Mg (0.1% DM) RESULTS AND DISCUSSION The influence of [P, S and Mg](#) on [palm frond fermented by the fungus Phanerochaete chrysosporium](#) and their effects [on rumen fluid characteristics](#) are shown in Table 1. The [addition of P, S and Mg](#) had no effect (p>0.05) on the rumen fluid, i.e., 6.80-6.87 is a normal pH for optimal rumen microbial activity. The same [results were reported by](#) Suyitman [et al.](#) (2013), who found that the pH of rumen fluid was 6.60-6.84 when S and P were supplemented during the ammonization of palm leaves. Van Soest (1994) determined that the optimal pH for the maintenance of cellulolytic activity was 6.6±0.5; Keidane and Birgele (2003) found that the pH ranges for the optimal activities of proteinases plus peptidases, cellulases and deaminases were 5.5-7.0, 6.2-7.0 and 6.5-7, respectively; they also found that the optimal pH for VFA synthesis was 6.0-8.0. Chiba (2009) determined that the rumen pH was 6-7 and Kamra (2005) determined that a pH of 6-6.9 was optimal for rumen microbial growth. The pH values of all treatments were nearly the same [due to the balance between VFA and NH₃](#) in [rumen fluid](#). The rumen [pH](#) represents a balance between its buffering [capacity and the acidity or alkalinity of fermentation products](#) (VFA and NH₃) and pH is affected by the products of fermentation (such as NH₃-N). In Church's study, the pH value was directly proportional to the concentration of NH₃-N (Church, 1988). The VFA concentration was lowest after treatment A at 95.791 mM, which indicates a low fermentation ratio. Supplementation with P, P and S and P, S and Mg ([treatments B, C and D](#), respectively) increased [the](#) concentration of [the](#) total VFA in the rumen fluid; this was also reported by Zain et al. (2010a,b,c). These findings indicate that mineral supplementation

can improve the growth and proliferation of rumen microbes, thereby increasing fermentation. VFA production is affected by the type and amount of forage, the rumen pH and the rate of passage through the rumen (Peters et al., 1989); the type of carbohydrate, the physical form of the feed, intake, the feeding frequency and the use of additives (France and Dijkstra, 2005) and the substrate, rumen microbial populations and ecology (Bannink et al., 2008). Treatment D (supplementation with P, S and Mg) produced the highest concentration of VFA at 111.007 mM. Supplementation with P, S and Mg thus encourages rumen microbe growth. The nutrient content of FOPFs is not sufficient for rumen microbes and supplementation with P, S and Mg encourages optimal growth. In the rumen, sulfur is critical for the synthesis of S-containing amino acids used in microbial protein synthesis and livestock are frequently deficient in S. P supplementation is essential for the animal and the microbes and sufficient Mg is a precondition for the optimal utilization of low-quality forages (Leng, 1990; Zain et al., 2010a,b,c). P deficiency is widespread in tropical countries (McDowell et al., 1983). Normal levels of Ca, P, Mg and Mo in the rumen increase the digestion of fibre (especially cellulose) via microbial activity. This increases the digestibility of cellulose and the quantity of volatile fatty acids produced (Adriani and Mushawwir, 2009). The NH₃ concentration was 4.136-4.895 mM, which is within the normal range for supporting the growth of microbial protein synthesis. Paengkoum et al. (2006) found that a 5-20 mg/dL concentration of N-NH₃ was optimal for digestion, which is equivalent to 3.57-14.28 mM. Microbial growth is not optimal if NH₃ levels are less than 3.6 mM (McDonald et al., 2010). In our study, the lowest NH₃ concentration was 4.136 mM (treatment A); a decrease in the concentration of NH₃ with increasing concentrations of VFA was reflected in a higher content of total microbes, which increased microbial protein synthesis after treatment D due to the increased use of nitrogen by the rumen bacteria. Similar results were reported by Zain et al. (2010 a,b,c). Rumetor (2009) argued that a decrease in N-NH₃ resulted in a reduction in protein digestibility, whereas increased levels of VFA are due to an increase in the digestibility of carbohydrates. This mechanism suggests that the types of microbes change with decreasing NH₃-N; the population of proteolytic microbes is reduced while the population of cellulolytic microbes increases. Microbial protein synthesis is affected by NH₃ availability in the rumen because it is a major source of nitrogen and essential for microbial protein synthesis. An environment containing approximately 80% ammonia is required for the growth of rumen bacteria (Baldwin and Allison, 1983). The effects of the addition of the minerals P, S and Mg on palm frond fermentation by the fungus *Phanerochaete chrysosporium* and their effects on microbial protein synthesis are shown in Table 2. Supplementation with P, S and Mg had no effect ($p>0.05$) on the rumen bacterial and protozoan populations. This is because the addition of P, S and Mg did not affect the pH of the rumen fluid. The protozoan population in this study ranged from 5. 851-5.864 cells/mL (log 10). The same results were reported by Badarina et al. (2013); in their study, coffee husks fermented by *Pleurotus ostreatus* had a protozoa count of 5.69-5.71 cells/mL (log 10) following supplementation. In this study, the population of bacteria was 6.843-6.987 cells/mL (log 10), which was in agreement with Stewart (1991), who found that bacterial counts were 109-1012 cells/mL in rumen contents. Supplementation with P, S and Mg had no effect on the total bacteria because supplementation with P, S and Mg did not affect the number of protozoa. This result shows that the numbers of protozoa and bacteria are in equilibrium due to a decreased number of protozoa, which increased the number of total bacteria. Supplementation with P, S and Mg during the fermentation of OPFs did not affect microbial protein synthesis. This can be concluded because all treatments (with the same nutrient content) allowed rumen microbial degradation to occur. Microbial protein synthesis (MPS) rates ranged from 36.878-46.816 mg/ml. The lowest MPS, 36.878 mg/ml, was found for treatment A (without the addition of minerals). The highest MPS was obtained for treatment D (with supplementation of fermented OPFs with P, S and Mg). This indicates that the minerals P, S and Mg are required for rumen microbial growth and is in agreement Table 1: Supplementation of P, S and Mg in the oil palm fronds fermented and their effects on rumen fluid characteristic -----

Treatments	A	B	C	D	Measured parameters
					VFA (mM)
	6.875±0.05	6.875±0.05	6.800±0.08	6.875±0.05	95.791±11.47b
	104.823±2.41ab	102.050±6.52ab	111.007±8.98a	4.895±0.28ab	4.817±0.47a
	4.773±0.48ab	4.136±0.17b	<u>Means in the same row with different letters (a and b) are significantly (p<0.05)</u>		

A: Oil palm frond fermented without the addition of mineral (control) B: Oil palm frond fermented+mineral P (0.4% DM) C: Oil palm frond fermented+mineral P (0.4% DM)+S (0.3% DM) D: Oil palm frond fermented+mineral P (0.4% DM)+S (0.3% DM)+Mg (0.1% DM) Table 2: Supplementation of P, S and Mg in the oil palm fronds fermented and their effects on Microbial Protein Synthesis (MPS) -----

Treatments	A	B	C	D	Measured parameters
					Protozoa (cell/mL) (log 10)
	5.851±0.05	5.855±0.02	5.854±0.07	5.864±0.01	Bakteri (cell/mL) (log 10)
	6.987±0.04	6.866±0.12	6.852±0.12	6.843±0.12	MPS (mg/ml)
	43.938±7.63	44.067±9.64	46.816±10.85	<u>Means in the same row with different letters (a and b) are significantly (p<0.05)</u>	

A: Oil palm frond fermented without the addition of mineral (control) B: Oil palm frond fermented+mineral P (0.4% DM) C: Oil palm frond fermented+mineral P (0.4% DM)+S (0.3% DM) D: Oil palm frond fermented+mineral P (0.4% DM)+S (0.3% DM)+Mg (0.1% DM) with Slyter et al. (1986), who showed that sulfur stimulates rumen microbial growth. Mg and Co also play roles in the growth of microbes and in the functioning of enzyme systems (Little, 1986); these minerals are often deficient in low-quality forage (Leng, 1990). The optimal rumen microbial growth leading to high microbial protein synthesis in treatment D increased the fermentability of fermented OPFs in the rumen. This was accompanied by the highest total VFA of 111.007 mM. This result agrees with those of Zain et al. (2010 a,b,c). Sanchez (2009) argued that it is difficult to determine the proportion of substrate degradation that contributes to energy production and the proportion that contributes to microbial protein synthesis because microbial growth is supported by the fermentation of substrate and the fermentation of the substrate

occurs via microbial growth. Microbial protein synthesis depends on determining the correct amount of nitrogen in the feed, the absorption of ammonia and amino acids, the flow of material out of the rumen, the amino acids required by the microbes and rumen fermentation according to the type of feed. Conclusion: Supplementation of fermented OPFs [with P, S and Mg](#) to optimize [the rumen](#) bioprocess results in the highest concentration of VFA and the greatest amount of microbial protein synthesis. ACKNOWLEDGEMENTS This research is [part of the Ph.D. dissertation of Dewi Febrina in Animal Nutrition, Faculty of Animal Science Andalas University](#). We thank the undergraduate students who collected the data and analyzed samples at the Laboratory of Ruminant Nutrition in the Faculty of Animal Science at Andalas University. REFERENCES Adriani, L. and A. Mushawwir, 2009. Level of Blood Glucose, Lactose and Dairy Cattle Milk Yield at Different Level of Macro Mineral Supplementation. *J. Indon. Trop. Anim. Agric.*, 34: 88-95. Badarina, I., D. Evvyernie, T. Toharmat, E.N. Herliyana and L.K. Darusman, 2013. Nutritive value of coffee husk fermented with *Pleurotus ostreatus* as ruminant feed. *Media Peternakan*, 36: 58-63. Bal, M.A. and D. Ozturk, 2006. Effect of sulfur containing supplements on ruminal fermentation and microbial protein synthesis. *Res. J. Anim. and Vet. Sci.*, 1: 33-36. Baldwin, R.L. and M.J. Allison, 1983. Rumen metabolism. *J. Anim. Sci.*, 57: 2209-2215. Bannink, A., J. France, S. Lopez, W.J.J. Gerrits, E. Kebreab, S. Tamminga and J. Dijkstra, 2008. Modelling the implications of feeding strategy on rumen fermentation and functioning of the rumen wall. *Anim. Feed Sci. Technol.* 143: 3-26. Bravo, D., D. Sanvant, C. Bogaert and F. Meschy, 2003. Quantitative aspects of phosphorus absorption in ruminant. *Reprods. Nutr. Dev.*, 43: 271-284. INRA. EDP, Sciences. Chiba, L.I., 2009. *Animal Nutrition Handbook*. Second Revision. Church, D.C., 1988. *The Ruminant Animal Digestive Physiology and Nutrition*. Prentice Hall, Englewood Cliff, New York. Dorado, J., G. Almendros, S. Camarero, A.T. Martiinez, T. Vares and Hatakka, 1999. Transformation of wheat straw in the course of solid-state fermentation by four ligninolytic basidiomycetes. *Enzymes Microb. Technol.*, 25: 605-612. Febrina, D., 2014. Biological delignification by *Phanerochaete chrysosporium* with addition of mineral Ca and its effect on nutrient content of oil palm frond. Research Report (unpublished). Febrina, D., J. Novirman, Z. Mardiati, Khasrad and M. Rini, 2014. Biological delignification by *Phanerochaete chrysosporium* with addition of mineral Mn and its effect on nutrient content of oil palm frond. The 16th AAAP Animal Science Congress November 10-14, 2014. Yogyakarta, Indonesia, pp: 1723-1726. Fragoeiro, S. and N. Magan, 2005. Enzymatic activity, osmotic stress and degradation of pesticide mixtures in soil extract liquid broth inoculated with *Phanerochaete chrysosporium* and *Trametes versicolor*. *Environ. Microbiol.*, 7: 348-355. France, J. and J. Dijkstra, 2005. Volatile Fatty Acid Productions. In: *Quantitative Aspect of Ruminant Digestion and Metabolism*. 2nd Ed. CAB. International, Cambridge, USA, ISBN. Imsya, E.B. Laconi, K.G. Wiryawan and Y. Widyastuty, 2013. Identification of phenolic compounds and its antioxidant activity from lignin and palm oil frond fermented with *Phanerochaete chrysosporium*. Proceedings of the 4th International Conference on Sustainable Animal Agriculture for Developing Countries (SAADC 2013) 27-31 July 2013. Lanzhou University Lanzhou, China, pp: 310-312. Kamra, D.N., 2005. Rumen microbial ecosystem. Special Section: Microbial Diversity. *Curr. Sci.*, 1: 124-135. Keidane, D. and E. Birgele, 2003. The efficacy of feed on the intra ruminal and intra abomasal pH dynamics in goats. *Vet. IR Zootechnica*, 22: 55-61. Komisarczuk, S. and M. Durand, 1991. Effect of Mineral on Microbial Metabolism and Ruminant Digestion. INRA. Publ. Versailles. Leng, R.A., 1990. Factor affecting the utilization of "poor quality" forage by ruminants particularly under tropical condition. In Smith RH, edition: *Nutrition Research Review*, Vol 3. Cambridge University Press, USA. Little, D.A., 1986. The mineral content of ruminant feed and the potential for the mineral supplementation in South-East Asia with particular reference to Indonesia. R.M. Dixon Ed. IDP, Camberra. Mc Donald, P., R.A. Edwards, J.F.D. Greenhalgh and C.A. Morgan, 2010. *Animal Nutrition*. 7th Edition. Longman. Scientific and Technical John Willey and Sons. Inc. New York. McDowell, L.R., J.H. Conrad, G.L. Ellis and J.K. Loosli, Sayed, A.N., N.A. Gazia, A.M.A. Ellah and S.M.A. Raheem, 1983. *Minerals for Grazing Ruminants in Tropical Regions*. Department of Animal Science, Center for Tropical Agriculture, University of Florida, Gainesville and the U.S. Agency for International Development. Ogimoto, K. and S. Imai, 1981. *Atlas of Rumen Microbiology*. Japan Scientific Societies Press. Tokyo, pp: 201-221. Paengkoum, P., J.B. Liang, Z.A. Jalan and M. Basery, 2006. Utilization of steam-treated oil palm fronds in growing saanen goats: II. Supplementation with Energy and Urea. *Asian-Aust. J. Anim. Sci.*, 19: 1623-1631. Peters, J.P., J.A.Z. Leedle and J.B. Paulissen, 1989. Factor affecting the in vitro production of volatile fatty acids by mixed bacterial populations from the bovine rumen. *J. Anim. Sci.*, 67: 1593-1602. Preston, T.R. and R.A. Leng, 1987. *Matching Ruminant Production System with Available Resources in The Tropics and Sub-tropics*. Penambol Books. Armidale. Rabinovich, M.L., A.V. Bolobova and Vasilchenko, 2004. Fungal decomposition of natural aromatic effect of organic matter digestibility on obligatory faecal phosphour loss in lactating goats, determined from balance data structures and xenobiotics. A Review: *Appl. Biochem. Microbiol.*, 40: 1-17. Rodehutsord, M. Heuvers and H. Pfeffer, 2000. Effect of organic matter digestibility on obligatory faecal phosphour loss in lactating goats, determined from balance data. *Anim. Sci.*, 70: 561-568. Rumetor, S.D., 2009. Supplementation of Bangun- bangun leaf (*Coleus amboinicus* Lour) and Zinc- Vitamin E In Ration on in vitro rumen metabolism of Etawah Grade Goats. *J. Ilmu Peternakan*, 4: 74-82. Sanchez, C., 2009. Lignocellulosic residues: biodegradation and bioconversion by fungi. *Biotechnol. Advan*, 27: 185-194. Slyter, L, L.W. Chaluppa, R.R. Oltjen and J.M. Weafer, 1986. Sulfur influences on rumen microorganism in vitro and in sheep and calves. *J. Anim. Sci.*, 63: 1949. Stewart, C.S., 1991. *The Rumen Bacteria*. In: *Rumen Microbial Metabolism and Rumen Digestion*. J.P. Jouany (Ed.). Institut National De La Recherche Agronomique, Paris. Suyitman, Lili and W. Evitayani, 2013. S and P mineral supplementation of ammoniated palm leaves as ruminant feed. *Pak. J. Nutr.*, 12: 903-906. Van Soest, P.J., 1994. *Nutritional Ecology of The Ruminant*. 2nd ed. Comstock Publishing Associates. A. Division of Cornell University Press. Ithaca and London. Wen, X.H., Y.N. Jia and J.X. Li, 2009. Degradation of tetracycline and

oxytetracycline by crude lignin peroxidase prepared from *Phanerochaete chrysosporium*-A white rot fungus. *Chemosphere*, 75: 1003-1007. Yu, M., G. Zeng, Y.N. Chen, H.Y. Yu, D. Huang and L. Tang, 2009. Influence of *Phanerochaete chrysosporium* on microbial communities and lignocellulose degradation during solid-state fermentation of rice straw. *Process Biochem.*, 44: 17-22. Zahari, M.W., O. Abu Hassan, H.K. Wong and J.B. Liang, 2003. Utilization of oil palm frond: base diets for beef and dairy production in Malaysia. *Asian-Aust. J. Anim. Sci.*, 16: 625-636. Zain, M., J. Rahman and Khasrad, 2014. Effect of palm oil by products on in vitro fermentation and nutrient digestibility. *Anim. Nutr. and Feed Technol.*, 14: 175- 181. Zain, M.N., Jamarun and Zulkarnaini, 2010a. Effect of phosphorus and sulfur supplementation in growing beef cattle diet based on rice straw ammoniated. *Asian J. Sci. Res.*, 3: 184-188. Zain, M., N. Jamarun and A.S. Tjakradijaya, 2010b. Tjakradidjaja. Phosphorus supplementation of ammoniated rice straw on rumen fermentability, synthesised microbial protein and degradability in vitro. *World Academy of Sci. Eng. and Technol.*, Vol. 4. Zain, M., Nurhaita and N. Jamarun, 2010c. Effect of sulfur supplementation on in vitro fermentability and degradability of ammoniated rice straw. *Pak. J. Nutr.*, 9: 413-415. *Pak. J. Nutr.*, 15 (3): 299-304, 2016 *Pak. J. Nutr.*, 15 (3): 299-304, 2016 *Pak. J. Nutr.*, 15 (3): 299-304, 2016 *Pak. J. Nutr.*, 15 (3): 299-304, 2016 300 301 302 303 304