

A.I.C.1(g)

Turnitin Originality Report

Processed on: 01-Mar-2020 10:00 PM +08

ID: 1266840618

Word Count: 6231

Submitted: 1

11 By Sri Melia

2% match
(publications)

[Analianasari, M Apriyani. ""](#),
[Journal of Physics: Conference Series, 2018](#)

Similarity Index

12%

Similarity by Source

Internet Sources:	N/A
Publications:	12%
Student Papers:	N/A

1% match
(publications)

[E Mirwandono, M Sitepu, T H Wahyuni, Hasnudi, N Ginting, G AW Siregar, I Sembiring. "Nutrition quality test of fermented waste vegetables by bioactivator local microorganisms \(MOL\) and effective microorganism \(EM4\)", IOP Conference Series: Earth and Environmental Science, 2018](#)

1% match (publications)

[Irmayanti, Liya Fitriyana. "Fortification of Yellow Potatoes Sweet Flour and Jamblang Fruit Paste \(Syzygiumcumini\) in the Making of Biscuits on it's Sensoric Characteristics and Mineral Content", Journal of Physics: Conference Series, 2019](#)

1% match (publications)

[F. Karadas, V. Pirgozliev *, T. Acamovic, M.R. Bedford. " The effects of dietary phytase activity on the concentration of Coenzyme Q in the liver of young turkeys and broilers ", British Poultry Abstracts, 2005](#)

1% match (publications)

[Marco A. de Paula Brotto, Thomas M. Nosek, Ralph C. Kolbeck. "Influence of Ageing on the Fatigability of Isolated Mouse Skeletal Muscles from Mature and Aged Mice", Experimental Physiology, 2002](#)

1% match (publications)

["Advertisements", World's Poultry Science Journal, 2019](#)

1% match (publications)

[A M N Fitratullah, F Maruddin, F N Yuliati, K I Prahesti, M Taufik. " Addition of red dragon fruit \(\) on yogurt: Effect on lactic acid content, pH, and the inhibition of growth ", IOP Conference Series: Earth and Environmental Science, 2019](#)

1% match (publications)

["IoT based Smart Farming using Cloud Computing and Machine Learning",](#)

International Journal of Innovative Technology and Exploring Engineering, 2019
<p>1% match (publications)</p> <p>Indri Juliyarsi, Arief, Akmal Djamaan, Endang Purwati. "Characteristics based of edible film made from whey with isolated lactic acid bacteria from tempoyak as probiotics packaging", IOP Conference Series: Earth and Environmental Science, 2019</p>
<p>1% match (publications)</p> <p>"Proceeding of the 2nd International Conference on Tropical Agriculture", Springer Science and Business Media LLC, 2018</p>
<p>< 1% match (publications)</p> <p>M S H Aziz, G J Manuhara, R Utami, L U Khasanah. " The Application of Active Paper Incorporated with Oleoresin of Cinnamon Leaf () Distillation Residues on Maintaining Dragon Fruits () Quality during Storage ", IOP Conference Series: Materials Science and Engineering, 2018</p>
<p>< 1% match (publications)</p> <p>Sri Melia, Endang Purwati, Yuherman ., Jaswandi ., Salam N. Aritonang, Mangatas Silaen. "Characterization of the Antimicrobial Activity of Lactic Acid Bacteria Isolated from Buffalo Milk in West Sumatera (Indonesia) Against Listeria monocytogenes", Pakistan Journal of Nutrition, 2017</p>
<p>< 1% match (publications)</p> <p>A W Pangistika, Z Bachruddin, A Kurniawati, R Utomo, C Hanim. " The effect of addition of turmeric (L.) on the rancidity process of concentrate feed based on lactic acid bacteria fermentation during aerobic storage ", IOP Conference Series: Earth and Environmental Science, 2019</p>
<p>< 1% match (publications)</p> <p>Salam N Aritonang, Elly Roza, Evy Rossi. "The effect of whippy cream adding on the quality of frozen soyghurt as symbiotic ice cream", IOP Conference Series: Earth and Environmental Science, 2019</p>
<p>< 1% match (publications)</p> <p>Olajire Fagbola, Olusola J Oyetunji, Oluwale Osonubi, Kalemani Mulongoy. "Greenhouse evaluation of two woody hedgerows as affected by arbuscular mycorrhizal fungus (AMF), soil quality and moisture", Archives of Agronomy and Soil Science, 2005</p>
<p>< 1% match (publications)</p> <p>Eveline, M Audina. " Utilization of Super Red Dragon Fruit Peel ((F.A.C. Weber) Britton & Rose) in the Making of Fermented Beverage ", IOP Conference Series: Earth and Environmental Science, 2019</p>
<p>< 1% match (publications)</p> <p>R Talon, D Walter, M.C Montel. "Growth and effect of staphylococci and lactic acid bacteria on unsaturated free fatty acids", Meat Science, 2000</p>
<p>< 1% match (publications)</p> <p>Liss Dyah Dewi Arini. "Pemanfaatan Bakteri Baik dalam Pembuatan Makanan Fermentasi yang Bermanfaat untuk Kesehatan", Biomedika, 2017</p>
<p>< 1% match (publications)</p> <p>Desmelati, Sumarto, Dewita, Dahlia, Syafrijal, P A Sari. " Determination of</p>

Nano-Collagen Quality from Sea Cucumber ", IOP Conference Series: Earth and Environmental Science, 2020

< 1% match (publications)

Afriani Sandra, Yulianti Fitri Kurnia, Ade Sukma, Endang Purwati. " The chemical characteristics of yoghurt (Lactobacillus fermentum MGA40-6 and Streptococcus thermophilus) with additional puree from fruit (Melastoma malabathricum, L.) ", IOP Conference Series: Earth and Environmental Science, 2019

International Journal of Engineering & Technology, , Vol. 8, 11-Special Issue, 2018 International Journal of Engineering & Technology Website: www.sciencepubco.com/index.php/IJET doi: Research paper, Short communication, Review, Technical paper Influence of Use Lactobacillus fermentum L23 and Streptococcus thermophilus with Dragon Fruit Extract (Hylocereus Polyrrhisuz) to Quality of Microbiology, Chemistry and Organoleptic Value of Yoghurt [1]Endang Purwati*, [2]Dhiva Rezzy Pratama, [3]Sri Melia, [4]Hendri Purwanto [1]Professor of Biotechnology/ Technology of Product Husbandry Faculty of Animal Science University of Andalas, Padang West Sumatera Indonesia, [2]Bachelor Student Department of Technology of Product Husbandry Faculty of Animal Science, University of Andalas Padang, West Sumatera Indonesia, [3] Lecture Departement of Technology of Product Husbandry Faculty of Animal Science University of Andalas, Padang West Sumatera Indonesia, [4]Laboratory Biotechnology/ Technology of Product Husbandry Faculty of Animal Science, University of Andalas, Padang West Sumatera Indonesia Abstract The difference between starter concentration and dragon fruit (Hylocereus polyrrhisuz) extract concentration on the quality of microbiology, Chemistry, and organoleptic value of yoghurt. The method of research was the experimental method by using a Randomized Block Design (RBD) 3x4 factorial pattern with 2 replicates as a group. Factor A starter concentration that is A1: 4%, A2: 5%, A3: 6% and Factor is dragon fruit extract that is B1: 0%, B2: 1%, B3: 2%, B4: 3% . The parameters measured were Total Lactic Acid Bacteria colonies, Water content, protein content, fat content, antioxidant activity, pH Value and organoleptic value of yoghurt. The results show that there was a real interaction ($P < 0.05$) concentration of the starter (factor A) and concentration of the extract dragon fruit (factor B) against total lactic acid bacteria colonies. but there was no significant interaction ($P > 0.05$) on water content, protein content, fat content, antioxidant activity, pH value and organoleptic value of yoghurt. factor A can give a real effect ($P < 0.05$) to antioxidant activity and pH value of yoghurt. While factor B can give a very real effect ($P < 0.01$) on water content, protein content, fat content, antioxidant activity and pH value of yoghurt. Then for organoleptic value there is no significant effect ($P > 0.05$) from each factor to organoleptic value of yoghurt. The best percentage addition for yoghurt was 5% at starter concentration and 2% at dragon fruit (Hylocereus polyrrhisuz) extract concentration with total lactic acid bacteria colonies 170×10^8 CFU/ml, water content 81.22%, protein content 4.65%, fat content 3.58%, antioxidant activity 44.53%, pH value 4.4 and organoleptic value still in range preferred by panelists. Keywords — chemical quality, dragon fruit extract, microbiology, starter,

yoghurt. Yoghurt is a product obtained from pasteurized milk, then I. INTRODUCTION fermented with certain bacteria until acidity, odor and taste are One food product that comes from livestock and is favored by typical, with or without the addition of other ingredients [3]. some people, namely milk. Milk is a food ingredient that has high Yoghurt is milk that is fermented by using microbial mixed cultures, nutritional value, so that it becomes a medium that is highly favored generally Lactobacillus bulgaricus and Streptococcus thermophilus, by microorganisms for its growth and development [1]. With the so as to produce consistency resembling pudding. In addition to the growth of various microbes can change the quality of milk which is use of Lactobacillus bulgaricus bacteria can also be used other characterized by changes in taste, aroma, color and appearance that bacteria such as Lactobacillus bacteria isolated from buffalo causes milk to be damaged. Therefore, to improve the quality of milk[4]. milk so as not to become damaged quickly, various techniques can Through isolation and identification of buffalo milk in Agam be carried out both processing and preserving, one type of Regency, West Sumatra, 88 strains of LAB were produced, most of which were characterized as rod, gram positive, catalase-negative, preservation in milk is generally known as fermentation. Products homofermentative and heterofermentative. One of the isolates from fermented milk which are commonly known include yoghurt included Lactobacillus fermentum L23 (A 3.3) type of LAB which [2]. showed high inhibition against Listeria monocytogenes [5]. Then according to [6] which states that Lactobacillus fermentum strain L23 can produce bacteriocin and is stable at hot temperatures with molecular mass. Lactobacillus bulgaricus substitution in yoghurt with Lactobacillus fermentum L23 bacteria is expected to change the texture of yoghurt to be more dense and can produce bacteriocin so that it can inhibit the growth of pathogenic bacteria in food. While the addition of Streptococcus thermophilus to milk can produce lactic acid which can also preserve milk. During fermentation, this bacterium produces acetaldehyde which forms the aroma in yoghurt [7]. Yoghurt is very beneficial for people who are not resistant to milk sugar (lactose), which is known as a sufferer of "lactose intolerance". Therefore yoghurt can be linked to the need for functional food for the general public. According to the Regulation of the Head of the Food and Drug Supervisory Agency [8], it is stated that functional food is processed food containing one or more functional components based on scientific studies having certain physiological functions, proven to be harmless and beneficial to health. Thus to improve the quality of yoghurt as a functional food, yoghurt can be combined with other components of fruits, such as by adding dragon fruit extract. This is because dragon fruit contains important components that can increase the antioxidant content of yoghurt. Dragon fruit (Hylocereuspolyrhizus) is a horticultural plant that was developed in Indonesia. Dragon fruit contains bioactive substances which are beneficial for the body including antioxidants (ascorbic acid, beta carotene and anthocyanin) and contain dietary fiber in the form of pectin. With its sweet and refreshing taste, dragon fruit is also rich in benefits such as lowering cholesterol and balancing blood sugar, binding cancer-causing carcinogens and improving digestion. Then than that dragon fruit also contains oligosaccharides, including raffinose, stakiose and fructo-oligosaccharides [9]. According to previous research [10], the use of dragon fruit extracts on yoghurt ranged from 7% could reduce pH

and increase the acidity of yoghurt and increase the inhibitory power of *E. coli* along with the addition of dragon fruit concentration. II. METHOD OF RESEARCH A. Preparation of Extract Dragon Fruit Extract dragon fruit were made according to [11] dragon fruit peel was peeled, the flesh was chopped and then oven using 60°C until it dries. After drying the blender until it's shaped like flour. Weigh 50 grams of dragon fruit flour, put it into erlemenyer and add 70% polar ethanol solvent as much as 500 ml. The maceration results are filtered using filter paper, separated between powder and solution. The filtrate obtained was evaporated vacuum using a rotary evaporator to remove the solvent. B. Preparation Yoghurt Starter In this study preparation yoghurt starter based on method of [12]. C. Yoghurt Making In this study, yoghurt making based on method of [12]. D. Data Analysis This experiment used experimental method designed with Randomized Block Design (RBD) factorial pattern 3 x 4 and 2repetition. The treatment is a starter addition factor, consisting of (A1) addition of 4% stater. (A2) 5% starter addition.(A3) addition of 6% starter.Factor B is the addition of extract red dragon fruit, consisting of (B1) addition of 0%, (B2) addition of 1%, (B3) addition of 2%, (B4) addition of 3%. The treatments that showed the results were significantly different (F count> F table 0.05). Then further tests were performed using Duncan's Multiple Range Test (DMRT). In this study, the measured variables are the Total Lactic Acid Bacteria colonies, Water content, protein content, fat content, antioxidant activity, pH Value and organoleptic value of yoghurt. III. RESULT OF RESEARCH A. Total Lactic Acid Bacteria Colonies The results of statistical analysis showed that there were interactions (P <0.05) between difference in the addition of starter concentration (factor A) and fruit extract concentration dragon (factor B) to total yoghurt lactic acid bacteria. This shows that an increase in the amount of starter and extract of dragon fruit affects the total BAL. Average Total Lactic Acid Bacteria in each treatment presented in Table 1. Table 1. Average total lactic acid bacteria ($\times 10^8$ CFU/ml) difference in addition starter concentration (Factor A) and concentration of dragon fruit extract (Factor) in various treatments. Factor A B1 Factor B B2 B3 B4 Averag e A1 126cd 154bc 179ab 112de 142 A2 101def 203a 170ab 83ef 139 A3 73f 100def 147bc 91ef 91 Average 100 152 165 95 In Table 1, show the interaction is significantly different (P <0.05) Starter (Factor A) with dragon fruit extract (factor B) on total lactic acid bacteria. The results of the variance analysis show the interaction between factor A and factor B on the total effect of lactic acid bacteria was significantly different (P <0.05). Duncan's multiple range test results showed that the interaction of the treatment of total lactic acid bacteria on A2B2 203×10^8 CFU / ml treatment was significantly different (P <0.05) against treatment A1B3 179×10^8 CFU / ml, A2B3 170×10^8 CFU / ml, A1B2 154×10^8 CFU / ml, A3B3 147×10^8 CFU / ml, A1B1 126×10^8 CFU / ml, A1B4 112×10^8 CFU / ml, A3B4 91×10^8 CFU / ml, A2B4 $\times 10^8$ CFU / ml, A3B1 73×10^8 CFU / ml. Different real (P <0.05) total lactic acid bacteria caused by starter concentration and addition of dragon fruit extract. Starter interaction (Factor A) and fruit extract naga (factor B) shows the total increase in lactic acid bacteria substitution of starter (factor A) and dragon fruit extract (Factor B) on yoghurt. The table above shows the total average of lactic acid bacteria by giving 4%, 5% and 6% starter and giving 0%, 1%, 2% 3% extract Dragon fruit ranges from 73×10^8 CFU / gr - 203×10^8 CFU / ml. Total low BAL in A3B1 treatment, namely yoghurt with 3% starter

addition (A3) and without the [addition of dragon fruit extract](#) due to the absence of extract dragon fruit that functions as a prebiotic that will spur growth probiotic bacteria *Lactobacillus fermentum* L23 and *Streptococcus thermophilus*. The higher the addition of starter causes the nutrients for BAL not sufficient for BAL growth in yoghurt [13]. Addition of dragon fruit extract red can increase the growth rate of BAL because of dragon fruit extract reds have simple sugar compounds such as oligosaccharides which are prebiotics, a group of oligosaccharides found in dragon fruit extract, namely raffinosa. This is in accordance with the opinion of [14] stating that BAL need carbohydrates to be fermented, these bacteria will not be able to live on a medium that contains very little carbohydrate. [15] added that [red dragon fruit](#) (*Hylocereus polyrhizus*) [contains carbohydrates and other](#) compounds, and is rich in antioxidants. Carbohydrate on [red dragon fruit can be used](#) by BAL as a [source](#) of energy during fermentation. The total height of BAL in A2B2 treatment was 203×10^8 CFU / ml due to the addition of 4% starter has been maximally utilized dragon fruit extract as a prebiotic to increase the total BAL. Treatment A2B2 differs not significantly ($P > 0.05$) with A1B3 so it can be known [addition of 1% dragon fruit extract](#) has increased [the total](#) LAB. Yoghurt with the addition of 4% starter concentration (A2) and [red dragon fruit](#) extract at [a concentration of 1%](#) (B2) [is the best medium for growth the microorganisms](#) used, the condition is seen increasingly the total number of lactic acid bacteria. In treatment A2B4 83×10^8 CFU / ml and A3B4 91×10^8 CFU / ml decreased due to poor environmental conditions acid. The pH of yoghurt from the results of the study decreased 4.3 - 4.2, so the buildup H^+ ions in cells will cause electrolyte imbalances in cells bacteria so that bacteria will try to remove H^+ which results bacteria release large amounts of ATP and cause cell death [16]. [17] stated that besides produce [lactic acid, lactic acid](#) bacteria [also produce other](#) metabolites [such as](#) hydrogen peroxide [and](#) other antibiotic compounds which, besides being [able to](#) inhibits [the growth of pathogenic bacteria](#), in large accumulations it is possible to inhibit lactic acid bacteria too. Antibacterial activity from dragon fruit extract causes the work [of lactic acid bacteria in the](#) description Lactose [is](#) inhibited, so [the growth of](#) BAL is less stable. But this is the case not too influential, the amount of lactic acid bacteria found in yoghurt it is in accordance with the requirements of probiotic drinks. Requirements for probiotic drinks contain more than 10^8 cells of probiotic bacteria per milliliter of life [18] and according to World Health Organization [19] is 10^9 CFU / ml. Every treatment produces the total BAL that meets the minimum standard is 10^7 CFU / ml [20]. The average amount of lactic acid bacteria found in yoghurt is 73×10^8 CFU / ml - 203×10^8 CFU / ml. This shows that raw materials and the substrate used can produce yoghurt with good quality [21].

B. Water content The results of statistical analysis show that there is no interaction ($P > 0.05$) between Factor A (Starter) and Factor B (Dragon fruit extract) on water content. The average value of water content in each treatment is presented in Table 2. Table 2. Average yoghurt water content (%) difference in the addition of starter concentration (factor A) and concentration of dragon fruit extract (factor B) on various treatment. Factor A B1 Factor B B2 B3 B4 Average A1 83.92 82.65 81.59 80.85 82.25 A2 83.83 82.03 81.22 81.80 82.04 A3 83.98 82.60 81.62 81.71 82.48 Average 83.91a 82.43b 81.48c 81.22d

In Table 2, shows that yoghurt with differences the addition of starter

concentration (Factor A) has an unreal effect ($P > 0.05$) and have a significant effect ($P < 0.05$) on addition concentration of dragon fruit extract on yoghurt water content. Addition starter concentration (Factor A) has no significant effect ($P > 0.05$) on Yoghurt water content, this is due to Lactobacillus fermentum L23 and Streptococcus thermophilus is a fermented type of lactic acid bacteria homofermentative wherein the results of metabolism do not produce water. So that the activity of lactic acid bacteria has no real effect on water content. As seen in the results of the addition of concentration research starter 4% (A1) with an average of 82.25%, 5% (A2) with an average of 82.04% and starter 6% (A3) with an average of 82.48%. In yoghurt with the addition of dragon fruit extract (Factor B) in the Table 4, shows that there is an influence on water content ranging from 81.22% - 83.91%. The results of variance analysis showed that yoghurt was added Dragon fruit extract (Factor B) had a very significant effect ($P < 0.05$) on yoghurt water content, which is the highest in treatment without addition Dragon fruit extract 0% (B1) with an average moisture content of 83.91% and the lowest on addition treatment of dragon fruit extract 3% (B4) with an average water content 81.22%. According to the opinion of [22] who states that pectin is an anion polysaccharide complex that is present on the primary and intercellular cell walls in high-level plants. Pectin widely used as a functional component in the food industry because of its ability to form a gel. So that it increases the addition of dragon fruit decreases in water content. The results in this study the yoghurt water content obtained ranged 81.10 - 83.61%, not much different from the results of research [23], namely ranges from 81.26 - 85.56% with the addition of cinnamon bark extract. [24] states that commercial yoghurt uses addition of emulsifiers, flavors, and preservatives, so that the water content lower commercial yoghurt is produced. According to [25] the composition of commercial yoghurt has a moisture content of 75-80%. This is because commercial yoghurt uses a lot of additional ingredients make commercial yoghurt water content lower.

C. Protein Content The results of statistical analysis showed that there was no interaction ($P > 0.05$) between the difference in the addition of starter concentration (factor A) and the concentration of dragon fruit extract (factor B) on the protein content of yoghurt. The absence of interaction between the addition of starter concentration (factor A) and the concentration of dragon fruit extract (factor B) was due to the pH produced from the starter and dragon fruit extract still at the isolate point which ranged from 4.6. So that shows no interaction on the protein content of yoghurt. The average level of yoghurt protein in each treatment is presented in Table 3. Table 3. Average yoghurt protein content (%) difference in addition of starter concentration (factor A) and concentration of dragon fruit extract (factor B) in various treatments.

Factor A	B1	Factor B	B2	B3	B4	Average
A1	2.75	3.74	3.96	3.99	3.61	A2
2.75	3.80	4.65	4.37	3.89	A3	2.47
3.71b	4.33a	4.25a				3.58
						4.38
						4.41
						3.71
						Average
						2.66c

In Table 3, shows that the average protein content in the treatment of starter concentration (factor A) ranged between 3.61% - 3.89%. The treatment of the addition of starter concentration (factor A) did not give a significant effect ($P > 0.05$) on the protein content of yoghurt. In contrast, it was not significant that yoghurt protein content due to the addition of starter concentration was caused by the pH value of yoghurt with the addition of a starter concentration (Factor A) of 4-6% resulting in a pH value of 4.4-4.7 yoghurt. Regarding the pH of

yoghurt, it is still in the range of milk protein isolation points, from 4.3 to 4.6 [26]. This condition results in no effect on the protein content of yoghurt. In accordance with the opinion the pH of the isolate of protein molecules will form the same number of positive and negative ions. So that the protein condition is relatively stable, which results in no effect on yoghurt protein.

D. Fat Content The results of statistical analysis showed that there was no interaction ($P > 0.05$) between the difference in the addition of starter concentration (factor A) and the concentration of dragon fruit extract (factor B) on the fat content of yoghurt. The absence of interaction between the addition of starter concentration (factor A) and the concentration of dragon fruit extract (factor B) was due to the absence of fat content in dragon fruit extract. So that shows no interaction on the fat content of yoghurt. The average fat content of yoghurt in each treatment is presented in Table 4. Table 4. Average yoghurt fat content (%) difference in addition of starter concentration (factor A) and concentration of dragon fruit extract (factor B) in various treatments.

Faktor A	B1	Faktor B	B2	B3	B4	Average
A1	4.08	3.73	3.53	3.42	3.69	A2
A2	4.12	3.84	3.58	3.34	3.72	A3
A3	4.18	3.69	3.29	3.34	3.63	Average
	4.13a	3.75b	3.47c	3.37c		

In Table 4, shows that yoghurt with different starter concentrations (Factor A) did not give a significant effect ($P > 0.05$) on fat content in yoghurt products. The average fat content of yoghurt with different starter concentrations (Factor A) ranged between 3.63%-3.72%. Different was not significant ($P > 0.05$) yoghurt fat content due to the addition of starter concentration was caused by an increase in starter delivery up to 6% (A3) did not cause changes to the fat content of yoghurt. So there is only a minimal starter role in breaking milk fat into free fatty acids. In accordance with the opinion of [27] which states that during the fermentation process, the bacteria will experience three main reactions by decomposing the milk component, which is decomposing lactose into lactic acid (fermentation), hydrolyzing casein into peptides and free amino acids (proteolysis) and break down milk fat into free fatty acids (lipolysis).

E. Antioxidant Activity The results of statistical analysis showed that there was no interaction ($P > 0.05$) between the difference in the addition of starter concentration (factor A) and the concentration of dragon fruit extract (factor B) on the antioxidant activity of yoghurt. The absence of interaction between the addition of starter concentration (factor A) and the concentration of dragon fruit extract (factor B) was due to both factors having their respective antioxidant activity. So that shows no interaction on the antioxidant activity of yoghurt. The average antioxidant activity of yoghurt in each treatment is presented in Table 5.

Factor	Factor B	Average
A1	19.33	30.84
A2	19.70	32.26
A3	19.92	31.82
Average	19.65a	

In Table 5, shows that yoghurt with a difference in the addition of starter concentration (Factor A) had a significant effect ($P < 0.05$) on antioxidant activity in yoghurt. The average antioxidant activity of yoghurt with the difference in the addition of starter concentration (Factor A) ranged between 37.48% -38.91%. Where is the highest activity in the treatment of starter starter 6% (A3) with antioxidant activity of yoghurt averaging 38.91% and the lowest antioxidant activity in the treatment of giving starter 4% (A1) with

antioxidant activity of yoghurt average of 37.48%. This increase in antioxidant activity is also associated with a total increase in lactic acid bacteria, *Lactobacillus*. So that the higher the starter concentration, the higher the ability of LAB to break lactose into lactic acid and the more H⁺ ions donated to free radicals can increase primary antioxidant activity. As shown in Table 5, the addition of starter as much as 6% can produce antioxidant activity of 38.91%. This is in accordance with the opinion of [28] which states that the ability to change sugar to lactic acid by lactic acid bacteria is synergistic by giving H⁺ ions to free radicals thus increasing the primary antioxidant activity. In addition, the ability of lactic acid bacteria to break down proteins (proteolytics) into small peptides (bioactive peptides) which have antioxidant activity is also associated with increased antioxidant activity in yoghurt [29].

F. pH value The results of statistical analysis show that there is no interaction ($P > 0.05$) between starter (Factor A) and dragon fruit extract (Factor B) on pH value. However, each factor showed a significant effect ($P < 0.05$). The average pH value for each treatment is presented in Table 6.

Table 6. Level of pH Value difference in addition of starter concentration (Factor A) and concentration of dragon fruit extract (Factor B) on various treatments.

	Factor A B1	Factor B B2	B3	B4	Average
A1	4.8	4.6	4.7	4.6	4.7a
A2	4.7	4.5	4.4	4.3	4.5b
A3	4.6	4.4	4.3	4.2	4.4b
Average	4.7a	4.5b	4.5b	4.3c	

In Table 6, showed that yoghurt with a difference in the addition of starter concentration (Factor A) had a significant effect ($P < 0.05$) on the pH value of yoghurt. Average pH value of yoghurt with the difference in the addition of starter concentration (Factor A) ranges from 4.4 - 4.7. Where is the highest activity in the treatment of giving starter 4% (A1) with a yoghurt pH value averaging 4.7 and the lowest pH value on treatment of giving starter 6% (A3) with a pH value of yoghurt on average 4.4. As shown in the results of the study, on the addition of a starter highest 6% (Factor A) produces the lowest pH of 4.4. Acids the organic formed is dissociated acids in the form H⁺ ions. The more acid produced, the more too H⁺ ions are formed so that the pH measurement by the pH meter electrode shows a decreasing value. The more number of BAL more and more metabolites in the form of lactic acid can dissociate in H⁺ ions so that the pH becomes lower, as evidenced by the greater the proportion of red dragon fruit extract increases the total acid of yoghurt.

G. Organoleptic Value 1. Taste The results of the statistical analysis showed that there was no significant interaction ($P > 0.05$) between the differences in the introduction of starter concentration (Factor A) and the addition of dragon fruit extract (factor B) to the assessment of the taste of yoghurt flavor. The average score of organoleptic taste in each treatment is presented in Table 7.

Table 7. Average taste of organoleptic test on yoghurt (%) difference in giving starter concentration (factor A) and concentration of dragon fruit extract (factor B) on various treatments.

	Factor A B1	Factor B B2	B3	B4	Average
A1	1.84	2.04	1.64	1.76	1.82
A2	1.56	1.88	1.88	1.84	1.79
A3	1.96	1.96	1.92	1.92	1.94
Average	1.95	2.01	1.81	1.84	

Based on the results of Friedman diversity analysis showed that the treatment of adding starter concentration differences (Factor A) and addition of dragon fruit extract (Factor B) did not give a significant effect ($P > 0.05$) on the flavor of organoleptic yoghurt with an average organoleptic value ranging between 1.64 - 2.08. The absence of a significant effect ($P > 0.05$) on the treatment of adding starter

concentration differences (Factor A) to the organoleptic value of yoghurt flavor, due to the addition of starter concentrations of 4% to 6% did not give a different effect on the taste of yoghurt produced on this research. As seen in the results of the study, the addition of starters 4% to 6% shows a score of 1.82 - 2.04 which is still within the panelists' preferred range. 2. Aroma The results of statistical analysis showed that there was no significant interaction ($P > 0.05$) between the differences in the introduction of starter concentration (Factor A) and the addition of dragon fruit extract (factor B) to the assessment of organoleptic aroma of nyoghurt. The average score for organoleptic aroma in each treatment is presented in Table 8. Table 8. Average of organoleptic aroma test on yoghurt (%) giving additional starter concentration (factor A) and concentration of dragon fruit extract (factor B) on various treatments.

Factor A	B1	Factor B	B2	B3	B4	Average
A1	1.80	2.04	1.88	1.96	1.92	A2
1.92	1.96	2.00	2.20	2.02	A3	1.76
2.04	2.04	1.88	1.93	Average	1.83	
2.01	1.97	2.01	Ased	on the results of	Friedman diversity	analysis

showed that the treatment of adding starter concentration differences (Factor A) and addition of dragon fruit extract (Factor B) did not give a significant effect ($P > 0.05$) on organoleptic aroma of yoghurt with an average organoleptic value ranging from 1.76 - 2,20. The absence of a significant effect ($P > 0.05$) on the treatment of adding starter concentration differences (Factor A) to the organoleptic value of the aroma of yoghurt, due to the addition of starter concentrations of 4% to 6% did not give a different effect on the aroma of yoghurt produced on this research. As seen in the results of the study, the addition of a starter of 4% to 6% shows a score of 1.92 - 2.02 which is still in the range of panelists' preference. 3. Textsture The results of statistical analysis showed that there was no significant interaction ($P > 0.05$) between the difference in the introduction of starter concentration (Factor A) and the addition of dragon fruit extract (factor B) to the organoleptic texture of yoghurt. The average organoleptic texture for each treatment is presented in Table 9. Table 9. Average of organoleptic texture test on yoghurt (%) difference in giving starter concentration (factor A) and concentration of dragon fruit extract (factor B) on various treatments.

Factor	Factor B	Average	A	B1	B2	B3	B4	Average
1.67	1.95	1.96	1.92	Based on the results of	Friedman diversity	analysis	showed	

that the treatment of adding starter concentration differences (Factor A) and addition of dragon fruit extract (Factor B) did not give a significant effect ($P > 0.05$) on organoleptic yoghurt texture with an average organoleptic value ranging from 1.56 - 2.04. The absence of a significant effect ($P > 0.05$) on the treatment of adding starter concentration differences (Factor A) to the organoleptic value of yoghurt texture, due to the addition of starter concentrations of 4% to 6% did not give a different effect on the texture of yoghurt produced on this research. As seen in the results of the study, that the addition of a starter of 4% to 6% shows a score of 1.79 - 1.92 which is still within the panelists' preferred range. The absence of the effect of starter concentration is also influenced by the type and number of microorganisms in the starter [30]. According to [31] the type and number of microorganisms in the starter that are used greatly play a role in the formation and taste and texture of yoghurt, besides that the duration of fermentation and ambient temperature also influence yoghurt making. Organoleptic value of yoghurt texture in this study shows the level of similarity of the results of [32] research on the texture of sesame-beet milk yoghurt

which showed that the most viscous yoghurt texture value of 2.06 was produced in the percentage of 14% sesame seeds with fruit extract concentration bit 15%, while the texture value of yoghurt is the lowest or dilute of 1.40 resulting in the percentage of 10% sesame seeds with a concentration of 5% fruit extract. The results showed that the organoleptic value of the yoghurt treatment was respectively 1.56-2.04 in the range of likes, meaning that the treatment of the research on the [organoleptic value of the texture was](#) still favored by [the](#) panelists. IV. CONCLUSION [The](#) results of this study concluded that there was a real interaction at concentration of the starter (factor A) and concentration of the extract dragon fruit (factor B) against total lactic acid bacteria colonies. but there was no significant interaction on water content, protein content, fat content, antioxidant activity, pH value and organoleptic value of yoghurt. factor A can give a real effect to antioxidant activity and pH value of yoghurt. While factor B can give a very real effect on water content, protein content, fat content, antioxidant activity and pH value of yoghurt. Then for organoleptic value there is no significant effect from each factor to organoleptic value of yoghurt. The best percentage addition for yoghurt was 5% at starter concentration and 2% at dragon fruit (*Hylocereus polyrhizus*) extract concentration with total lactic acid bacteria colonies 170×10^8 CFU / ml, water content 81.22%, protein content 4.65%, fat content 3.58 %, antioxidant activity 44.53%, pH value 4.4 and organoleptic value still in range preferred by panelists. ACKNOWLEDGMENT [This research was supported by](#) Laboratory of Biotechnology / [Technology of](#) Product Husbandry, Faculty of Animal Science, University of Andalas, Padang, Indonesia and Grant research Professor (RGB) University of Andalas Nomor : 41/UN.16.17/PP.RGB/LPPM/2018 chairman : Prof. Drh. Hj. Endang Purwati, MS., Ph.D. REFERENCES [1] Saleh, E. 2004. Dasar Pengolahan Susu dan Hasil Ikutan Ternak. Jurnal Program Studi Produksi Ternak Fakultas Pertanian Universitas Sumatera Utara. [2] Melia, S., Jaswandi, Yuherman, E. Purwati. 2018. Selection of buffalo milk lactic acid bacteria with probiotic potential. Asian Journal of Pharmaceutical and Clinical Research 11(6):186 · June 2018. [3] Surajudin., Kusuma., dan P. Dwi. 2006. Yoghurt, Susu Fermentasi yang Menyehatkan. Agromedia Pustaka (Revisi). Jakarta. [4] Purwati, E., James Hallyward, Indri Juliyarsi, Sri Melia, Hendri Purwanto and Puji Hartini. 2018. Effect of Addition Cinnamon Bark Extract (*Cinnamomum burmannii*) of Water Content, Total Lactic Acid Bacteria Colonies, Antioxidant Activity and Cholesterol Levels from Goat's Milk Yoghurt. Journal of Adv Research in Dynamical & Control Systems, Vol. 10, 04-Special Issue, 2018 [5] Melia, S., E. Purwati, Yuherman, Jaswandi, S. N. Aritonang and M. Silaen. 2017. Characterization of the Antimicrobial Activity of Lactic Acid Bacteria Isolated from Buffalo Milk in West Sumatera (Indonesia) Against *Listeria monocytogenes* Pakistan Journal of Nutrition ISSN 1680-5194 DOI: 10.3923/pjn.2017.645.650. [6] Pascual, L., F. Ruiz., W. Giordano and I. L. Barberis. 2010. Vaginal colonization and activity of the probiotic bacterium *Lactobacillus fermentum* L23 in a murine model of vaginal tract infection. [7] Khomsan, Ali. 2009. Rahasia Sehat Dengan Makanan Berkhasiat. Jakarta : PT Kompas Media Nusantara, hal. 10-12. [8] Badan Pengawasan Obat dan Makanan. 2005. Peraturan Kepala Badan Pengawasan Obat dan Makanan Terkait Pangan Fungsional. Badan Pengawasan Obat dan Makanan, Jakarta. [9] Khalili, R. M. A., A. B. C. Abdullah and A. A. Manaf. 2014. Isolation and characterization of oligo-

saccharides composition in organically grown red pitaya, white pitaya and papaya. *Int J Pharm Pharmaceut Sci* 6: 131-136. [10] Fitratullah, A. M. N. 2017. Pengaruh Konsentrasi Penambahan Buah Naga Merah (*Hylocereus polyrhizus*) terhadap Daya Hambat *Escherichia coli*, pH dan Keasaman Yoghurt. Naskah Skripsi-S1. Fakultas Peternakan. Universitas Hasanuddin. Makassar. [11] Gulewicz, P., D. Ciesiolka., J. Frias., C. V. Valverde., S. Frejnagel., and K. Trojanowska. 2000. Simple Method of Isolation and Purification of β -galactosides from Legumes. *J. Agric Food Chem.* 48: 3120-3123. [12] Melia, S., E. Purwati, Yuherman, Jaswandi, H. Purwanto, P. Hartini. 2018. Prosedur Pembuatan Yoghurt Probiotik dengan Ekstrak Buah Naga Merah Sebagai Sumber Probiotik dan Antoksidan. Paten No. SID21018044802. [13] Purwati, E., S. N. Aritonang, S. Melia, I. Juliyarsi and H. Purwanto. 2016. Manfaat Probiotik Bakteri Asam Laktat Dadiah Menunjang Kesehatan Masyarakat. Lembaga Literasi Dayak (LID), Tangerang. ISBN 978-602-6381-09-5 [14] Widodo. 2003. Bioteknologi Industri Susu. Lacticia Press. Yogyakarta. [15] Nurul, S. R. dan R. Asmah. 2014. Variability in Nutritional Composition and Phytochemical Properties of Red Pitaya (*Hylocereus polyrhizus*) from Malaysia dan Australia. *International Food Research Journal* 21(4):1689-1697. [16] Oeitan, A. I Nugrahani dan K Netty 2013. Pembuatan Yoghurt Murbei Hitam (*Morus Nigra L*) : Proporsi Sari Buah dan Susu Sapi Terhadap Komponen Bioaktif dan Viabilitas Bakteri Asam Laktat Selama Penyimpanan. Okawa, M. Kinjo, J. 2001. DPPH (1,1-diphenyl-2-picrylhydrazyl) Radical Scavenging Activity of Flavonoid obtained from Some Medical Plants. *Biok Pharm Bull.* Vol 24 No 10. [17] Suseno, T.I.P., Surjoseputro, S. dan Anita, K. 2000. Minuman Probiotik Nira Siwalan: Kajian Lama Penyimpanan terhadap Daya Anti Mikroba *Lactobacillus casei* pada beberapa Bakteri Patogen. *J. Teknologi Pangan dan Gizi*, 1 (1): 1-13. [18] Badan Standardisasi Nasional. 2009. SNI 01-2981-2009. Syarat Mutu Yoghurt. Standar Nasional Indonesia, Jakarta. [19] FAO/WHO. 2001. Health and Nutritional Properties of Probiotics in Food Including Power of Milk with Live Lactic Acid Bacteria. Proceedings of the FAO/WHO Expert Consultation On Evaluation of Health and Nutritional Properties of Probiotic in Food Including Power of Milk With Live Lactic Acid Bacteria. Cordoba, Argentina. [20] Badan Standardisasi Nasional. 2011. SNI 01-3141-2011. Standardisasi Nasional Indonesia SNI Susu segar. Jakarta. [21] Kumalasari, K. E. D. Nurwantoro dan S. Mulyani. 2012. Pengaruh Kombinasi Susu dengan Air Kelapa terhadap Total Bakteri Asam Laktat, Total Gula, dan Keasaman Yoghurt Drink. Universitas Diponegoro Semarang. *Jurnal Aplikasi Teknologi Pangan* 1 (2): 48-53. [22] Suwoto., A. Septiana., dan G. Puspa. 2017. Ekstraksi Pektin Pada Kulit Buah Naga Super Merah (*Hylocereus Costaricensis*) Dengan Variasi Suhu Ekstraksi dan Jenis Pelarut. *Jurnal Ilmiah Teknik Kimia UNPAM, Tangerang Selatan.* Vol. 1 No. 2. [23] Hartini, P. 2016. Pengaruh Penambahan Ekstrak Kulit Kayu Manis (*Cinnamomum burmannii*) Terhadap Kadar Air, Nilai pH, Total Koloni Bakteri Asam Laktat dan Aktivitas Antioksidan Yogurt Susu Kambing. Skripsi. Universitas Andalas. Padang. [24] Danasaputra. 2004. Direktorat Pengolahan dan Pemasaran Hasil Peternakan. Departemen Pertanian, Jakarta. [25] Sugitha, M., dan Widarta, R. W. 2012. Teknologi Susu Daging dan Telur. Buku Arti, Denpasar. [26] Mattila-Sandholm, T., and M. Saarela. 2000. Functional Dairy Product. Woodhead Publishing Limited and CRC Press LLC. Fulda, Germany. [27] Smit, G., B. A. Smit, and W. J. M. Engels. 2005. Flavour formation by lactic acid bacteria and

biochemical flavour profiling of cheese products. FEMS Microbil.Rev. 29: 591-610. [28] Primurdia, E. G. 2013. Aktivitas Antioksidan Minuman Probiotik Sari Kurma (*Phoenix dactilyfera* L.) dengan isolat *L. Plantarum* dan *L. casei*. Jurnal Pangan dan Agroindustri Vol.2 No.3 p.98-109. [29] Samichah. 2014. Aktivitas Antioksidan Dan Penerimaan Organoleptik Yoghurt Sari Wortel (*Daucus Carrota* L). Program Studi Ilmu Gizi Fakultas Kedokteran : Universitas Diponegoro. Semarang. [30] Purwati, E., S. Syukur, Husmaini, H. Purwantodan R.P. Pasaribu, 2014. Molekuler Karakteristik Bakteri Asam Laktat Isolate Dadih Air Dingin Kabupaten Solok Sumatera Barat. Jurnal Vol. 40. No.2. Hal. 134-146 [31] Prasetyo, H. 2010. Pengaruh Penggunaan Starter Yoghurt Pada Level Tertentu Terhadap Karakteristik Yoghurt Yang Dihasilkan. Skripsi. Fakultas Pertanian. Universitas Sebelas Maret. Surakarta. [32] Guruh, M. K., dan Nanik S. 2017. Karakteristik Yoghurt Susu Wijen (*Sesamunindicum*) Dengan Penambahan Ekstrak Buah Bit (*Beta vulgaris*). Universitas Slamet Riyadi. Surakarta. Artikel. [International Journal of Engineering & Technology, , Vol. 8, 11-Special Issue, 2018 1](#) [International Journal of Engineering & Technology, , Vol. 8, 11-Special Issue, 2018 2](#) [International Journal of Engineering & Technology, , Vol. 8, 11-Special Issue, 2018 3](#) [International Journal of Engineering & Technology, , Vol. 8, 11-Special Issue, 2018 4](#) [International Journal of Engineering & Technology, , Vol. 8, 11-Special Issue, 2018 5](#) [International Journal of Engineering & Technology, , Vol. 8, 11-Special Issue, 2018 6](#) 1 2 3 4 5 6