

## Effect Supplementation of Glutamate as Additive Feed in Improving Broiler Performance

Vebera Maslami<sup>1</sup>, Mirnawati<sup>2</sup>, Jamsari<sup>3</sup>, Yuliaty Shafan Nur<sup>2</sup> ~~and~~, Yetti Marlida<sup>2\*</sup>

<sup>1</sup>Department of Agricultural Science, Faculty of Agriculture, Andalas University, 25163, Indonesia

<sup>2</sup>Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, Andalas University, 25163, Indonesia

<sup>3</sup>Department of Crop Science, Faculty of Agriculture, ~~Universities~~ Andalas, 25163, Indonesia

**Commented [E1]:** Check the word

Corresponding author's E-mail: [yettimarlida@ansci.unand.ac.id](mailto:yettimarlida@ansci.unand.ac.id)

### ABSTRACT

Feed additives are ingredients that are added to the ration as growth promoters and enhancement of the immune system. Glutamate is a feed additive that improves performance by improving the quality of the small intestine and enhancing the immune system. The purpose of this study was to ~~see-know~~ the effect of ~~giving-including~~ glutamate as a feed additive in improving broiler performance. The material used in this study was broiler strain MB 202 from PT Charoen Phokphand Indonesia as many as 240 birds. The design used was a Completely Randomized Design (CRD) trial design, with ~~six~~6 treatments and ~~four~~4 replications, so that there were 24 cage plots as experimental units. Each experimental unit consisted of 10 chickens. ~~The~~ Glutamate doses ~~is given in groups were~~, A (0.4% commercial glutamate); B (0% glutamate); C (0.2% glutamate); D (0.4% glutamate); E (0.6% glutamate); F (~~0-~~ 8% glutamate). The results ~~indicated~~~~showed~~ that glutamate up to 0.8% had ~~a~~-significant effects ( $P < 0.05$ ) on feed intake (~~4588.68 g~~), body weight gain (~~2693.95 g~~) and feed conversion ratio (1.70%), but it ~~does~~~~did~~ not affect the percentage of carcass, ~~but still affects the percentage of carcass but wasis still within the normal range.~~ ~~The~~ ~~It is~~ ~~concludedsion~~ that ~~giving-including the~~ glutamate ~~to~~in broiler ~~diets~~ can improve broiler performance.

**Commented [E2]:** 0.8%?  
Or 0% and 8%?

**Commented [E3]:** Don't use statistical data in this section.

**Keywords:** Feed additive, Glutamate, Growth promoters, Immune system, Performance

**Commented [E4]:** Arrange base on alphabet

**Commented [E5R4]:** The first letter should be capital

## INTRODUCTION

~~Meat needs continue to increase along with increasing public awareness of the importance of animal protein for the body. One of the animal protein that is consumed by many people is broiler chicken.~~ The broiler is one of the meat-producing livestock that is quite potent in meeting the people's needs for animal protein needs. This is because broiler meat is relatively cheap and easy to obtain compared to other animal proteins. According to ~~(Ensminger et al., (2004)~~ broilers have a fast and efficient growth in converting feed into meat. ~~In addition to several advantages, broilers also have weaknesses, which tend to be susceptible to disease attacks which results in a decrease in broiler performance.~~

~~Addition s~~Supplementation of additional feed is one solution to prevent disease attacks by increasing immunity and growth promoters. According to Madhupriya et al. (2018) Feed additives are ingredients ~~and~~ added to rations to improve animal immunity and performance. Antibiotic Growth Promoter (AGP) is an antibiotic feed additive added to broiler feed. However, the use of AGP in broiler feed is prohibited because residues in broiler meat are harmful to public health. Residual hazards can reduce human resistance to consumption of certain types of antibiotics and other hazards (Ruegg, 2013; Singh et al., 2014). This ~~has~~ led to the enactment of the World Health Organization's regulation regarding the prohibition on the use of antibiotics in animal feed, hence the use of antibiotics as feed additives in broiler feed. So it is necessary to look for additional feed ingredients that are safer for livestock and the community. One safe and harmless feed additive is glutamate.

Glutamate is a feed additive that ~~improves~~ growth promotes and enhances the broiler immune system. Glutamate functions as a constituent of proteins, a substrate in the synthesis of amino acids, as a precursor to several non-essential amino acids and ~~helped~~ the ~~body's~~ metabolism (Young and Ajami, 2000). Giving glutamate to broilers can improve the digestive tract, thus increasing feed efficiency. According to ~~Newholms et al. (2003) and Reeds et al. (2000)~~ Glutamate can improve intestinal development by increasing intestinal villi length and increasing maintenance of intestinal integrity. In addition, according to ~~Li et al. (2007)~~ Glutamate regulates ~~I~~nductible ~~N~~itric ~~O~~xide ~~S~~ynthase (~~I~~NOS) in certain tissues. The expression of INOS ~~was~~ considered ~~as a~~ fundamental mechanism in the protection of parasites,

Commented [36]: c

Commented [E7]: delete

Commented [E8]: rephrase it

Commented [E9]: Please update the references from previous years.

Commented [310]: This is not in references

bacteria, fungi, malignant cells, intracellular protozoa and viruses in different animal species, including mammals and birds. ~~From the description above, it was necessary to do research~~The present study aimed to ~~know-see~~ the effect of giving glutamate as a feed additive in broilers.

## MATERIALS AND METHODS

### ~~Time and Place~~ Place of study

This research was conducted ~~on~~ May-September 2018. Maintenance of broilers was conducted at the Poultry Division Field Laboratory, Faculty of Animal Husbandry, Andalas University.

### Material

The livestock used in this study were broiler MB 202 from PT. Charoen Pokphand Indonesia, which was 1 day old with 240 birds. The chickens ~~were~~ placed randomly into 24 cage plots (open cages), each measuring 1x1 m<sup>2</sup>. Each plot ~~was~~ filled with 10 chickens, equipped with food and drinking water. This broiler cage ~~was~~ also equipped with 1 bulb (35 Watt). The ration ~~was~~ prepared by itself from feed ingredients consisting of corn, CP511 commercial feed, fish meal, soy flour, bran, palm oil and mineral premix. Nutrient content and feed composition ~~was indicated can be seen~~ in Table 1- and Table 2.

### Method

Making glutamate ~~was~~ done by fermentation with *Lactobacillus plantarum*. The nutritional composition of fermentation media used in this study was KH<sub>2</sub>PO<sub>4</sub>, MgSO<sub>4</sub> · 7H<sub>2</sub>O, FeSO<sub>4</sub> · 7H<sub>2</sub>O, MnSO<sub>4</sub> · 4-5H<sub>2</sub>O, 9% sugar cane, 5 μ / L biotin, 90% tofu water, 10% distilled water, water starter and starter 9%. The duration of fermentation in this study was 36 hours at 36°C. After fermentation the media ~~were~~ centrifuged for 20 minutes and 10,000 rpm at a temperature of -4°C. After that, the supernatant was concentrated using an evaporation technique in an oven with a temperature of 40°C for 48 hours. The shrinking material ~~was~~ then calculated for the acid content using glutamate HPLC ~~and ready to be given to broilers~~ (Maslami et al., 2018).

### Maintenance of broilers

**Commented [311]:** Add the ethical approval and any certificate code in this part of article with a separated title.

**Commented [312]:** Add country

**Commented [E13]:** Chang it

**Commented [314]:** The language needs to revision

Maintenance ~~was~~ carried out for ~~five~~ weeks. Feed and drinking water ~~were~~ ~~are not~~ available ~~restricted~~ (*ad libitum*). Glutamate treatment ~~was~~ given to chickens aged ~~two~~ weeks to ~~six~~ weeks. Glutamate ~~was~~ added to drinking water. ~~Weighing~~ ~~The~~ body weight was carried out at the beginning of the study, ~~and~~ every following weeks, and ~~also~~ at the end of the study. Weighing the rest of the feed ~~was~~ done every weekend. Calculations were carried out, ~~for~~ feed intake, body weight gain, feed conversion ratio and carcass percentage.

Formatted: Font: Italic

### Experiment and ~~D~~data ~~A~~analysis

This study used ~~s~~ a completely randomized design. Each treatment was repeated ~~4~~four times. Each treatment and each replication consisted ~~s~~ of 10 birds. The observed variables were ~~just broiler~~ performance ~~traits~~. Production performance included ~~s~~ average feed consumption, body weight gain, final weight, feed conversion value, and carcass percentage. The variable performance included average feed intake, body weight gain, feed conversion ratio and carcass percentage. The data obtained were analyzed using analysis of variance (Analysis of Variance / ANOVA) and if there were differences between treatments were tested further using the DMRT (Duncan Multiple Range Test) tests at a level of 5% (Steel and Torrie, 1991). ~~As a treatment~~, the dose of glutamate in drinking water ~~was~~ A (0.4% Commercial Glutamate), B (0.0% Glutamate), C (0.2% Glutamate), D (0.4% Glutamate), E (0.6% Glutamate) and F (0.8% Glutamate).

Commented [315]: These sentences are same as each other

Commented [316]: What do you mean?

### RESULTS AND ~~D~~DISCUSSION

The results ~~indicated~~ ~~showed~~ that giving of glutamate had a significant effect ( $P < 0.05$ ) on feed intake, body weight gain and feed conversion ratio. While the ~~giving of glutamate did not~~ have a significant effect ( $P > 0.05$ ) on the percentage of the carcass. The average feed intake (FI), body weight gain (BWG), feed conversion ratio (FCR) and carcass percentage (CP) by giving glutamate ~~was indicated~~ ~~can be seen~~ in ~~T~~table 3.

Commented [317]: You should update the references from 2016-2019 (at least 3-4 references should be added)

Commented [318]: Please abbreviated words from methods and materials and then use in all sentences

### Feed ~~I~~intake

~~From T~~table 3. It can be seen that ~~treatment~~ ~~group~~ B was not significantly different ( $P > 0.05$ ) in the treatment of C, D and E (~~Table~~ 3). Judging from the average value there ~~was~~ a tendency to increase feed consumption with an increase in the dose of glutamic acid which can

be the same as the consumption of ration by providing commercial glutamate. Increased consumption of rations with the addition of glutamate caused by glutamate can improve the quality of the digestive tract which can increase the absorption of nutrients. According to Ebadiasl (2011) glutamate in small bowel function as the formation of intestinal mucous villi cells. - ~~Added~~ Bartell and Batal (2007) reported that - glutamate supplementation can improve intestinal development with an increase in the relative weight of the duodenum and jejunum. Improving the development of the small intestine can increase the capacity of the digestive tract to accommodate the volume of the feed so that, many rations can be consumed by broilers. According to Ibrahim (2008), larger intestines can increase ration consumption because there will be a larger volume of food that can be accommodated, digested, and absorbed.

In addition, the increase in ration consumption in this study was due to a decrease in stress on broilers. Reducing stress by adding glutamate because it can synthesize amino acids which played have a role in reducing stress levels. According to Young and Ajami (2000) states that glutamate is an amino acid builder protein which is a precursor for ~~e~~Gamma ~~A~~amino ~~B~~butyric ~~A~~acid (GABA). Glutamate will be converted into GABA if broilers ~~were~~are under stress. According to Wang et al. (2015) GABA serves to reduce stress so that it can maintain appetite and increase the efficiency of the use of body nutrients in livestock. Thus giving glutamate to broiler chickens can reduce stress caused by increased feed consumption. Feed intake in this study was between 4478.75-4588.68 g lower than that obtained by Bezerra et al. (2015) which ~~was~~is 5303 g for 42 days of maintenance. The same was reported by Olubodun et al. (2015) that administration of 0.5% glutamate to broilers resulted in consumption of 4083 g.

#### Body ~~W~~weight ~~G~~gain

~~From Table 3, there was an increased in~~The body weight gain was increased as an increasing dose of glutamate. Giving glutamate in the treatment of F can increase weight gain, but not significantly different ( $P > 0.05$ ) with treatment E. Body weight gain ~~was~~is influenced by feed consumption, the higher the consumption of food, the more weight gain ~~was~~is generated. According to Fadilah (2005) ~~states~~is that one that influences the size of broiler body weight gain ~~was~~is feed intake because, feed intake ~~had~~s a positive correlation with body weight gain. Increased glutamate dose in this treatment ~~is~~caused by an increase in absorption of nutrients in the small intestine. This is supported by the opinion of Shakeri et al. (2014) stated that

Formatted: Heading 4, Left, Line spacing: single

Commented [319]: What do you mean?

administration of glutamate in broiler feed ~~can~~ improve small bowel development, intestinal villi length and nutrient absorption. Glutamate in the small intestine plays a role in the mechanism of intestinal mucosal defense by increasing the ~~mucosal layer~~ ~~mucous layer~~ (Akiba et al., 2009). Increasing the inner surface and surface area of the small intestine will increase the digestion and absorption of feed extracts by the small intestine (Yao et al., 2006).

Increasing weight by giving glutamate because glutamate is a precursor of several non-essential amino acids, so it will meet the needs of several other essential amino acids. According to Blachier et al. (2009) stated that glutamate is a precursor to other non-essential amino acids, such as arginine, glutamine, and proline. With the fulfillment of the need for essential amino acids will increase the growth of broilers in accordance with its function in increasing growth. According to Ajinamoto (2007) administration of glutamate in broiler chicken feed ~~can~~ increase the growth of connective tissue so that, it can increase broiler body weight gain.

**Commented [320]:** It is not clear for understanding  
Rephrase it

**Commented [321]:** It is not clear for understanding  
Rephrase it

The highest weight gain in this study was found in treatment F (2646.45 g / bird) which could match treatment A with commercial glutamate (2638.35 g / bird). The body weight gain in broiler ~~chickens in~~ of this study was higher than that reported by Olubodun et al. (2005) ~~administration of glutamate in broiler chicken feed can increase broiler weight gain with a maintenance period of at~~ 42 days ~~of age~~ (2375 g / ~~head~~bird) ~~compared to those without glutamate administration~~. Increased body weight gain by adding glutamate to broilers ~~compared to without glutamate diet was reported by~~ (Porto et al., (2015). Furthermore, according to Bezerra et al. (2015) the addition of 1.76% glutamate could match the increase in broiler body weight gain without a reduction in crude protein ration. Furthermore, according to Bezerra et al. (2015) addition of 1.76% glutamate ~~can~~ increase ~~broiler~~ body weight gain ~~of broilers~~ with a crude low protein ration.

### Feed ~~c~~Conversion ~~R~~atio

The results ~~indicated~~ ~~showed~~ that treatment ~~groups~~ (D, E and F) gave the same low FCR, even matching treatment A with commercial glutamate. FCR ~~was~~ low by giving glutamate along with increased feed intake and body weight gain. According to Usman (2009); ~~and~~ Zuidhof et al. (2014) the value of FCR ~~was~~ influenced by the amount of feed intake and body weight gain. Another factor is the increase in feed conversion with increasing doses because glutamate can improve the nutritional quality of feed and the digestive system. ~~The~~ ~~It~~ ~~same~~ ~~was~~

reported by Andriyanto et al. (2015) ~~factors~~ the factors that can influence the conversion value of rations are the quality of nutrients. Decreased FCR caused by administration of glutamate ~~can~~ improve the quality of the digestive system by increasing intestinal length thereby increasing absorption of feed nutrients (Olubodun et al., 2015). ~~With~~ increasing absorption of feed nutrients ~~which~~ causes an increase in body weight so that it will increase the efficiency of feed use. A low FCR indicates an increase in feed efficiency (Saputra 2013).

**Commented [322]:** It does not need to reference/s

The lowest FCR in this study was ~~between~~ 1.76-1.70%, lower than Shakeri et al. (2014) stated that the broiler FCR in broilers with administration of glutamate was 1.95%. Similar results were conveyed by Zulkifli et al. (2016) giving 1% glutamate to broiler ~~can~~ reduce FCR to 1.81%. Furthermore, according to Olubodun et al. (2015) added that 0.5% of the mixture of glutamate and glutamine could reduce the FCR of broiler chicken ~~FCR in 1-42 days~~ to 1.78%.

**Commented [323]:** Just one value should be indicated

### Carcass Percentage

The results of statistical analysis ~~indicated~~ showed that the administration of glutamate ~~was~~ did not significantly ~~different~~ affect ( $P > 0.05$ ) ~~with~~ the percentage of broiler carcass. The effect of giving glutamate on carcass percentage can be seen in ~~T~~table 3. The average carcass percentage of each treatment from treatment A to treatment F was 67.23, 68.76, 73.21, 70.70, 70.13, and 70.47% respectively. The percentage of carcass ~~was~~ is the ratio between carcass weight and live weight multiplied by 100%. The percentage of the carcass in this study ~~was~~ is still in the range of 67.23-73.21%. ~~and~~ supported by Resnawati (2004) who stated the percentage of the carcass from heavy cuts was 68-71.8%.

### CONCLUSION

Addition of glutamate up to 0.8% ~~could~~ ~~can~~ improve the performance of broilers ~~by feed intake (4588.68 g), body weight gain (2693.95 g) and feed conversion ratio (1.70%)~~. However, administration of glutamate ~~did~~ does not affect the percentage of the carcass. The percentage of carcass obtained in this study ~~was~~ is still in the normal range between 67.23-73.21%.

**Commented [E24]:** Please do not use data in this section Also make an strong conclusion and message from the most important results

### DECLARATIONS

#### Acknowledgments

~~We are~~ All authors are very grateful to the Minister of Research, Technology and Higher Education of the Republic of Indonesia for support and finance through PMDSU No: No: 1387 /

E4. 2015 and support through the B-Class World Class Professor Program Scheme 123.57 / D2.3 / KP / 2018.

#### Competing interests

The authors declare that they have no competing interests.

#### Author's contribution

All the authors have made contribution to the study.

**Commented [325]:** Indicate the role of each author separately.

#### Consent to publish

All the authors gave their informed consent prior to their inclusion in the study.

#### REFERENCES

Ajinomoto. Animal Nutrition. 2007. Influence of glutamic acid on broiler carcass quality.

**Commented [326]:** Please check our published papers and also the marginal comments to make the references revised.

Akiba Y, Watanabe C, Mizumori M and Kaunitz JD (2009). Luminal l-glutamate enhances duodenal mucosal defense mechanisms via multiple glutamate receptors in rats. *Am. J. Physiol. Gastrointest. Liver Physiol.* 297: 781–791. DOI: 10.1152/ajpgi.90605.2008.

**Commented [327]:** Initial letter of First name?

Andriyanto, Satyaningtjas AS, Yufiadri R, Wulandari R, Darwin VM and Siburian SNA (2015). Performan dan pencernaan pakan ayam broiler yang diberi hormon testosteron dengan dosis bertingkat. *J. Acta Veterinaria Indonesiana*, 3 (1): 29-37. ~~ISSN 2337-3202~~  
~~E-ISSN 2337-4373~~

**Commented [328]:** Full words of journal

Bartell S and Batal A (2007). The effect of supplemental glutamine on growth performance, development of the gastrointestinal tract, and humoral immune response of broilers. *Poultry Sci.*, 86:1940-1947. DOI: 10.1093/ps/86.9.1940.

Bezerra R.M, Costa FGP, Givisiez PEN, Freitas ER, Goulart CC, Santos RA, Souza JG, Brand~ao PA, Lima MR, Melo ML, Rodrigues VP, Nogueira ET and Vieir DVG (2016). Effect of L-glutamate acid supplementation on performance and nitrogen balance of broilers fed low protein diets. *Journal of animal physiology and animal nutrition*, 100: 590–600. DOI: 10.1111/jpn.2405.



- Blachier F, Boutry C, Bos C and Tomé D (2009). Metabolism and functions of L-glutamate in the epithelial cells of the small and large intestines. *Am J Clin Nutr*, 90(3):814-821. DOI: 10.3945/ajcn.2009.27462S.
- Ebadiasl, G. 2011. Effects of supplemental glutamine and glutamate on growth performance, gastrointestinal development, jejunum morphology and *Clostridium perfringens* count in caecum of broilers. Thesis.
- Ensminger ME, Scanes CG and Brant G (2004). *Poultry Science*. 4th Edition. Pearson Prentice Hall, New.
- Ibrahim S (2008). Hubungan ukuran-ukuran usus halus dengan berat badan broiler. *Agripet*, 8(2): 42- 46. DOI: 10.17969/agripet.v8i2.615
- Fadilah Y (2005). *Panduan Mengelola Peternakan Ayam Broiler Komersial*. Agromedia. Pustaka. Jakarta.
- Madhupriya V, Shamsudeen P, Raj Manohar G, Senthilkumar S, Soundarapandiyam V and Moorthy. Phyto feed additives in poultry nutrition – a review, *International Journal of Science. Environment*, ISSN 2278-3687.
- Maslami V, Marlida Y, Mirnawati, Jamsari and Nur YS (2018). Optimization of glutamate production from *Lactobacillus plantarum* originating from Minangkabau fermented food as a feed supplement for broiler. *Pakistan Journal of Nutrition*, 17(7): 336-343, DOI: 10.2923/pjn.2018.336.343.
- Nampoothiri K, Madhavan and Ashok P (1996). Urease Activity in a Glutamate Producing *Brevibacterium* sp. *Procrss Biochemistry*, 31: 471-475. DOI: 10.1016/S0032-9592(95)00090-9.
- Olubodun JO, Zulkifli I, Farjam AS, Hair-Bejo M and Kasim A (2015). Glutamine and glutamic acid supplementation enhances performance of broiler chickens under the hot and humid tropical condition. *Italian Journal of Animal Science*, 14 (3263): 25-29. DOI:10.4081/ijas.2015.3263.

- Porto ML, Givisiez PEN, Saraiva EP, Costa FGP, Moreira Filho ALB, Andrade MFS, Brandão PA and Guerra RR (2015). Glutamic Acid Improves Body Weight Gain and Intestinal Morphology of Broiler Chickens Submitted to Heat Stress. *Brazilian Journal of Poultry Science*, 17 (3): 355-362. DOI: 10.1590/1516-635x1703355-362.
- Reeds PJ, Burrin DG, Stoll B and Jahoor F (2000). Intestinal glutamate metabolism. *J Nutr.*, 130(4): 978-82. DOI: 10.1093/jn/130.4.978S.
- Resnawati. 2004. Bobot potongan karkas dan lemak abdomen ayam ras pedaging yang diberi ransum mengandung tepung cacing tanah. [http://peternakan.litbang.deptan.go.id/user/pros\\_04-75.pdf](http://peternakan.litbang.deptan.go.id/user/pros_04-75.pdf). Accessed: 22 September 2018].
- Ruegg PL (2013). Antimicrobial residues and resistance: Understanding and managing drug usage on dairy farms. University of WI, Dept. of Dairy Science, Madison.
- Saputra WY, Mahfudz LD dan Suthama N (2013). Pemberian pakan single step down dengan penambahan asam sitrat sebagai acidifier terhadap performa pertumbuhan broiler. *Animal Agriculture Journal*, 2(3): 61-72. <http://ejournal-s1.undip.ac.id/index.php/aaj>
- Shakeri M, Zulkifli I, Soleimani AF, Reilly ELO, Eckersall PD, Anna AA, Kumara S and Abdullah FFJ (2014). Response to dietary supplementation of L-glutamine and L-glutamate in broiler chickens reared at different stocking densities under hot, humid tropical conditions. *Poultry Science*, 93:1-9. DOI: 10.3382/ps.2014-03910.
- Singh S., Sanjay S, Neelam T, Nitesh K dan Ritu P (2014). Antibiotic residues: a global challenge. *An International Journal of Pharmaceutical Science. Pharma Science Monitor*, 5 (3):184-197. DOI: 10.5455/vetworld.2008.375-377
- Steel RGD and Torrie JH (1991). *Prinsip dan Prosedur Statistika*. Translated by Bambang Sumantri. PT. Gramedia Pustaka Utama. Jakarta
- Usman (2009). Pertumbuhan ayam buras periode grower melalui pemberian tepung biji buah merah (*Pandanus conoideus* LAMK) sebagai pakan alternatif. *Prosiding Seminar*

Nasional Teknologi Peternakan dan Veteriner. Balai Pengkajian Teknologi Pertanian Papua.

Wang DM, Chacher, Liu HY, Wang JK, Lin J and Liu JX (2015). Effects of  $\gamma$ -aminobutyric acid on feed intake, growth performance and expression of related genes in growing lambs. *Animal*, 9(3):445-448. DOI: 10.1017/S1751731114002651.

Yao Y, Xiaoyan T, Haibo X, Jinchun K, Ming X and Xiaobing W (2006). Effect of choice feeding on performance gastrointestinal development and feed utilization of broilers. *Asian-Aust. J. Anim. Sci.*, 19:91-96. DOI: 10.5713/ajas.2006.91.

Young VR and Ajami AM (2000). Glutamate: an amino acid of particular distinction. *Journal of Nutrition*, 130: 892–900. DOI: 10.1093/jn/130.4.892S.

Zuidhof MJ, Scheider BL, Carney VL, Korver DR and Robinson FE (2014). Growth, efficiency and yield of commercial broilers from 1957, 1978 and 2005. *Poult. Sci*, 93(12): 2970-2982. DOI: 10.3382/ps.2014-04291.

Zulkifli M, Shakeri and Soleimani AF (2016). Dietary supplementation of L-glutamine and L-glutamate in broiler chicks subjected to delayed placement. *Poultry Science*, 1: 1-7. DOI:10.3382/ps/pew267.

Commented [329]: Initial letter of First name?

**Table 1.** Nutrient content of feed ingredients for ~~research rations~~ broiler chickens during 35 days of rearing.

**Commented [330]:** Check it  
**Formatted:** Font: Not Bold

Feed ingredients	Nutritional content (% )and ME Kkal/kg					
	Crude Protein	Crude Fiber	Crude Fat	Ca	P	ME
Corn	9.55	3.8	2.18	0.38	0.33	3300
Bran	10.6	10.84	4.09	0.7	0.09	1592
Fish flour	41	2.8	1.52	5.55	2.6	2580
CP 511 ration	23	1.88	5.87	0.29	0.15	3200
Soybean Meal	40.16	3.58	1.37	0.63	0.32	2240
Coconut oil	0	0	100	0	0	8600
Mineral Premix	0	0	0	5.38	1.14	0

**Commented [331]:** Explain full words under the table  
**Commented [E32]:** Explain full words under the table

**Table 2.** The composition of broiler rations (%)

**Formatted:** Font: Not Bold  
**Formatted:** Font: Not Bold

Feed ingredients	Percentage of ration (%) and nutrient content (%)
Corn	40
Bran	7
Fish flour	17
CP 511 ration	20
Soybean Meal	14
Coconut oil	1.5
Mineral Premix	0.5
Total	100
Crude Protein	21.75
Crude Fiber	3.86
Crude Fat	4.12
Ca	1.50
P	0.83
ME (Kcal/kg)	2952,64

**Table 3.** Effect of glutamate on broiler chicken performance during the study

Treatment	FI (g)	BWG (g)	FCR (%)	CP (%)
A (0.4% Commercial Glutamate)	4591,13 <sup>b</sup> ± 63,67	2688,35 <sup>a</sup> ±24,73	1,71 <sup>b</sup> ±0,03	67,23 ± 3,52
B (0 % Glutamate)	4429,05 <sup>b</sup> ± 77,25	2391,50 <sup>c</sup> ±28,25	1,85 <sup>a</sup> ±0,05	68,76 ± 1,80
C (0.2 % Glutamate)	4478,75 <sup>ab</sup> ± 84,25	2450,63 <sup>c</sup> ±54,96	1,83 <sup>a</sup> ±0,05	73,21 ± 3,62
D (0.4 % Glutamate)	4530,28 <sup>ab</sup> ± 76,08	2573,60 <sup>b</sup> ±63,79	1,76 <sup>b</sup> ±0,07	70,70 ± 3,44
E (0.6 % Glutamate)	4503,70 <sup>ab</sup> ± 103,94	2634,15 <sup>ab</sup> ±31,66	1,71 <sup>b</sup> ±0,04	70,13 ± 3,04
F (0.8% Glutamate)	4588,68 <sup>a</sup> ± 26,24	2693,95 <sup>a</sup> ±74,06	1,70 <sup>b</sup> ±0,05	70,47 ± 4,39

**Commented [333]:** Indicate the study period

**Formatted:** Font: Not Bold

**Formatted:** Font: Not Bold

**Formatted:** Font: Not Bold

**Commented [334]:** BW BWG?  
Which one is true?  
Please check it and make it correct in text of article.

**Commented [E35]:** Explain full words under the table

**Commented [336]:** Delete it  
For all data

**Commented [337]:** Make it correct for all data!

**Effect of Supplementation of Glutamate as a Feed Additive Feed in Performance of Improving Broiler Chickens Performance**

Vebera Maslami<sup>1</sup>, Mirnawati<sup>2</sup>, Jamsari<sup>3</sup>, Yuliaty Shafan Nur<sup>2</sup> and Yetti Marlida<sup>2\*</sup>

<sup>1</sup>Department of Agricultural Science, Faculty of Agriculture, Andalas University, 25163, Indonesia

<sup>2</sup>Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, Andalas University, 25163, Indonesia

<sup>3</sup>Department of Crop Science, Faculty of Agriculture, University Andalas, 25163, Indonesia

\*Corresponding author's E-mail: [yettimarlida@ansci.unand.ac.id](mailto:yettimarlida@ansci.unand.ac.id)

**ABSTRACT**

Feed additives are ingredients that are added to the ration as growth promoters and enhancement of the immune system. Glutamate is a feed additive that improves performance by improving the quality of the small intestine and enhancing the immune system. The purpose of this study was to know the effect of including glutamate as a feed additive in improving broiler performance. The material used in this study was broiler strain MB 202 from PT Charoen Phokphand Indonesia as many as 240 birds. The design used was a Completely Randomized Design (CRD) trial design, with six treatments and four replications, so that there were 24 cage plots as experimental units. Each experimental unit consisted of 10 chickens. The Glutamate doses in groups were, A (0.4% commercial glutamate); B (0% glutamate); C (0.2% glutamate); D (0.4% glutamate); E (0.6% glutamate); F (0.8% glutamate). The results indicated that glutamate up to 0.8% had significant effects on feed intake body weight gain and feed conversion ratio (1.70%), but it did not affect the percentage of carcass. It is concluded that including the glutamate in broiler diets can improve broiler performance.

Commented [31]: Add percent

Commented [32]: During the age

**Key words:** Feed additive, Glutamate, Growth promoters, Immune system, Performance

## INTRODUCTION

The broiler is one of the meat-producing livestock that is quite potent in meeting the people's needs for animal protein needs. This is because broiler meat is relatively cheap and easy to obtain compared to other animal proteins. According to [Ensminger et al. \(2004\)](#) broilers have a fast and efficient growth in converting feed into meat. In addition, broilers also have weaknesses which tend to be susceptible to disease attacks which results in a decrease in broiler performance.

Supplementation of additional feed is one solution to prevent disease attacks by increasing immunity and growth promoters. According to [Madhupriya et al. \(2018\)](#) Feed additives are ingredients and added to rations to improve animal immunity and performance. Antibiotic Growth Promoter (AGP) is an antibiotic feed additive added to broiler feed. However, the use of AGP in broiler feed is prohibited because residues in broiler meat are harmful to public health. Residual hazards ~~can~~ could reduce human resistance to consumption of certain types of antibiotics and other hazards ([Ruegg, 2013](#); [Singh et al., 2014](#)). This had led to the enactment of the World Health Organization's regulation regarding the prohibition on the use of antibiotics in animal feed, hence the use of antibiotics as feed additives in broiler feed. So it is necessary to look for additional feed ingredients that are safer for livestock and the community. One safe and harmless feed additive is glutamate.

Glutamate is a feed additive that [can](#) improves growth promotes and enhances the broiler immune system. Glutamate functions as a constituent of proteins, a substrate in the synthesis of amino acids, as a precursor to several non-essential amino acids and helped the metabolism ([Young and Ajami, 2000](#)). Giving glutamate to broilers ~~can~~ could improve the digestive tract, thus increasing feed efficiency. According to [Newholms et al. \(2003\)](#); [Reeds et al. \(2000\)](#) glutamate ~~can~~ could improve intestinal development by increasing intestinal villi length and increasing maintenance of intestinal integrity. In addition, according to [Li et al. \(2007\)](#) glutamate ~~regulates~~ regulated Inductible Nitric Oxide Synthase (iNOS) in certain tissues. The expression of iNOS was considered as a fundamental mechanism in the protection of parasites, bacteria, fungi, malignant cells, intracellular protozoa and viruses in different animal species, including mammals and birds. ~~It was~~ The present study aimed to know the effect of giving glutamate as a feed additive in broilers.

## MATERIALS AND METHODS

Commented [33]: References?

Commented [a4]: Add the **ethical approval** and any certificate code in this part of article with a separated title.

### Place of study

This research was conducted on May-September 2018. Maintenance of broilers was conducted at the Poultry Division Field Laboratory, Faculty of Animal Husbandry, Andalas university, Indonesia.

### Materials

A total of 240 one-day old male broiler chicks (MB 202 from) were purchased from PT Charoen Pokphand, Indonesia. The chickens were placed randomly into 24 cage plots (open cages), each measuring 1×1 m<sup>2</sup>. Each plot was filled with 10 chickens, equipped with food and drinking water. This broiler cage was also equipped with 1 bulb (35 Watt). The ration was prepared by itself from feed ingredients consisting of corn, CP511 commercial feed, fish meal, soy flour, bran, palm oil and mineral premix. Nutrient content and feed composition was indicated in table 1 and table 2.

**Commented [35]:** What is it?

**Commented [36]:** What is different between these tables?  
Please indicate why two able

### Method

Making glutamate was done by fermentation with *Lactobacillus plantarum*. The nutritional composition of fermentation media used in this study was KH<sub>2</sub>PO<sub>4</sub>, MgSO<sub>4</sub> · 7H<sub>2</sub>O, FeSO<sub>4</sub> · 7H<sub>2</sub>O, MnSO<sub>4</sub> · 4-5H<sub>2</sub>O, 9% sugar cane, 5 μ / L biotin, 90% tofu water, 10% distilled water, water starter and starter 9%. The duration of fermentation in this study was 36 hours at 36°C. After fermentation the media were centrifuged for 20 minutes and 10,000 rpm at a temperature of -4°C. After that, the supernatant was concentrated using an evaporation technique in an oven with a temperature of 40°C for 48 hours. The shrinking material was then calculated for the acid content using glutamate HPLC (Maslami et al., 2018).

### Maintenance of broilers

Maintenance was carried out for five weeks. Feed and drinking water were available ad libitum. Glutamate treatment was given to chickens aged two weeks to six weeks. Glutamate was added to drinking water. The body weight was carried out at the beginning of the study and every following week, and also at the end of the study. Weighing the rest of the feed was done every weekend. Calculations were carried out for Feed Intake (FI), Body Weight Gain (BWG), Feed Conversion Ratio (FCR) and Carcass Percentage (CP).



### Experiment and data analysis

This study used a completely randomized design. Each treatment was repeated four times. Each treatment and each replication consisted of 10 birds. The observed variables were just performance traits. The variable performance included average feed intake, body weight gain, feed conversion ratio and carcass percentage. The data obtained were analyzed using analysis of variance (Analysis of Variance / ANOVA) and if there were differences between treatments were tested further using the DMRT (Duncan Multiple Range Test) tests at a level of 5% (Steel and Torrie, 1991). The dose of glutamate in drinking water is A (0.4% commercial glutamate), B (0.0% glutamate), C (0.2% glutamate), D (0.4% glutamate), E (0.6% glutamate) and F (0.8% glutamate).

### RESULTS AND DISCUSSION

The results indicated that giving of glutamate had a significant effect ( $P < 0.05$ ) on feed intake, body weight gain and feed conversion ratio. While the giving of glutamate did not have a significant effect ( $P > 0.05$ ) on the percentage of the carcass. The average FI, BWG, FCR and CP by giving glutamate was indicated in table 3.

#### Feed intake

It can be seen that group B was not significantly different ( $P > 0.05$ ) in the treatment of C, D and E (Table 3). Judging from the average value there was a tendency to increase feed consumption with an increase in the dose of glutamic acid which can be the same as the consumption of ration by providing commercial glutamate. Increased consumption of rations with the addition of glutamate caused by glutamate can improve the quality of the digestive tract which can increase the absorption of nutrients. According to Ebadiasl (2011) glutamate in small bowel function as the formation of intestinal mucous villi cells. Bartell and Batal (2007) reported that glutamate supplementation ~~can~~ could improve intestinal development with an increase in the relative weight of the duodenum and jejunum. Improving the development of the small intestine can increase the capacity of the digestive tract to accommodate the volume of the feed so that, many different rations can be consumed by broilers. According to Ibrahim (2008), larger intestines ~~can~~ could increase ration consumption because ~~there will be~~ a larger volume of food ~~that~~ can be accommodated, digested, and absorbed.

In addition, the increase in ration consumption in this study was due to a decrease in stress on broilers. Reducing stress by adding glutamate because it can synthesize amino acids which have a role in reducing stress levels. According to Young and Ajami (2000) states that glutamate is an amino acid builder protein which is a precursor for Gamma AminoButyric Acid (GABA). Glutamate will be converted into GABA if broilers were under stress. According to Wang et al. (2015) and Lener et al. (2017) GABA serves to reduce stress so that it can maintain appetite and increase the efficiency of the use of body nutrients in livestock. Thus giving glutamate to broiler chickens can reduce stress caused by increased-increasing feed consumption. Feed intake in this study was between 4478.75-4588.68 g lower than that obtained by Bezerra et al. (2015) which was 5303 g for 42 days of maintenance. The same was Similar study was reported-presented by Olubodun et al. (2015) that administration of 0.5% of glutamate to broilers resulted-and indicated in consumption of 4083 g feed.

### **Body weight gain**

The body weight gain was increased as an increasing dose of glutamate. Giving glutamate in the treatment of F can increase weight gain, but not significantly different ( $P > 0.05$ ) with treatment E. Body weight gain was influenced by feed consumption, the higher the consumption of food, the more weight gain was generated. According to Fadilah (2005) feed consumption has-had a positive effect on weight gain. Increased glutamate dose in this treatment caused by an increase in absorption of nutrients in the-small intestine. This is supported by the opinion of Shakeri et al. (2014) stated that administration of glutamate in broiler feed could improve small bowel development, intestinal villi length and nutrient absorption. Glutamate in the small intestine plays a role in the mechanism of intestinal mucosal defense by increasing the mucosal layer (Akiba et al., 2009). Increasing the inner surface and surface area of the small intestine will increase the digestion and absorption of feed extracts by the small intestine (Yao et al., 2006).

Giving glutamate can increase broiler weight because glutamate is a precursor for non-essential amino acids, so it will meet the needs of several other essential amino acids. According to Blachier et al. (2009) stated that glutamate is a precursor to other non-essential amino acids, such as arginine, glutamine, and proline. With the fulfillment of the need for non-essential amino acids will increase the growth of broilers. According to Ajinamoto (2007); Maslami et al. (2018)

administration of glutamate in broiler chicken feed could increase the growth of connective tissue so that, it can increase broiler body weight gain.

The highest weight gain in this study was found in treatment F (2646.45 g / bird) which could match treatment A with commercial glutamate (2638.35 g / bird). The body weight gain in broiler chickens of this study was higher than that reported by Olubodun et al. (2005) at 42 days of age (2375 g / bird). Increased body weight gain by adding glutamate to broilers diet was reported by Porto et al. (2015). Furthermore, according to Bezerra et al. (2015) the addition of 1.76% glutamate could match the increase in broiler body weight gain without a reduction in crude protein ration. ~~Furthermore~~ ~~Moreover~~, according to ~~Bezerra et al. (2015)~~ ~~to this study~~ addition of 1.76% glutamate could increase body weight gain of broilers with a crude low protein ration.

#### Feed conversion ratio

The results indicated that treatment groups (D, E and F) gave the same low FCR, even matching treatment A with commercial glutamate. FCR was low by giving glutamate along with increased feed intake and body weight gain. A low FCR indicates an increase in feed efficiency (Razak et al., 2016). According to Usman (2009) and Zuidhof et al. (2014) the values of FCR ~~was~~ ~~were~~ influenced by the amount of feed intake and body weight gain. Another factor is the increase in feed conversion with increasing doses because glutamate can improve the nutritional quality of feed and the digestive system. ~~It was reported by Andriyanto et al. (2015) the factors could influence the conversion value of rations are the quality of nutrients~~. Decreased FCR caused by administration of glutamate could improve the quality of the digestive system by increasing intestinal length thereby increasing absorption of feed nutrients (Olubodun et al., 2015). Increasing absorption of feed nutrients causes an increase in body weight so that it will increase the efficiency of feed use. ~~A low FCR indicates an increase in feed efficiency~~.

The lowest FCR in this study was 1.70%, lower than Shakeri et al. (2014) stated that the FCR in broilers with administration of glutamate was 1.95%. Similar results were conveyed by Zulkifli et al. (2016) giving 1% glutamate to broiler could reduce FCR to 1.81%. Furthermore, according to Olubodun et al. (2015) ~~that~~ added ~~that~~ 0.5% of the mixture of glutamate and glutamine ~~to broiler chicken diet and could reported~~ reduce ~~the in~~ FCR of broiler chicken to (1.78%).

Commented [37]: It is not clear for understanding

Commented [38]: It is not percent

Commented [39]: It is not percent

Commented [310]: It is not percent

Commented [311]: It is not percent

### Carcass percentage

The results of statistical analysis indicated that the administration of glutamate did not significantly affect ( $P > 0.05$ ) on the percentage of broiler carcass. The effect of giving glutamate on carcass percentage ~~can be seen~~ presented in table 3. The average carcass percentage of each treatment from treatment A to treatment F was 67.23, 68.76, 73.21, 70.70, 70.13, and 70.47% respectively. The percentage of carcass was the ratio between carcass weight and live weight multiplied by 100%. The percentage of the carcass in this study was in the range of 67.23-73.21% ~~which was~~ supported by Resnawati (2004) who stated the percentage of the carcass from heavy cuts was 68-71.8%.

### CONCLUSION

Addition of glutamate up to 0.8% could improve the performance of broilers. However, administration of glutamate did not affect the percentage of the carcass. ~~The percentage of carcass obtained in this study was in the normal.~~

### DECLARATIONS

#### Acknowledgments

All authors are very grateful to the Minister of Research, Technology and Higher Education of the Republic of Indonesia for support and finance through PMDSU No: No: 1387 / E4. 2015 and support through the B-Class World Class Professor Program Scheme 123.57 / D2.3 / KP / 2018.

#### Competing interests

The authors declare that they have no competing interests.

#### Author's contribution

Vebera Maslami and Yetti Marlida conducted the research and ~~compile~~ prepared data and wrote the articles. Mirnawati, Jamsari and Yuliaty Shafan Nur checked and ~~compiled~~ confirmed the final form of article and revised the article.

#### Consent to publish

All ~~the~~ authors gave their informed consent prior to their inclusion in the study.

**Commented [312]:** This methods and it is not results.  
Move it

**Commented [313]:** Is it related to Glutamate?

**Commented [314]:** What does it mean?  
Change it

**Commented [315]:** Add another important sentence about conclusion like a message for future studies

**Commented [316]:** the age of rearing should be indicated

**Commented [a17]:** Who write the article and do statistical analysis?

## REFERENCES

- Ajinomoto, AN- (2007). Influence of glutamic acid on broiler carcass quality. Available at: <http://www.lysine.com/pdf/poultry/pr19.pdf>. (Accessed August 20, 2015).
- Akiba Y, Watanabe C, Mizumori M and Kaunitz JD (2009). Luminal l-glutamate enhances duodenal mucosal defense mechanisms via multiple glutamate receptors in rats. *American Journal of Physiology-Gastrointest Liver Physiology*, 297: 781–791. DOI: <https://doi.org/10.1152/ajpgi.90605.2008>.
- Andriyanto, Satyaningtjas AS, Yufiadri R, Wulandari R, Darwin VM and dan-Dan Sibirian SNA (2015). Performan dan pencernaan pakan ayam broiler yang diberi hormon testosteron dengan dosis bertingkat. *Journal - Acta Veterinaria Indonesiana*, 3 (1): 29-37.
- Bartell S and Batal A (2007). The effect of supplemental glutamine on growth performance, development of the gastrointestinal tract, and humoral immune response of broilers. *Poultry Sci.*, 86:1940-1947. DOI: 10.1093/ps/86.9.1940.
- Bezerra R-M, Costa FGP, Givisiez FGP, Givisiez PEN, Freitas ER, Goulart CC, Santos RA, Souza JG, Brand-ao PA, Lima MR, Melo ML, Rodrigues VP, Nogueira ET and Vieir DVG (2016). Effect of L-glutamate acid supplementation on performance and nitrogen balance of broilers fed low protein diets. *Journal of animal physiology and animal nutrition*, 100: 590–600. DOI: 10.1111/jpn.2405.
- Blachier F, Boutry C, Bos C and Tomé D (2009). Metabolism and functions of L-glutamate in the epithelial cells of the small and large intestines. *Am J Clin Nutr*, 90(3):814-821 DOI: 10.3945/ajcn.2009.27462S.
- Ebadiasl, G- (2011). Effects of supplemental glutamine and glutamate on growth performance, gastrointestinal development, jejunum morphology and Clostridium perfringens count in caecum of broilers. Thesis.
- Ensminger ME, Scanes CG and Brant G (2004). *Poultry Science*. 4th Edition. Pearson Prentice Hall, New.
- Ibrahim S (2008). Hubungan ukuran-ukuran usus halus dengan berat badan broiler. *Agripet*, 8(2): 42- 46. DOI: 10.17969/agripet.v8i2.615

**Commented [a18]:** Please write the reference according to the journal format. You can check one of journal's articles using this link:

[http://iwpr.science-line.com/attachments/article/49/J%20World%20Poult%20Res%209\(2\)%20109-116,%202019.pdf](http://iwpr.science-line.com/attachments/article/49/J%20World%20Poult%20Res%209(2)%20109-116,%202019.pdf)

**Commented [a19]:** Change like sample: [http:// www.bradley.edu/computorg/psiohi/DS9/ep/503r.htm](http://www.bradley.edu/computorg/psiohi/DS9/ep/503r.htm); Internet; accessed 8 October 199.

**Formatted:** No underline

**Field Code Changed**

**Formatted:** No underline

**Commented [a20]:** Use comma

**Commented [321]:** DOI: <https://>  
Please make this format for all references

**Commented [a22]:** Use English words

**Commented [a23]:** Use English words

**Commented [a24]:** Full name of journal

**Commented [a25]:** When the number of authors is more than 10, et al is used.

**Commented [a26]:** English words

**Commented [a27]:** Mention page/s or web address

**Commented [328]:** pages

Fadilah Y (2005). Panduan Mengelola Peternakan Ayam Broiler Komersial. Agromedia. Pustaka. Jakarta.

**Commented [329]:** Mention page/s or web address

Lener MS, Niciu MJ, Ballard ED, Park M, Park LT, Nugent A and Zarate CA. (2017). Glutamate and GABA systems in the pathophysiology of major depression and antidepressant response to ketamine. *Biological Psychiatry*. 81(10): 886–897. DOI: 10.1016/j.biopsych.2016.05.005

Li P., Yin YL and Li D. 2007. Amino acids and immune function. *British Journal of Nutrition* 98: 237- 252. DOI: 10.1017/S000711450769936X

Madhupriya V, Shamsudeen P, Raj Manohar G, Senthilkumar S, Soundarapandiyan V and Moorthy. Phyto feed additives in poultry nutrition – a review, *International Journal of Science, Environment, ISSN-2278-3687*.

**Commented [330]:** Year?

**Commented [331]:** ?

Maslami V, Marlida Y, Mirnawati, Jamsari and Nur YS (2018). Optimization of glutamate production from *Lactobacillus plantarum* originating from Minangkabau fermented food as a feed supplement for broiler. *Pakistan Journal of Nutrition*, 17(7): 336-343. DOI: 10.2923/pjn.2018.336.343.

**Commented [332]:** Check the anme of journal and add the volume and issue and pages

Maslami V, Marlida Y, Mirnawati, Jamsari, Nur YS, Adzitey F and Huda N. (2018). A review on potential of glutamate producing lactic acid bacteria of west Sumatera's fermented food origin, As feed additive for broiler chicken. *The Journal of World's Poultry Research*. 8(4):120-126. PII: S2322455X1800017-8

Nampoothiri K, Madhavan and Ashok P (1996). Urease Activity in a Glutamate Producing *Brevibacterium* sp. *Proccrs Biochemistry*, 31: 471-475. DOI: 10.1016/S0032-9592(95)00090-9.

Olubodun JO, Zulkifli I, Farjam AS, Hair-Bejo M and Kasim A (2015). Glutamine and glutamic acid supplementation enhances performance of broiler chickens under the hot and humid tropical condition. *Italian Journal of Animal Science*, 14 (3263): 25-29. DOI:10.4081/ijas.2015.3263.

Porto ML, Givisiez PEN, Saraiva EP, Costa FGP, Moreira Filho ALB, Andrade MFS, Brandão PA and Guerra RR (2015). Glutamic Acid Improves Body Weight Gain and Intestinal Morphology of Broiler Chickens Submitted to Heat Stress. *Brazilian Journal of Poultry Science*, 17 (3): 355-362. DOI: 10.1590/1516-635x1703355-362.

Razak AD Kiramang K dan Hidayat MN. 2016. Pertambahan bobot badan, konsumsi ransum dan konversi ransum ayam ras pedaging yang diberikan tepung daun sirih (*Piper Betle Linn*) sebagai imbuhan pakan. *Jurnal Ilmu dan Industri Peternakan*, 3: 135-147. ISSN: 2442-2541

Reeds PJ, Burrin DG, Stoll B and Jahoor F (2000). Intestinal glutamate metabolism. *J Nutr.*, 130(4): 978-82. DOI: 10.1093/jn/130.4.978S.

Commented [333]: Full words

Resnawati (2004). Bobot potongan karkas dan lemak abdomen ayam ras pedaging yang diberi ransum mengandung tepung cacing tanah. [http://peternakan.litbang.deptan.go.id/user/pros\\_04-75.pdf](http://peternakan.litbang.deptan.go.id/user/pros_04-75.pdf). (Accessed: 22 September 2018).

Ruegg PL (2013). Antimicrobial resi-dues and resistance: Understand-ing and managing drug usage on dairy farms. University of WI, Dept. of Dairy Science, Madison.

Commented [334]: Full words

Saputra WY, Mahfudz LD dan Suthama N (2013). Pemberian pakan single step down dengan penambahan asam sitrat sebagai acidifier terhadap performa pertumbuhan broiler. *Animal Agriculture Journal*, 2(3): 61-72. Available at: <http://ejournal-s1.undip.ac.id/index.php/aaj>

Shakeri M, Zulkifli I, Soleimani AF, Reilly ELO, Eckersall PD, Anna AA, kumara S and Abdullah FFJ (2014). Response to dietary supplementation of L-glutamine and L-glutamate in broiler chickens reared at different stocking densities under hot, humid tropical conditions. *Poultry Science*, 93:1-9. DOI: 10.3382/ps.2014-03910.

Singh S., Sanjay S, Neelam T, Nitesh K dan Ritu P (2014). Antibiotic residues: a global challenge. *An International Journal of Pharmaceutical Science. Pharma Science Monitor*, 5 (3):184-197. DOI: 10.5455/vetworld.2008.375-377

Steel RGD and Torrie JH (1991). Prinsip dan Prosedur Statistika. Translated by Bambang Sumantri. PT. Gramedia Pustaka Utama. Jakarta

Commented [335]: Mention page/s or web address

Usman (2009). Pertumbuhan ayam buras periode grower melalui pemberian tepung biji buah merah (*Pandanus conoideus* LAMK) sebagai pakan alternatif. Prosiding Seminar Nasional Teknologi Peternakan dan Veteriner. Balai Pengkajian Teknologi Pertanian Papua.

Wang DM, Chacher, Liu HY, Wang JK, Lin J and Liu JX (2015). Effects of  $\gamma$ -aminobutyric acid on feed intake, growth performance and expression of related genes in growing lambs. *Animal*, 9(3): 445-448. DOI: 10.1017/S1751731114002651.

Yao Y, Xiaoyan T, Haibo X, Jinchen K, Ming X and Xiaobing W (2006). Effect of choice feeding on performance gastrointestinal development and feed utilization of broilers.

[Asian-Aust. J. Anim. Sci.](#), 19:91-96. DOI: 10.5713/ajas.2006.91.

**Commented [336]:** Full words

Young VR and Ajami AM (2000). Glutamate: an amino acid of particular distinction. *Journal of Nutrition*, 130: 892–900. DOI: 10.1093/jn/130.4.892S.

Zuidhof MJ, Scheider BL, Carney VL, Korver DR and Robinson FE (2014). Growth, efficiency and yield of commercial broilers from 1957, 1978 and 2005. [Poult. Sci.](#), 93(12): 2970-2982. DOI: 10.3382/ps.2014-04291.

**Commented [a37]:** Full name of journal

**Commented [338]:** Full words

Zulkifli M, Shakeri M and Soleimani AF (2016). Dietary supplementation of L-glutamine and L-glutamate in broiler chicks subjected to delayed placement. *Poultry Science*, 1: 1-7. DOI:10.3382/ps/pew267.



**Table 1.** Nutrient content of feed ingredients for broiler chickens

Feed ingredients	Nutritional content ( % ) and energy metabolism (Kkal/kg)					
	Crude Protein	Crude Fiber	Crude Fat	Ca	P	ME
Corn	9.55	3.8	2.18	0.38	0.33	3300
Bran	10.6	10.84	4.09	0.7	0.09	1592
Fish flour	41	2.8	1.52	5.55	2.6	2580
CP 511 ration	23	1.88	5.87	0.29	0.15	3200
Soybean Meal	40.16	3.58	1.37	0.63	0.32	2240
Coconut oil	0	0	100	0	0	8600
Mineral Premix	0	0	0	5.38	1.14	0

Energy metabolism: ME; calcium: Ca; Phosphor: P

**Commented [E39]:** Explain full words under the table  
Author: I explained

**Commented [340]:** Full words under the table

**Table 2.** The composition of [rations of broiler rations chickens](#)

Feed ingredients	Percentage of ration (%) and nutrient content (%)
Corn	40
Bran	7
Fish flour	17
CP 511 ration	20
Soybean Meal	14
Coconut oil	1.5
Mineral Premix	0.5
Total	100
Crude Protein	21.75
Crude Fiber	3.86
Crude Fat	4.12
Ca	1.50

**Commented [341]:** Full words under the table

**Commented [342]:** Full words under the table

P	0.83
ME (Kcal/kg)	2952,64

Commented [343]: Full words under the table

Commented [344]: Full words under the table

**Table 3.** Effect of dietary inclusion of glutamate on broiler chicken performance for 35 days

Commented [345]: Percent?

Treatment	FI (g)	BWG (g)	FCR (%)	CP (%)
A (0.4% Commercial Glutamate)	4591.13 <sup>a</sup> ± 63.67	2688.5 <sup>a</sup> ±24.73	1.71 <sup>b</sup> ±0.03	67.23 ± 3.52
B (0 % Glutamate)	4429.05 <sup>b</sup> ± 77.25	2391.50 <sup>c</sup> ±28.25	1.85 <sup>a</sup> ±0.05	68.76 ±1.80
C (0.2 % Glutamate)	4478.75 <sup>ab</sup> ± 84.25	2450.63 <sup>c</sup> ±54.96	1.83 <sup>a</sup> ±0.05	73.21 ±3.62
D (0.4 % Glutamate)	4530.28 <sup>ab</sup> ± 76.08	2573.60 <sup>b</sup> ±63.79	1.76 <sup>b</sup> ±0.07	70.70 ± 3.44
E (0.6 % Glutamate)	4503.70 <sup>ab</sup> ± 103.94	2634.15 <sup>ab</sup> ±31.66	1.71 <sup>b</sup> ±0.04	70.13 ± 3.04
F (0.8% Glutamate)	4588.68 <sup>a</sup> ± 26.24	2693.95 <sup>a</sup> ±74.06	1.70 <sup>b</sup> ±0.05	70.47 ± 4.39

Feed intake: FI; body weight gain: BWG; feed conversion ratio: FCR; Carcass percentage: CP

**Effect of Supplementation of Glutamate as a Feed Additive Feed-on Performance of Improving Broiler Chickens Performance**

Vebera Maslami<sup>1</sup>, Mirnawati<sup>2</sup>, Jamsari<sup>3</sup>, Yuliaty Shafan Nur<sup>2</sup> and Yetti Marlida<sup>2\*</sup>

<sup>1</sup>Department of Agricultural Science, Faculty of Agriculture, Andalas University, 25163, Indonesia

<sup>2</sup>Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, Andalas University, 25163, Indonesia

<sup>3</sup>Department of Crop Science, Faculty of Agriculture, University Andalas, 25163, Indonesia

\*Corresponding author's E-mail: [yettimarlida@ansci.unand.ac.id](mailto:yettimarlida@ansci.unand.ac.id)

**ABSTRACT**

Feed additives are ingredients that are added to the ration as growth promoters and enhancement of the immune system. Glutamate is a feed additive that improves performance by improving the quality of the small intestine and enhancing the immune system. The purpose of this study was to know the effect of including glutamate as a feed additive in improving broiler performance. The material used in this study was broiler strain MB 202 from PT Charoen Phokphand Indonesia as many as 240 birds. The design used was a Completely Randomized Design (CRD) trial design, with six treatments and four replications, so that there were 24 cage plots as experimental units. Each experimental unit consisted of 10 chickens. The Glutamate doses in groups were, A (0.4% commercial glutamate); B (0% glutamate); C (0.2% glutamate); D (0.4% glutamate); E (0.6% glutamate); F (0.8% glutamate). The results indicated that glutamate up to 0.8% had significant effects on feed intake body weight gain and feed conversion ratio (1.70%), but it did not affect the percentage of carcass. It is concluded that including the 0.8% glutamate in broiler diets can improve broiler performance with 35 days of maintenance.

**Key words:** Feed additive, Glutamate, Growth promoters, Immune system, Performance

## INTRODUCTION

The broiler is one of the meat-producing livestock that is quite potent in meeting the people's needs for animal protein needs. This is because broiler meat is relatively cheap and easy to obtain compared to other animal proteins. According to [Ensminger et al. \(2004\)](#) broilers have a fast and efficient growth in converting feed into meat. In addition, broilers also have weaknesses which tend to be susceptible to disease attacks which results in a decrease in broiler performance.

Supplementation of additional feed is one solution to prevent disease attacks by increasing immunity and growth promoters. According to [Madhupriya et al. \(2018\)](#) Feed additives are ingredients and ~~added~~ to rations to improve animal immunity and performance. Antibiotic Growth Promoter (AGP) is an antibiotic feed additive added to broiler feed. However, the use of AGP in broiler feed is prohibited because residues in broiler meat are harmful to public health. Residual hazards ~~can~~ could reduce human resistance to consumption of certain types of antibiotics and other hazards ([Ruegg, 2013](#); [Singh et al., 2014](#)). This had led to the enactment of the World Health Organization's regulation regarding the prohibition on the use of antibiotics in animal feed, hence the use of antibiotics as feed additives in broiler feed. So it is necessary to look for additional feed ingredients that are safer for livestock and the community. One safe and harmless feed additive is glutamate.

Glutamate is a feed additive that can ~~improves~~ growth promotes and enhances the broiler immune system. Glutamate functions was known as a constituent of proteins, a substrate in the synthesis of amino acids, as a precursor to several non-essential amino acids and helped the metabolism ([Young and Ajami, 2000](#)). According to [Newholms et al. \(2003\)](#); [Reeds et al. \(2000\)](#) glutamate ~~can~~ could improve intestinal conditions by increasing intestinal villi length and increasing maintenance of intestinal integrity. In addition, according to [Li et al. \(2007\)](#) glutamate ~~regulates~~ regulated Inductible Nitric Oxide Synthase (iNOS) in certain tissues. The expression of iNOS was considered as a fundamental mechanism in the protection of parasites, bacteria, fungi, malignant cells, intracellular protozoa and viruses in different animal species, including mammals and birds. ~~It was~~ The present study aimed to know the effect of giving glutamate as a feed additive in broilers.

## MATERIALS AND METHODS

### Place of study

This research was conducted on May-September 2018. Maintenance of broilers was conducted at the Poultry Division Field Laboratory, Faculty of Animal Husbandry, Andalas university, Indonesia.

### Materials

A total of 240 one-day old male broiler chicks (MB 202 from) were purchased from PT Charoen Pokphand, Indonesia. The chickens were placed randomly into 24 cage plots (open cages), each measuring 1×1 m<sup>2</sup>. Each plot was filled with 10 chickens, equipped with food and drinking water. This broiler cage was also equipped with 1 bulb (35 Watt). The ration was prepared by itself from feed ingredients consisting of corn, commercial feed (CP511), fish meal, soy flour, bran, palm oil and mineral premix. Nutrient content and feed composition was indicated in table 1 and table 2.

### Method

Making glutamate was done by fermentation with *Lactobacillus plantarum*. The nutritional composition of fermentation media used in this study was KH<sub>2</sub>PO<sub>4</sub>, MgSO<sub>4</sub> · 7H<sub>2</sub>O, FeSO<sub>4</sub> · 7H<sub>2</sub>O, MnSO<sub>4</sub> · 4-5H<sub>2</sub>O, 9% sugar cane, 5 μ / L biotin, 90% tofu water, 10% distilled water, water starter and starter 9%. The duration of fermentation in this study was 36 hours at 36°C. After fermentation the media were centrifuged for 20 minutes and 10,000 rpm at a temperature of -4°C. After that, the supernatant was concentrated using an evaporation technique in an oven with a temperature of 40°C for 48 hours. The shrinking material was then calculated for the acid content using glutamate HPLC (Maslami et al., 2018).

### Maintenance of broilers

Maintenance was carried out for five weeks. Feed and drinking water were available ad libitum. Glutamate treatment was given to chickens aged two weeks to six weeks. Glutamate was added to drinking water. The body weight was carried out at the beginning of the study and every following week, and also at the end of the study. Weighing the rest of the feed was done every weekend. Calculations were carried out for Feed Intake (FI), Body Weight Gain (BWG), Feed Conversion Ratio (FCR) and Carcass Percentage (CP).

**Commented [31]:** What is it?  
Author: CP 511 Broiler trademark

Please mention the corporation and also the country in parentheses

### Experiment and data analysis

This study used a completely randomized design. Each treatment was repeated four times. Each treatment and each replication consisted of 10 birds. The observed variables were just performance traits. The variable performance included average feed intake, body weight gain, feed conversion ratio and carcass percentage. The data obtained were analyzed using analysis of variance (Analysis of Variance / ANOVA) and if there were differences between treatments were tested further using the DMRT (Duncan Multiple Range Test) tests at a level of 5% (Steel and Torrie, 1991). The dose of glutamate in drinking water is A (0.4% commercial glutamate), B (0.0% glutamate), C (0.2% glutamate), D (0.4% glutamate), E (0.6% glutamate) and F (0.8% glutamate).

### RESULTS AND DISCUSSION

The results indicated that giving of glutamate had a significant effect ( $P < 0.05$ ) on feed intake, body weight gain and feed conversion ratio. While the giving of glutamate did not have a significant effect ( $P > 0.05$ ) on the percentage of the carcass. The average FI, BWG, FCR and CP by giving glutamate was indicated in table 3.

#### Feed intake

It can be seen that group B was not significantly different ( $P > 0.05$ ) in the treatment of C, D and E (Table 3). Judging from the average value there was a tendency to increase feed consumption with an increase in the dose of glutamic acid which can be the same as the consumption of ration by providing commercial glutamate. Increased consumption of rations with the addition of glutamate caused by glutamate can improve the quality of the digestive tract which can increase the absorption of nutrients. According to Ebadiasl (2011) glutamate in small bowel function as the formation of intestinal mucous villi cells. Bartell and Batal (2007) reported that glutamate supplementation ~~can~~ could improve intestinal development with an increase in the relative weight of the duodenum and jejunum. Improving the development of the small intestine can increase the capacity of the digestive tract to accommodate the volume of the feed so that, many different rations can be consumed by broilers. According to Ibrahim (2008), larger intestines ~~can~~ could increase ration consumption because ~~there will be~~ a larger volume of food ~~that~~ can be accommodated, digested, and absorbed.

In addition, the increase in ration consumption in this study was due to a decrease in stress on broilers. Reducing stress by adding glutamate because it can synthesize amino acids which have a role in reducing stress levels. According to Young and Ajami (2000) states that glutamate is an amino acid builder protein which is a precursor for Gamma AminoButyric Acid (GABA). Glutamate will be converted into GABA if broilers were under stress. According to Wang et al. (2015) and Lener et al. (2017) GABA serves to reduce stress so that it can maintain appetite and increase the efficiency of the use of body nutrients in livestock. Thus giving glutamate to broiler chickens can reduce stress caused by increased-increasing feed consumption. Feed intake in this study was between 4478.75-4588.68 g lower than that obtained by Bezerra et al. (2015) which was 5303 g for 42 days of maintenance. The same was Similar study was reported-presented by Olubodun et al. (2015) that administration of 0.5% of glutamate to broilers resulted-and indicated in consumption of 4083 g feed.

### **Body weight gain**

The body weight gain was increased as an increasing dose of glutamate. Giving glutamate in the treatment of F can increase weight gain, but not significantly different ( $P > 0.05$ ) with treatment E. Body weight gain was influenced by feed consumption, the higher the consumption of food, the more weight gain was generated. According to Fadilah (2005) feed consumption has-had a positive effect on weight gain. Increased glutamate dose in this treatment caused by an increase in absorption of nutrients in the-small intestine. This is supported by the opinion of Shakeri et al. (2014) stated that administration of glutamate in broiler feed could improve small bowel development, intestinal villi length and nutrient absorption. Glutamate in the small intestine plays a role in the mechanism of intestinal mucosal defense by increasing the mucosal layer (Akiba et al., 2009). Increasing the inner surface and surface area of the small intestine will increase the digestion and absorption of feed extracts by the small intestine (Yao et al., 2006).

Giving glutamate can increase broiler weight because glutamate is a precursor for non-essential amino acids, so it will meet the needs of several other essential amino acids. According to Blachier et al. (2009) stated that glutamate is a precursor to other non-essential amino acids, such as arginine, glutamine, and proline. With the fulfillment of the need for non-essential amino acids will increase the growth of broilers. According to Ajinamoto (2007) and Maslami et al.

(2018) administration of glutamate in broiler chicken feed could increase the growth of connective tissue so that, it can increase broiler body weight gain.

The highest weight gain in this study was found in treatment F (2646.45 g / bird) which could match treatment A with commercial glutamate (2638.35 g / bird). The body weight gain in broiler chickens of this study was higher than that reported by Olubodun et al. (2005) at 42 days of age (2375 g / bird). Increased body weight gain by adding glutamate to broilers diet was reported by Porto et al. (2015). Furthermore, according to Bezerra et al. (2015) the addition of 1.76% glutamate could match the increase in broiler body weight gain without a reduction in crude protein ration. ~~Furthermore~~ ~~Moreover~~, according ~~to Bezerra et al. (2015)~~ ~~to this study~~ addition of 1.76% glutamate could increase body weight gain of broilers with a crude low protein ration.

#### Feed conversion ratio

The results indicated that treatment groups (D, E and F) gave the same low FCR, even matching treatment A with commercial glutamate. FCR was low by giving glutamate along with increased feed intake and body weight gain. A low FCR indicates an increase in feed efficiency (Razak et al., 2016). According to Usman (2009) and Zuidhof et al. (2014) the values of FCR ~~was~~ ~~were~~ influenced by the amount of feed intake and body weight gain. Another factor is the increase in feed conversion with increasing doses because glutamate can improve the nutritional quality of feed and the digestive system. It was reported by Andriyanto et al. (2015) **the nutritional quality of feed could influence broiler feed conversion.** Decreased FCR caused by administration of glutamate could improve the quality of the digestive system by increasing intestinal length thereby increasing absorption of feed nutrients (Olubodun et al., 2015). Increasing absorption of feed nutrients causes an increase in body weight so that it will increase the efficiency of feed use. ~~A low FCR indicates an increase in feed efficiency.~~

The lowest FCR in this study was 1.70% lower than Shakeri et al. (2014) stated that the FCR in broilers with administration of glutamate was 1.95%. Similar results were conveyed by Zulkifli et al. (2016) giving 1% glutamate to broiler could reduce FCR to 1.81%. Furthermore, according to Olubodun et al. (2015) ~~that~~ added ~~that~~ 0.5% of the mixture of glutamate and glutamine ~~to broiler chicken diet and could reported~~ reduce ~~the in~~ FCR of broiler chicken ~~to~~ (1.78%).

**Commented [32]:** It is not percent  
Author: I have revised it

**Commented [33]:** It is not percent  
Author: I have revised it

**Commented [34]:** It is not percent  
Author: I have revised it

**Commented [35]:** It is not percent  
Author: I have revised it



## Carcass percentage

The results of statistical analysis indicated that the administration of glutamate did not significantly affect ( $P > 0.05$ ) on the percentage of broiler carcass. The effect of giving glutamate on carcass percentage ~~can be seen~~ presented in table 3. The average carcass percentage of each treatment from treatment A to treatment F was 67.23, 68.76, 73.21, 70.70, 70.13, and 70.47% respectively. ~~The percentage of carcass was the ratio between carcass weight and live weight multiplied by 100%.~~ The percentage of the carcass in this study was at the range of 67.23-73.21% ~~which was and~~ supported by Resnawati (2004) ~~who stated the percentage of broiler carcass between 68-71.8%.~~

## CONCLUSION

Addition of glutamate up to 0.8% could improve the performance of broilers ~~for 35 days of maintenance.~~ However, administration of glutamate did not affect the percentage of the carcass. ~~Further researches should to be conduct to indicate the ideal glutamate dose in broilers diet.~~ ~~The percentage of carcass obtained in this study was in the normal.~~

## DECLARATIONS

### Acknowledgments

All authors are very grateful to the Minister of Research, Technology and Higher Education of the Republic of Indonesia for support and finance through PMDSU No: No: 1387 / E4. 2015 and support through the B-Class World Class Professor Program Scheme 123.57 / D2.3 / KP / 2018.

### Competing interests

The authors declare that they have no competing interests.

### Author's contribution

~~Vebera Maslami and Yetti Marlida conducted the research, and compile prepared data, performed statistical analysis and wrote the articles.~~ Mirnawati, Jamsari and Yuliaty Shafan Nur ~~checked and compiled confirmed the final form of article and revised the article.~~

### Consent to publish

**Commented [36]:** This methods and it is not results.  
Move it  
Author : I have revised it

**Commented [37]:** Is it related to Glutamate?  
Author: relation to glutamate is to support that the percentage of carcasses is still in normal conditions with glutamate administration

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Highlight

**Formatted:** Font: Times New Roman, 12 pt, Highlight

**Formatted:** Highlight

All the authors gave their informed consent prior to their inclusion in the study.

## REFERENCES

- Ajinomoto, AN- (2007). Influence of glutamic acid on broiler carcass quality. Poultry Research Report 19. DOI: <http://www.lysine.com/pdf/poultry/prr19.pdf>.
- Akiba Y, Watanabe C, Mizumori M and Kaunitz JD (2009). Luminal l-glutamate enhances duodenal mucosal defense mechanisms via multiple glutamate receptors in rats. American Journal of Physiology-Gastrointest Liver Physiology, 297: 781–791. DOI: 10.1152/ajpgi.90605.2008.
- Andriyanto, -Satyaningtjas AS, Yufiadri R, Wulandari R, Darwin VM and dan and Siburian SNA (2015). Performance and digestibility of broiler chicken feed fed with testosterone in multilevel doses. Journal - Acta Veterinaria Indonesiana, 3 (1): 29-37.
- Bartell S and Batal A (2007). The effect of supplemental glutamine on growth performance, development of the gastrointestinal tract, and humoral immune response of broilers. Poultry Science, 86:1940-1947. DOI: <https://10.1093/ps/86.9.1940>.
- Bezerra R-M, et al (2016). Effect of L-glutamate acid supplementation on performance and nitrogen balance of broilers fed low protein diets. Journal of animal physiology and animal nutrition, 100: 590–600. DOI: 10.1111/jpn.2405.
- Blachier F, Boutry C, Bos C and Tome D (2009). Metabolism and functions of L-glutamate in the epithelial cells of the small and large intestines. Am J Clin Nutr, 90(3):814-821 DOI: <https://10.3945/ajcn.2009.27462S>.
- Ebadias, G- (2011). Effects of supplemental glutamine and glutamate on growth performance, gastrointestinal development, jejunum morphology and Clostridium perfringens count in caecum of broilers. Thesis. Available at: <https://core.ac.uk/download/pdf/11987364.pdf>
- Ensminger ME, Scanes CG and Brant G (2004). Poultry Science. 4th Edition. Pearson Prentice Hall, New.
- Ibrahim S (2008). Hubungan ukuran-ukuran usus halus dengan berat badan broiler. Agripet, 8(2): 42- 46. DOI: <https://10.17969/agripet.v8i2.615>

**Commented [a8]:** Please write the reference according to the journal format. You can check one of journal's articles using this link: [http://iwpr.science-line.com/attachments/article/49/J%20World%20Poult%20Res%209\(2\)%20109-116,%202019.pdf](http://iwpr.science-line.com/attachments/article/49/J%20World%20Poult%20Res%209(2)%20109-116,%202019.pdf)

**Commented [a9]:** When the number of authors is more than 10, et al is used. Please just write the name of first ten author and then use et al

- Fadilah Y (2005). Panduan Mengelola Peternakan Ayam Broiler Komersial. Agromedia. Pustaka. Jakarta.
- Lener MS, Niciu MJ, Ballard ED, Park M, Park LT, Nugent A and Zarate CA (2017). Glutamate and GABA systems in the pathophysiology of major depression and antidepressant response to ketamine. *Biological Psychiatry*, 81(10): 886–897. DOI: <https://10.1016/j.biopsych.2016.05.005>
- Li P., Yin YL and Li D. 2007. Amino acids and immune function. *British Journal of Nutrition* 98: 237- 252. DOI: <https://10.1017/S000711450769936X>
- Madhupriya V, Shamsudeen P, Raj Manohar G, Senthilkumar S, Soundarapandiyam V and Moorthy (2018). Phyto feed additives in poultry nutrition – a review, *International Journal of Science, Environment and Technology*, Available at: <http://www.ijset.net/journal/2109.pdf>
- Maslami V, Marlida Y, Mirnawati, Jamsari and Nur YS (2018). Optimization of glutamate production from *Lactobacillus plantarum* originating from Minangkabau fermented food as a feed supplement for broiler. *Pakistan Journal of Nutrition*, 17(7): 336-343. DOI: <https://10.2923/pjn.2018.336.343>.
- Maslami V, Marlida Y, Mirnawati, Jamsari, Nur YS, Adzitey F and Huda N (2018). A review on potential of glutamate producing lactic acid bacteria of west Sumatera's fermented food origin, As feed additive for broiler chicken. *Journal of World's Poultry Research*, 8(4):120-126. PII: S2322455X1800017-8
- Nampoothiri K, Madhavan and Ashok P (1996). Urease Activity in a Glutamate Producing *Brevibacterium* sp. *Procrss Biochemistry*, 31: 471-475. DOI: [https://10.1016/S0032-9592\(95\)00090-9](https://10.1016/S0032-9592(95)00090-9).
- Olubodun JO, Zulkifli I, Farjam AS, Hair-Bejo M and Kasim A (2015). Glutamine and glutamic acid supplementation enhances performance of broiler chickens under the hot and humid tropical condition. *Italian Journal of Animal Science*, 14 (3263): 25-29. DOI: <https://10.4081/ijas.2015.3263>.
- Porto ML, Givisiez PEN, Saraiva EP, Costa FGP, Moreira Filho ALB, Andrade MFS, Brandão PA and Guerra RR (2015). Glutamic Acid Improves Body Weight Gain and Intestinal Morphology of Broiler Chickens Submitted to Heat Stress. *Brazilian Journal of Poultry Science*, 17 (3): 355-362. DOI: <https://10.1590/1516-635x1703355-362>.

Razak AD Kiramang K and Hidayat MN. 2016. Increase in body weight, feed consumption and feed conversion of broilers fed with betel leaf flour (*Piper Betle* Linn) as feed additives. 3: 135-147. Available at: <http://journal.uin-alauddin.ac.id/index.php/jiip/article/view/3924> ISSN: 2442-2541

Reeds PJ, Burrin DG, Stoll B and Jahoor F (2000). Intestinal glutamate metabolism. *Journal of Nutrition*, 130(4): 978-82. DOI: [https://10.1093/jn/130.4.978S](https://doi.org/10.1093/jn/130.4.978S).

Resnawati (2004). Bobot potongan karkas dan lemak abdomen ayam ras pedaging yang diberi ransum mengandung tepung cacing tanah. Available at: [http://peternakan.litbang.deptan.go.id/user/pros\\_04-75.pdf](http://peternakan.litbang.deptan.go.id/user/pros_04-75.pdf). (Accessed: 22 September 2018).

Ruegg PL (2013). Antimicrobial residues and resistance: Understanding and managing drug usage on dairy farms. University of Wisconsin, Department of Dairy Science, Madison.

Saputra WY, Mahfudz LD dan Suthama N (2013). Pemberian pakan single step down dengan penambahan asam sitrat sebagai acidifier terhadap performa pertumbuhan broiler. *Animal Agriculture Journal*, 2(3): 61-72. Available at: <http://ejournal-s1.undip.ac.id/index.php/aaaj>

Shakeri M, Zulkifli I, Soleimani AF, Reilly ELO, Eckersall PD, Anna AA, Kumara S and Abdullah FFJ (2014). Response to dietary supplementation of L-glutamine and L-glutamate in broiler chickens reared at different stocking densities under hot, humid tropical conditions. *Poultry Science*, 93:1-9. DOI: [https://10.3382/ps.2014-03910](https://doi.org/10.3382/ps.2014-03910).

Singh S, Sanjay S, Neelam T, Nitesh K dan Ritu P (2014). Antibiotic residues: a global challenge. *An International Journal of Pharmaceutical Science. Pharma Science Monitor*, 5 (3):184-197. DOI: [https://10.5455/vetworld.2008.375-377](https://doi.org/10.5455/vetworld.2008.375-377)

Steel RGD and Torrie JH (1991). Prinsip dan Prosedur Statistika. Translated by Bambang Sumantri. PT. Gramedia Pustaka Utama. Jakarta

Usman (2009). Pertumbuhan ayam buras periode grower melalui pemberian tepung biji buah merah (*Pandanus conoideus* LAMK) sebagai pakan alternatif. Prosiding Seminar Nasional Teknologi Peternakan dan Veteriner. Balai Pengkajian Teknologi Pertanian Papua.

- Wang DM, Chacher, Liu HY, Wang JK, Lin J and Liu JX (2015). Effects of  $\gamma$ -aminobutyric acid on feed intake, growth performance and expression of related genes in growing lambs. *Animal*, 9(3): 445-448. DOI: <https://10.1017/S1751731114002651>.
- Yao Y, Xiaoyan T, Haibo X, Jinchun K, Ming X and Xiaobing W (2006). Effect of choice feeding on performance gastrointestinal development and feed utilization of broilers. *Asian-Australasian Journal of Animal Sciences*, 19:91-96. DOI: <https://10.5713/ajas.2006.91>.
- Young VR and Ajami AM (2000). Glutamate: an amino acid of particular distinction. *Journal of Nutrition*, 130: 892–900. DOI: 10.1093/jn/130.4.892S.
- Zuidhof MJ, Scheider BL, Carney VL, Korver DR and Robinson FE (2014). Growth, efficiency and yield of commercial broilers from 1957, 1978 and 2005. *Poultry Science*, 93(12): 2970-2982. DOI: <https://10.3382/ps.2014-04291>.
- Zulkifli M, Shakeri M and Soleimani AF (2016). Dietary supplementation of L-glutamine and L-glutamate in broiler chicks subjected to delayed placement. *Poultry Science*, 1: 1-7. DOI: <https://10.3382/ps/pew267>.

**Table 1.** Nutrient content of feed ingredients for broiler chickens

Feed ingredients	Nutritional content ( % ) and energy metabolism (Kkal/kg)					
	Crude Protein	Crude Fiber	Crude Fat	Ca	P	ME
Corn	9.55	3.8	2.18	0.38	0.33	3300
Bran	10.6	10.84	4.09	0.7	0.09	1592
Fish flour	41	2.8	1.52	5.55	2.6	2580
Commercial Feed	23	1.88	5.87	0.29	0.15	3200
Soybean Meal	40.16	3.58	1.37	0.63	0.32	2240
Coconut oil	0	0	100	0	0	8600
Mineral Premix	0	0	0	5.38	1.14	0

ME: Energy metabolism, Ca: Calcium, P: Phosphor.

**Table 2.** The composition of rations of broiler chickens

Feed ingredients	Percentage of ration (%) and nutrient content (%)
Corn	40
Bran	7
Fish flour	17
Commercial Feed	20
Soybean Meal	14
Coconut oil	1.5
Mineral Premix	0.5

**Commented [310]:** Explain it under the table  
What is it

**Commented [311]:** Explain it under the table  
What is it

Total	100
Crude Protein	21.75
Crude Fiber	3.86
Crude Fat	4.12
Ca	1.50
P	0.83
ME (Kcal/kg)	2952,64

Ca: Calcium, P: Phosphor, ME: Energy metabolism.

**Table 3.** Effect of dietary inclusion of glutamate on broiler chicken performance for 35 days

Treatment	FI (g)	BWG (g)	FCR (%)	CP (%)
A (0.4% Commercial Glutamate)	4591.13 <sup>a</sup> ± 63.67	2688.5 <sup>a</sup> ±24.73	1.71 <sup>b</sup> ±0.03	67.23 ± 3.52
B (0 % Glutamate)	4429.05 <sup>b</sup> ± 77.25	2391.50 <sup>c</sup> ±28.25	1.85 <sup>a</sup> ±0.05	68.76 ± 1.80
C (0.2 % Glutamate)	4478.75 <sup>ab</sup> ± 84.25	2450.63 <sup>c</sup> ±54.96	1.83 <sup>a</sup> ±0.05	73.21 ± 3.62
D (0.4 % Glutamate)	4530.28 <sup>ab</sup> ± 76.08	2573.60 <sup>b</sup> ±63.79	1.76 <sup>b</sup> ±0.07	70.70 ± 3.44
E (0.6 % Glutamate)	4503.70 <sup>ab</sup> ± 103.94	2634.15 <sup>ab</sup> ±31.66	1.71 <sup>b</sup> ± 0.04	70.13 ± 3.04
F (0.8% Glutamate)	4588.68 <sup>a</sup> ± 26.24	2693.95 <sup>a</sup> ±74.06	1.70 <sup>b</sup> ±0.05	70.47 ± 4.39

**Commented [312]:** Percent?  
Author: yes

**Commented [313]:** You should indicate the superscript letter for these data in this column

**Commented [314]:** You should indicate the superscript letter for these data in this column

FI: Feed intake, BWG: body weight gain, FCR: feed conversion ratio, CP: Carcass percentage.

**Original article**

**Effect of Glutamate Supplementation as a Feed Additive on Performance of Broiler Chickens**

Vebera Maslami<sup>1</sup>, Mirnawati<sup>2</sup>, Jamsari<sup>3</sup>, Yuliaty Shafan Nur<sup>2</sup> and Yetti Marlida<sup>2\*</sup>

<sup>1</sup>*Department of Agricultural Science, Faculty of Agriculture, Andalas University, 25163, Indonesia*

<sup>2</sup>*Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, Andalas University, 25163, Indonesia*

<sup>3</sup>*Department of Crop Science, Faculty of Agriculture, University Andalas, 25163, Indonesia*

\*Corresponding author's E-mail: [yettimarlida@ansci.unand.ac.id](mailto:yettimarlida@ansci.unand.ac.id)

**ABSTRACT**

Feed additives are ingredients that are added to the ration as growth promoters and enhancement of the immune system. Glutamate is a feed additive that improves performance by improving the quality of the small intestine and enhancing the immune system. The purpose of this study was to know the effect of including glutamate as a feed additive in improving broiler performance. The material used in this study was broiler strain MB 202 from PT Charoen Phokphand Indonesia as many as 240 birds. The design used was a Completely Randomized Design (CRD) trial design, with six treatments and four replications, so that there were 24 cage plots as experimental units. Each experimental unit consisted of 10 chickens. The Glutamate doses in groups were, A (0.4% commercial glutamate); B (0% glutamate); C (0.2% glutamate); D (0.4% glutamate); E (0.6% glutamate); F (0.8% glutamate). The results indicated that glutamate up to 0.8% had significant effects on feed intake body weight gain and feed conversion ratio (1.70%), but it did not affect the percentage of carcass. It is concluded that including the 0.8% glutamate in broiler diets can improve broiler performance with 35 days of maintenance.

**Key words:** Feed additive, Glutamate, Growth promoters, Immune system, Performance



## INTRODUCTION

The broiler is one of the meat-producing livestock that is quite potent in meeting the people's needs for animal protein needs. This is because broiler meat is relatively cheap and easy to obtain compared to other animal proteins. According to [Ensminger et al. \(2004\)](#) broilers have a fast and efficient growth in converting feed into meat. In addition, broilers also have weaknesses which tend to be susceptible to disease attacks which results in a decrease in broiler performance.

Supplementation of additional feed is one solution to prevent disease attacks by increasing immunity and growth promoters. According to [Madhupriya et al. \(2018\)](#) Feed additives are ingredients and add to rations to improve animal immunity and performance. Antibiotic Growth Promoter (AGP) is an antibiotic feed additive added to broiler feed. However, the use of AGP in broiler feed is prohibited because residues in broiler meat are harmful to public health. Residual hazards could reduce human resistance to consumption of certain types of antibiotics and other hazards ([Ruegg, 2013](#); [Singh et al., 2014](#)). This had led to the enactment of the World Health Organization's regulation regarding the prohibition on the use of antibiotics in animal feed, hence the use of antibiotics as feed additives in broiler feed. So it is necessary to look for additional feed ingredients that are safer for livestock and the community. One safe and harmless feed additive is glutamate.

Glutamate is a feed additive that can improve growth promotes and enhances the broiler immune system. Glutamate functions was known as a constituent of proteins, a substrate in the synthesis of amino acids, as a precursor to several non-essential amino acids and helped the metabolism ([Young and Ajami, 2000](#)). According to [Newholms et al. \(2003\)](#); [Reeds et al. \(2000\)](#) glutamate could improve intestinal conditions by increasing intestinal villi length and increasing maintenance of intestinal integrity. In addition, according to [Li et al. \(2007\)](#) glutamate regulated Inductible Nitric Oxide Synthase (iNOS) in certain tissues. The expression of iNOS was considered as a fundamental mechanism in the protection of parasites, bacteria, fungi, malignant cells, intracellular protozoa and viruses in different animal species, including mammals and birds. The present study aimed to know the effect of giving glutamate as a feed additive in broilers.

## **MATERIALS AND METHODS**

### **Place of study**

This research was conducted on May-September 2018. Maintenance of broilers was conducted at the Poultry Division Field Laboratory, Faculty of Animal Husbandry, Andalas university, Indonesia.

### **Materials**

A total of 240 one-day old male broiler chicks (MB 202 from) were purchased from PT Charoen Pokphand, Indonesia. The chickens were placed randomly into 24 cage plots (open cages), each measuring 1×1 m<sup>2</sup>. Each plot was filled with 10 chickens, equipped with food and drinking water. This broiler cage was also equipped with 1 bulb (35 Watt). The ration was prepared by itself from feed ingredients consisting of corn, commercial feed (CP511 PT, Charoen Pokphand Indonesia), fish meal, soy flour, bran, palm oil and mineral premix. Nutrient content and feed composition was indicated in table 1 and table 2.

### **Method**

Making glutamate was done by fermentation with *Lactobacillus plantarum*. The nutritional composition of fermentation media used in this study was KH<sub>2</sub>PO<sub>4</sub>, MgSO<sub>4</sub> · 7H<sub>2</sub>O, FeSO<sub>4</sub> · 7H<sub>2</sub>O, MnSO<sub>4</sub> · 4-5H<sub>2</sub>O, 9% sugar cane, 5 μ / L biotin, 90% tofu water, 10% distilled water, water starter and starter 9%. The duration of fermentation in this study was 36 hours at 36°C. After fermentation the media were centrifuged for 20 minutes and 10,000 rpm at a temperature of -4°C. After that, the supernatant was concentrated using an evaporation technique in an oven with a temperature of 40°C for 48 hours. The shrinking material was then calculated for the acid content using glutamate HPLC (Maslami et al., 2018).

### **Maintenance of broilers**

Maintenance was carried out for five weeks. Feed and drinking water were available ad libitum. Glutamate treatment was given to chickens aged two weeks to six weeks. Glutamate was added to drinking water. The body weight was carried out at the beginning of the study and every following week, and also at the end of the study. Weighing the rest of the feed was done every

weekend. Calculations were carried out for Feed Intake (FI), Body Weight Gain (BWG), Feed Conversion Ratio (FCR) and Carcass Percentage (CP).

### **Experiment and data analysis**

This study used a completely randomized design. Each treatment was repeated four times. Each treatment and each replication consisted of 10 birds. The observed variables were just performance traits. The variable performance included average feed intake, body weight gain, feed conversion ratio and carcass percentage. The data obtained were analyzed using analysis of variance (Analysis of Variance / ANOVA) and if there were differences between treatments were tested further using the DMRT (Duncan Multiple Range Test) tests at a level of 5% (Steel and Torrie, 1991). The dose of glutamate in drinking water is A (0.4% commercial glutamate), B (0.0% glutamate), C (0.2% glutamate), D (0.4% glutamate), E (0.6% glutamate) and F (0.8% glutamate).

### **RESULTS AND DISCUSSION**

The results indicated that giving of glutamate had a significant effect ( $P < 0.05$ ) on feed intake, body weight gain and feed conversion ratio. While the giving of glutamate did not have a significant effect ( $P > 0.05$ ) on the percentage of the carcass. The average FI, BWG, FCR and CP by giving glutamate was indicated in table 3.

#### **Feed intake**

It can be seen that group B was not significantly different ( $P > 0.05$ ) in the treatment of C, D and E (Table 3). Judging from the average value there was a tendency to increase feed consumption with an increase in the dose of glutamic acid which can be the same as the consumption of ration by providing commercial glutamate. Increased consumption of rations with the addition of glutamate caused by glutamate can improve the quality of the digestive tract which can increase the absorption of nutrients. According to Ebadiasl (2011) glutamate in small bowel function as the formation of intestinal mucous villi cells. Bartell and Batal (2007) reported that glutamate supplementation could improve intestinal development with an increase in the relative weight of the duodenum and jejunum. Improving the development of the small intestine can increase the capacity of the digestive tract to accommodate the volume of the feed so that, many different rations can be consumed by broilers. According to Ibrahim (2008), larger

intestines could increase ration consumption because a larger volume of food can be accommodated, digested, and absorbed.

In addition, the increase in ration consumption in this study was due to a decrease in stress on broilers. Reducing stress by adding glutamate because it can synthesize amino acids which have a role in reducing stress levels. According to Young and Ajami (2000) stated that glutamate is an amino acid builder protein which is a precursor for Gamma AminoButyric Acid (GABA). Glutamate will be converted into GABA if broilers were under stress. According to Wang et al. (2015) and Lener et al. (2017) GABA served to reduce stress so that it can maintain appetite and increase the efficiency of the use of body nutrients in livestock. Thus giving glutamate to broiler chickens can reduce stress by increasing feed consumption. Feed intake in this study was between 4478.75-4588.68 g lower than that obtained by Bezerra et al. (2015) which was 5303 g for 42 days of maintenance. Similar study was presented by Olubodun et al. (2015) that administrated 0.5% of glutamate to broilers and indicated consumption of 4083 g feed.

### **Body weight gain**

The body weight gain was increased as an increasing dose of glutamate. Giving glutamate in the treatment of F can increase weight gain, but not significantly different ( $P > 0.05$ ) with treatment E. Body weight gain was influenced by feed consumption, the higher the consumption of food, the more weight gain was generated. According to Fadilah (2005) feed consumption had a positive effect on weight gain. Increased glutamate dose in this treatment caused by an increase in absorption of nutrients in small intestine. This is supported by the opinion of Shakeri et al. (2014) stated that administration of glutamate in broiler feed could improve small bowel development, intestinal villi length and nutrient absorption. Glutamate in the small intestine plays a role in the mechanism of intestinal mucosal defense by increasing the mucosal layer (Akiba et al., 2009). Increasing the inner surface and surface area of the small intestine will increase the digestion and absorption of feed extracts by the small intestine (Yao et al., 2006).

Giving glutamate can increase broiler weight because glutamate is a precursor for non-essential amino acids, so it will meet the needs of several other essential amino acids. According to Blachier et al. (2009) stated that glutamate is a precursor to other non-essential amino acids,

such as arginine, glutamine, and proline. With the fulfillment of the need for non-essential amino acids will increase the growth of broilers. According to Ajinamoto (2007) and Maslami et al. (2018) administration of glutamate in broiler chicken feed could increase the growth of connective tissue so that, it can increase broiler body weight gain.

The highest weight gain in this study was found in treatment F (2646.45 g / bird) which could match treatment A with commercial glutamate (2638.35 g / bird). The body weight gain in broiler chickens of this study was higher than that reported by Olubodun et al. (2005) at 42 days of age (2375 g / bird). Increased body weight gain by adding glutamate to broilers diet was reported by Porto et al. (2015). Furthermore, according to Bezerra et al. (2015) the addition of 1.76% glutamate could match the increase in broiler body weight gain without a reduction in crude protein ration. Moreover, according to this study addition of 1.76% glutamate could increase body weight gain of broilers with a crude low protein ration.

#### **Feed conversion ratio**

The results indicated that treatment groups (D, E and F) gave the same low FCR, even matching treatment A with commercial glutamate. FCR was low by giving glutamate along with increased feed intake and body weight gain. A low FCR indicates an increase in feed efficiency (Razak et al., 2016). According to Usman (2009) and Zuidhof et al. (2014) the values of FCR were influenced by the amount of feed intake and body weight gain. Another factor is the increase in feed conversion with increasing doses because glutamate can improve the nutritional quality of feed and the digestive system. It was reported by Andriyanto et al. (2015) the nutritional quality of feed could influence broiler feed conversion. Decreased FCR caused by administration of glutamate could improve the quality of the digestive system by increasing intestinal length thereby increasing absorption of feed nutrients (Olubodun et al., 2015). Increasing absorption of feed nutrients causes an increase in body weight so that it will increase the efficiency of feed use.

The lowest FCR in this study was 1.70 lower than Shakeri et al. (2014) stated that the FCR in broilers with administration of glutamate was 1.95 Similar results were conveyed by Zulkifli et al. (2016) giving 1% glutamate to broiler could reduce FCR to 1.81 Furthermore, according to Olubodun et al. (2015) that added 0.5% of the mixture of glutamate and glutamine to broiler chicken diet and reported reduce in FCR (1.78).

### **Carcass percentage**

The results of statistical analysis indicated that the administration of glutamate did not significantly affect ( $P > 0.05$ ) on the percentage of broiler carcass. The effect of giving glutamate on carcass percentage presented in table 3. The average carcass percentage of each treatment from treatment A to treatment F was 67.23, 68.76, 73.21, 70.70, 70.13, and 70.47% respectively. The percentage of the carcass in this study was at the range of 67.23-73.21% which was supported by Resnawati (2004) who studied in dietary addition of glutamate in broiler chickens and stated the percentage of carcass between 68-71.8%. Glutamate is a building block of proteins that can increase protein synthesis in muscle tissue. Formation of protein in muscle tissue will affect the percentage of carcasses. According to Reeds et al. (2000), glutamate is one of the building blocks of amino acids from proteins that make up the body's protein.

### **CONCLUSION**

Addition of glutamate up to 0.8% could improve the performance of broilers for 35 days of maintenance. However, administration of glutamate did not affect the percentage of the carcass. Further researches should to be conduct to indicate the ideal glutamate dose in broilers diet.

### **DECLARATIONS**

#### **Acknowledgments**

All authors are very grateful to the Minister of Research, Technology and Higher Education of the Republic of Indonesia for support and finance through PMDSU No: No: 1387 / E4. 2015 and support through the B-Class World Class Professor Program Scheme 123.57 / D2.3 / KP / 2018.

#### **Competing interests**

The authors declare that they have no competing interests.

#### **Author's contribution**

Vebera Maslami and Yetti Marlida conducted the research, prepared data and performed statistical analysis. Mirnawati, Jamsari and Yuliaty Shafan Nur wrote the article. All authors checked and confirmed the final form of article.

### Consent to publish

All authors gave their informed consent prior to their inclusion in the study.

### REFERENCES

- Ajinomoto AN (2007). Influence of glutamic acid on broiler carcass quality. Poultry Research Report 19. Available at: <http://www.lysine.com/pdf/poultry/prr19.pdf>.
- Akiba Y, Watanabe C, Mizumori M and Kaunitz JD (2009). Luminal l-glutamate enhances duodenal mucosal defense mechanisms via multiple glutamate receptors in rats. *American Journal of Physiology-Gastrointest Liver Physiology*, 297: 781–791. DOI: <https://10.1152/ajpgi.90605.2008>.
- Andriyanto, Satyaningtjas AS, Yufiadri R, Wulandari R, Darwin VM and and Siburian SNA (2015). Performance and digestibility of broiler chicken feed fed with testosterone in multilevel doses. *Journal Acta Veterinaria Indonesiana*, 3 (1): 29-37.
- Bartell S and Batal A (2007). The effect of supplemental glutamine on growth performance, development of the gastrointestinal tract, and humoral immune response of broilers. *Poultry Science*, 86:1940-1947. DOI: <https://10.1093/ps/86.9.1940>.
- Bezerra RM, Costa GP, Givisiez PEN, Freitas ER, Goulart CC, Santos RA, Souza JG, Brandao PA, Lima MR, Melo ML, etal. (2016). Effect of L-glutamate acid supplementation on performance and nitrogen balance of broilers fed low protein diets. *Journal of animal physiology and animal nutrition*, 100: 590–600. DOI: <https://10.1111/jpn.2405>.
- Blachier F, Boutry C, Bos C and Tome D (2009). Metabolism and functions of L-glutamate in the epithelial cells of the small and large intestines. *Am J Clin Nutr*, 90(3):814-821 DOI: <https://10.3945/ajcn.2009.27462S>.
- Ebadiasl G (2011). Effects of supplemental glutamine and glutamate on growth performance, gastrointestinal development, jejunum morphology and *Clostridium perfringens* count in caecum of broilers. Thesis. Available at: <https://core.ac.uk/download/pdf/11987364.pdf>
- Ensminger ME, Scanes CG and Brant G (2004). *Poultry Science*. 4th Edition. Pearson Prentice Hall, New.

- Ibrahim S (2008). Hubungan ukuran-ukuran usus halus dengan berat badan broiler. *Agripet*, 8(2): 42- 46. DOI: <https://10.17969/agripet.v8i2.615>
- Fadilah Y (2005). *Panduan Mengelola Peternakan Ayam Broiler Komersial*. Agromedia. Pustaka. Jakarta.
- Lener MS, Niciu MJ, Ballard ED, Park M, Park LT, Nugent A and Zarate CA (2017). Glutamate and GABA systems in the pathophysiology of major depression and antidepressant response to ketamine. *Biological Psychiatry*, 81(10): 886–897. DOI: <https://10.1016/j.biopsych.2016.05.005>
- Li P., Yin YL and Li D. 2007. Amino acids and immune function. *British Journal of Nutrition* 98: 237- 252. DOI: <https://10.1017/S000711450769936X>
- Madhupriya V, Shamsudeen P, Raj Manohar G, Senthilkumar S, Soundarapandiyan V and Moorthy (2018). Phyto feed additives in poultry nutrition – a review, *International Journal of Science, Environment and Technology*, Available at: <http://www.ijset.net/journal/2109.pdf>
- Maslami V, Marlida Y, Mirnawati, Jamsari and Nur YS (2018). Optimization of glutamate production from *Lactobacillus plantarum* originating from Minangkabau fermented food as a feed supplement for broiler. *Pakistan Journal of Nutrition*, 17(7): 336-343. DOI: <https://10.2923/pjn.2018.336.343>.
- Maslami V, Marlida Y, Mirnawati, Jamsari, Nur YS, Adzitey F and Huda N (2018). A review on potential of glutamate producing lactic acid bacteria of west Sumatera's fermented food origin, As feed additive for broiler chicken. *Journal of World's Poultry Research*, 8(4):120-126. PII: S2322455X1800017-8
- Nampoothiri K, Madhavan and Ashok P (1996). Urease Activity in a Glutamate Producing *Brevibacterium* sp. *Procrss Biochemistry*, 31: 471-475. DOI: [https://10.1016/S0032-9592\(95\)00090-9](https://10.1016/S0032-9592(95)00090-9).
- Olubodun JO, Zulkifli I, Farjam AS, Hair-Bejo M and Kasim A (2015). Glutamine and glutamic acid supplementation enhances performance of broiler chickens under the hot and humid tropical condition. *Italian Journal of Animal Science*, 14 (3263): 25-29. DOI: <https://10.4081/ijas.2015.3263>.
- Porto ML, Givisiez PEN, Saraiva EP, Costa FGP, Moreira Filho ALB, Andrade MFS, Brandão PA and Guerra RR (2015). Glutamic Acid Improves Body Weight Gain and Intestinal Morphology of Broiler Chickens Submitted to Heat Stress. *Brazilian Journal of Poultry Science*, 17 (3): 355-362. DOI: <https://10.1590/1516-635x1703355-362>.



- Razak AD Kiramang K and Hidayat MN (2016). Increase in body weight, feed consumption and feed conversion of broilers fed with betel leaf flour (*Piper Betle* Linn) as feed additives, 3: 135-147. Available at: <http://journal.uin-alauddin.ac.id/index.php/jiip/article/view/3924>
- Reeds PJ, Burrin DG, Stoll B and Jahoor F (2000). Intestinal glutamate metabolism. *Journal of Nutrition*, 130(4): 978-82. DOI: <https://10.1093/jn/130.4.978S>.
- Resnawati H (2004). Bobot potongan karkas dan lemak abdomen ayam ras pedaging yang diberi ransum mengandung tepung cacing tanah. Available at: [http://peternakan.litbang.deptan.go.id/user/pros\\_04-75.pdf](http://peternakan.litbang.deptan.go.id/user/pros_04-75.pdf). (Accessed: 22 September 2018).
- Ruegg PL (2013). Antimicrobial residues and resistance: Understanding and managing drug usage on dairy farms. University of Wisconsin, Department of Dairy Science, Madison.
- Saputra WY, Mahfudz LD dan Suthama N (2013). Pemberian pakan single step down dengan penambahan asam sitrat sebagai acidifier terhadap performa pertumbuhan broiler. *Animal Agriculture Journal*, 2(3): 61-72. Available at: <http://ejournal-s1.undip.ac.id/index.php/aaaj>
- Shakeri M, Zulkifli I, Soleimani AF, Reilly ELO, Eckersall PD, Anna AA, Kumara S and Abdullah FFJ (2014). Response to dietary supplementation of L-glutamine and L-glutamate in broiler chickens reared at different stocking densities under hot, humid tropical conditions. *Poultry Science*, 93:1-9. DOI: <https://10.3382/ps.2014-03910>.
- Singh S, Sanjay S, Neelam T, Nitesh K dan Ritu P (2014). Antibiotic residues: a global challenge. *An International Journal of Pharmaceutical Science. Pharma Science Monitor*, 5 (3):184-197. DOI: <https://10.5455/vetworld.2008.375-377>
- Steel RGD and Torrie JH (1991). *Prinsip dan Prosedur Statistika*. Translated by Bambang Sumantri. PT. Gramedia Pustaka Utama. Jakarta
- Usman (2009). Pertumbuhan ayam buras periode grower melalui pemberian tepung biji buah merah (*Pandanus conoideus* LAMK) sebagai pakan alternatif. *Prosiding Seminar Nasional Teknologi Peternakan dan Veteriner*. Balai Pengkajian Teknologi Pertanian Papua.

- Wang DM, Chacher, Liu HY, Wang JK, Lin J and Liu JX (2015). Effects of  $\gamma$ -aminobutyric acid on feed intake, growth performance and expression of related genes in growing lambs. *Animal*, 9(3): 445-448. DOI: <https://10.1017/S1751731114002651>.
- Yao Y, Xiaoyan T, Haibo X, Jinchun K, Ming X and Xiaobing W (2006). Effect of choice feeding on performance gastrointestinal development and feed utilization of broilers. *Asian-Australasian Journal of Animal Sciences*, 19:91-96. DOI: <https://10.5713/ajas.2006.91>.
- Young VR and Ajami AM (2000). Glutamate: an amino acid of particular distinction. *Journal of Nutrition*, 130: 892–900. DOI: <https://10.1093/jn/130.4.892S>.
- Zuidhof MJ, Scheider BL, Carney VL, Korver DR and Robinson FE (2014). Growth, efficiency and yield of commercial broilers from 1957, 1978 and 2005. *Poultry Science*, 93(12): 2970-2982. DOI: <https://10.3382/ps.2014-04291>.
- Zulkifli M, Shakeri M and Soleimani AF (2016). Dietary supplementation of L-glutamine and L-glutamate in broiler chicks subjected to delayed placement. *Poultry Science*, 1: 1-7. DOI: <https://10.3382/ps/pew267>.

**Table 1.** Nutrient content of feed ingredients for broiler chickens

Feed ingredients	Nutritional content ( % ) and energy metabolism (Kkal/kg)					
	Crude Protein	Crude Fiber	Crude Fat	Ca	P	ME
Corn	9.55	3.8	2.18	0.38	0.33	3300
Bran	10.6	10.84	4.09	0.7	0.09	1592
Fish flour	41	2.8	1.52	5.55	2.6	2580
Commercial Feed*	23	1.88	5.87	0.29	0.15	3200
Soybean Meal	40.16	3.58	1.37	0.63	0.32	2240
Coconut oil	0	0	100	0	0	8600
Mineral Premix	0	0	0	5.38	1.14	0

ME: Energy metabolism, Ca: Calcium, P: Phosphor

\*Commercial Feed: CP511 PT, Charoen Pokphand Indonesia

**Table 2.** The composition of ration in broiler chickens

Feed ingredients	Percentage of ration ( % ) and nutrient content ( % )
Corn	40
Bran	7
Fish flour	17
Commercial Feed*	20
Soybean Meal	14
Coconut oil	1.5
Mineral Premix	0.5
Total	100
Crude Protein	21.75
Crude Fiber	3.86
Crude Fat	4.12
Ca	1.50
P	0.83
ME (Kcal/kg)	2952,64

Ca: Calcium, P: Phosphor, ME: Energy metabolism,

\*Commercial feed: CP511 PT, Charoen Pokphand, Indonesia

Mineral Premix: mineral mixture PT. Medion Indonesia (Ca 5.38% and 1.44%)

**Table 3.** Effect of dietary inclusion of glutamate on broiler chicken performance for 35 days

Treatment	FI (g)	BWG (g)	FCR (%)	CP (%)
A (0.4% Commercial Glutamate)	4591.13 <sup>a</sup> ± 63.67	2688.5 <sup>a</sup> ±24.73	1.71 <sup>b</sup> ±0.03	67.23 ± 3.52
B (0 % Glutamate)	4429.05 <sup>b</sup> ± 77.25	2391.50 <sup>c</sup> ±28.25	1.85 <sup>a</sup> ±0.05	68.76 ±1.80
C (0.2 % Glutamate)	4478.75 <sup>ab</sup> ± 84.25	2450.63 <sup>c</sup> ±54.96	1.83 <sup>a</sup> ±0.05	73.21 ±3.62
D (0.4 % Glutamate)	4530.28 <sup>ab</sup> ± 76.08	2573.60 <sup>b</sup> ±63.79	1.76 <sup>b</sup> ±0.07	70.70 ± 3.44
E (0.6 % Glutamate)	4503.70 <sup>ab</sup> ± 103.94	2634.15 <sup>ab</sup> ±31.66	1.71 <sup>b</sup> ±0.04	70.13 ± 3.04
F (0.8% Glutamate)	4588.68 <sup>a</sup> ± 26.24	2693.95 <sup>a</sup> ±74.06	1.70 <sup>b</sup> ±0.05	70.47 ± 4.39

FI: Feed intake, BWG: body weight gain, FCR: feed conversion ratio, CP: Carcass percentage.

**Commented [31]:** You must indicate the minerals ingredients under the table

Which minerals and how much?

Example article under the table 1 of this article: (Premix)

[https://www.researchgate.net/profile/Daryoush-Babazadeh/publication/230728043\\_Effects\\_of\\_probiotic\\_prebiotic\\_and\\_synbiotic\\_intake\\_on\\_blood\\_enzymes\\_and\\_performance\\_of\\_Japanese\\_quails\\_Coturnix\\_Japonica/links/0fcfd5038ac5bd7f1a000000.pdf](https://www.researchgate.net/profile/Daryoush-Babazadeh/publication/230728043_Effects_of_probiotic_prebiotic_and_synbiotic_intake_on_blood_enzymes_and_performance_of_Japanese_quails_Coturnix_Japonica/links/0fcfd5038ac5bd7f1a000000.pdf)

or table 1 of this this article (Premix):

[https://www.researchgate.net/profile/Daryoush-Babazadeh/publication/236659655\\_Effects\\_of\\_saccharomyces\\_cerevisiae\\_thepax\\_and\\_their\\_combination\\_on\\_blood\\_enzymes\\_and\\_performance\\_of\\_Japanese\\_quails\\_Coturnix\\_japonica/links/00463518bea51ceaf000000/Effects-of-saccharomyces-cerevisiae-thepax-and-their-combination-on-blood-enzymes-and-performance-of-Japanese-quails-Coturnix-japonica.pdf](https://www.researchgate.net/profile/Daryoush-Babazadeh/publication/236659655_Effects_of_saccharomyces_cerevisiae_thepax_and_their_combination_on_blood_enzymes_and_performance_of_Japanese_quails_Coturnix_japonica/links/00463518bea51ceaf000000/Effects-of-saccharomyces-cerevisiae-thepax-and-their-combination-on-blood-enzymes-and-performance-of-Japanese-quails-Coturnix-japonica.pdf)

Move to | Categorise | Snooze | ...

Review Process, JWPR 1099 3

**SiencelinePublication**  
<http://www.science-line.com/index/>

**The Journal of World's Poultry Research**  
<http://www.jwpr-science-line.com/index/>

**Dear Respected Author**

It's my great pleasure to inform you that your paper encoded "JWPR-1099" has been reviewed by our selected reviewers and editors and accepted after **moderate revision** for fast publishing in [The Journal of World's Poultry Research \(JWPR\)](#). The following attachments are declaration form, the invoice and the images of the review results. Kindly pay **the publication fee (\$100)**, required to processing the paper **within 7 days** and then send the information of your payment (the receipt) to [editorjwpr@gmail.com](mailto:editorjwpr@gmail.com) or [editor@jwpr-science-line.com](mailto:editor@jwpr-science-line.com).

**We will send the full review results and the edited article after confirmation of the payment details.**

Please:

**Acknowledge the receipt of all the documents.**  
 Send the filled and signed **declaration form** for this article.  
 Pay **the publication fee (\$100 +VAT where applicable)**, required to final processing of your accepted paper.  
*If you want to pay the publication fee by Paypal you should pay \$20 more than the usual cost (\$100). Paypal will reduce \$20 from our account.*

k | Move to | Categorise | Snooze | ...

The Journal of World's Poultry Research - JWPR 1118 4

**YM** Yetti Marlida  
 Mon 29/04/2019 11:35  
 To: editor.jwpr <editorjwpr@gmail.com>

Thank you, tomorrow we will pay

Get [Outlook for Android](#)

...

[Reply](#) | [Forward](#)

**EJ** editor.jwpr <editorjwpr@gmail.com>  
 Sat 27/04/2019 07:01  
 To: Yetti Marlida

Dear Author,

Kindly inform us if you want to pay the fee and continue the review process otherwise the paper withdrawal from our database.


Kind Regards,  
 Daryoush Babazadeh  
 WVJ

**EJ** editor.jwpr  
 SiencelinePublication <http://www.science-line.com/index/> The Journal of World's Poultry Research <http://www.jwpr-science-line...> Fri 12/04/2019 06:12

Move to | Categorise | Snooze | ...

**Review results - JWPR 1118** 1

**EJ** editorjwpr <editorjwpr@gmail.com>  
 Thu 04/07/2019 00:22  
 To: Yetti Marlida

 1552518702--JWPR-1118.doc  
 127 KB

**SiencelinePublication**  
<http://www.science-line.com/index>

**Journal of World's Poultry Research**  
<http://www.jwpr.science-line.com/index/>

**Dear Respected Author,**

*We sincerely apologize for this lengthy process. The Sienceline publication has two departments in Iran and Turkey. The processing department of this publication is in Iran and because of some holidays this department was closed and the review process of your submitted article taken more time.*

It's my great pleasure to inform you that your paper encoded "JWPR-1118" has been reviewed by our selected reviewers and editors and accepted for publishing in [The Journal of World's Poultry Research](#) (JWPR) after **moderate revision**. The following attachments are sent with regards to you. Kindly revise the manuscript according to reviewer's comments, edited article and specially from [Authors Instruction](#) standpoint and submit the revised manuscript **within a week**.

Please:


- The language of paper should be revised one more time after necessary corrections by language editor.
- Acknowledge the receipt of all the documents.
- Highlight all corrections/additions with **green color** in revised manuscript.
- Please indicate the manuscript type

To include all authors name, affiliation and email address in revised word file.

Move to | Categorise | Snooze | ...

**Journal of World's Poultry Research - JWPR1131** 4

**EJ** editorjwpr <editorjwpr@gmail.com>  
 Tue 16/07/2019 19:29  
 To: Yetti Marlida

 1560124728--JWPR-1131(edi...  
 310 KB

**SiencelinePublication**  
<http://www.science-line.com/index>

**Journal of World's Poultry Research**  
<http://www.jwpr.science-line.com/index/>

**Dear Respected Author,**

It's my great pleasure to inform you that your paper encoded "JWPR-1131" has been reviewed by our selected reviewers and editors and accepted for publishing in [The Journal of World's Poultry Research](#) (JWPR) after **moderate revision**. The following attachment is sent with regards to you. Kindly revise the manuscript according to reviewer's comments, edited article and specially from [Authors Instruction](#) standpoint and submit the revised manuscript **within a week**.

Please:


- Acknowledge the receipt of all the documents.
- Highlight all corrections/additions with **green color** in revised manuscript.
- Please indicate the manuscript type
- Include all authors name, affiliation and email address in revised word file.
- Indicate the corresponding author
- Please add the Acknowledgements, Competing Interests, Ethics Committee Approval and Authors' Contributions to the revised form.  
 Example: (Preziosi et al., 2012; Mills et al., 2015)
- Add DOI number of references to the end of each reference.
- Design and send the [graphical abstract](#)

Fill and send the Declaration form

Move to | Categorise | Snooze | ...

Article code revision JWPR -1118 3

**EJ** editor jwpr <editorjwpr@gmail.com>  
Tue 20/08/2019 05:47  
To: Yetti Marlida

 1552518702--JWPR-1118 E2....  
137 KB

Dear Yetti Marlida,

Thank you for your collaboration to making the paper ready for publishing in JWPR.  
Kindly find the second revision of your paper encoded JWPR 1118. Please make it revise very carefully base on marginal comments and directly on the attached file which is the last version of your paper then send back the full revised manuscript as soon as possible for continuing the final process. The revisions should be highlighted with **yellow color**.  
**PLEASE DO NOT DELETE THE MARGINAL COMMENTS.**


The next issue of JWPR will be published on 25 June and we need to full revised article before 28 Aug 2019.  
--

With Kind Regards,  
Dr. Daryoush Babazadeh  
Editor in Chief  
Journal of World's Poultry Research (JWPR)  
Skype ID: drdaryoushb  
Tel: +90 5387708824 (Turkey)  
<http://www.jwpr.science-line.com/>  
<https://www.linkedin.com/groups/6583256>

Move to | Categorise | Snooze | ...

JWPR - Third Revision 2

**EJ** editor jwpr <editorjwpr@gmail.com>  
Sun 22/09/2019 20:28  
To: Yetti Marlida

 1552518702--JWPR-1118 E3....  
139 KB

Dear Author,

Thank you for your collaboration to making the paper ready for publishing in JWPR.  
Kindly find the third revision of your paper encoded JWPR 1118. Please make it revise very carefully base on marginal comments and directly on the attached file which is the last version of your paper then send back the full revised manuscript as soon as possible for continuing the final process.  
The revisions should be highlighted with **green color**.  
**PLEASE DO NOT DELETE THE MARGINAL COMMENTS.**

The next issue of JWPR will be published on 25 Sept and we needs to your urgent answer and full revisions.  
--

With Kind Regards,  
Dr. Daryoush Babazadeh  
Editor in Chief  
Journal of World's Poultry Research (JWPR)  
Skype ID: drdaryoushb  
Tel: +90 5387708824 (Turkey)  
[Website](#), [Linkedin](#), [Publons](#).

-----  
**News:** The updated [CiteScore](#), [SNIP](#) and [SJR](#) of Journal of World's Poultry Research are available in [Scopus](#) website.

Replv | Forward

Move to | Categorise | Snooze | ...

### graphical abstract - JWPR 1118


Some content in this message has been blocked because the sender isn't in your Safe senders list. | [I trust content from editorjwpr@gmail.com.](#) | [Show blocked content](#)


**EJ** editorjwpr <editorjwpr@gmail.com> Sun 22/09/2019 20:37  
To: Yetti Marlida


Dear Sir,

**Please send a graphical abstract for your accepted article within 5 days.**

**Graphical Abstract:** authors should provide a graphical abstract (a beautifully designed feature figure) to represent the paper aiming to catch the attention and interest of readers. Graphical abstract will be published online in the table of content. The graphical abstract should be *colored*, and kept within an area of 12 cm (width) x 6 cm (height) or with similar format. Image should have a minimum resolution of 300 dpi and line art 1200dpi. Note: *Height of the image should be no more than the width.* **Please avoid putting too much information into the graphical abstract as it occupies only a small space.** Authors can provide the graphical abstract in the format of PDF, Word, PowerPoint, jpg, or png, after a manuscript is accepted for publication. See more sample graphical abstracts in [archive](http://jwpr.science-line.com/index.php?option=com_content&view=article&id=2&Itemid=11) at [http://jwpr.science-line.com/index.php?option=com\\_content&view=article&id=2&Itemid=11](http://jwpr.science-line.com/index.php?option=com_content&view=article&id=2&Itemid=11).

 Inline image 1

 Inline image 1

 Inline image 2

We appreciate your interest in [Journal of World's Poultry Research \(JWPR\)](#)

--

With Kind Regards,  
Dr. Daryoush Babazadeh  
Editor in Chief  
[Journal of World's Poultry Research \(JWPR\)](#)

Move to | Categorise | Snooze | ...

### graphical abstract - JWPR 1118


Some content in this message has been blocked because the sender isn't in your Safe senders list. | [I trust content from editorjwpr@gmail.com.](#) | [Show blocked content](#)


**EJ** editorjwpr <editorjwpr@gmail.com> Sun 22/09/2019 20:37  
To: Yetti Marlida


Dear Sir,

**Please send a graphical abstract for your accepted article within 5 days.**

**Graphical Abstract:** authors should provide a graphical abstract (a beautifully designed feature figure) to represent the paper aiming to catch the attention and interest of readers. Graphical abstract will be published online in the table of content. The graphical abstract should be *colored*, and kept within an area of 12 cm (width) x 6 cm (height) or with similar format. Image should have a minimum resolution of 300 dpi and line art 1200dpi. Note: *Height of the image should be no more than the width.* **Please avoid putting too much information into the graphical abstract as it occupies only a small space.** Authors can provide the graphical abstract in the format of PDF, Word, PowerPoint, jpg, or png, after a manuscript is accepted for publication. See more sample graphical abstracts in [archive](http://jwpr.science-line.com/index.php?option=com_content&view=article&id=2&Itemid=11) at [http://jwpr.science-line.com/index.php?option=com\\_content&view=article&id=2&Itemid=11](http://jwpr.science-line.com/index.php?option=com_content&view=article&id=2&Itemid=11).

 Inline image 1

 Inline image 1

 Inline image 2

We appreciate your interest in [Journal of World's Poultry Research \(JWPR\)](#)

--

With Kind Regards,  
Dr. Daryoush Babazadeh  
Editor in Chief  
[Journal of World's Poultry Research \(JWPR\)](#)



### graphical abstract - JWPR 1118

Some content in this message has been blocked because the sender isn't in your Safe senders list. [I trust content from editorjwpr@gmail.com.](#) | [Show blocked content](#)

**EJ** editorjwpr <editorjwpr@gmail.com>  
Sun 22/09/2019 20:37  
To: Yetti Marlida



Dear Sir,

**Please send a graphical abstract for your accepted article within 5 days.**

**Graphical Abstract:** authors should provide a graphical abstract (a beautifully designed feature figure) to represent the paper aiming to catch the attention and interest of readers. Graphical abstract will be published online in the table of content. The graphical abstract should be *colored*, and kept within an area of 12 cm (width) x 6 cm (height) or with similar format. Image should have a minimum resolution of 300 dpi and line art 1200dpi. Note: *Height of the image should be no more than the width.* **Please avoid putting too much information into the graphical abstract as it occupies only a small space.** Authors can provide the graphical abstract in the format of PDF, Word, PowerPoint, jpg, or png, after a manuscript is accepted for publication. See more sample graphical abstracts in [archive](http://jwpr.science-line.com/index.php?option=com_content&view=article&id=2&Itemid=11) at [http://jwpr.science-line.com/index.php?option=com\\_content&view=article&id=2&Itemid=11](http://jwpr.science-line.com/index.php?option=com_content&view=article&id=2&Itemid=11).

Inline image 1

Inline image 1

inline image 2

We appreciate your interest in [Journal of World's Poultry Research \(JWPR\)](#)

--  
With Kind Regards,  
Dr. Daryoush Babazadeh  
Editor in Chief  
[Journal of World's Poultry Research \(JWPR\)](#)  
Client ID: d4d4a4a4a4