

Artikel Roni Pazla PJN1 2018

by Roni Pazla

FILE	RONI_PAZLA_PJN1_2018.PDF (543.85K)	WORD COUNT	4699
TIME SUBMITTED	01-MAR-2021 01:41PM (UTC+0700)	CHARACTER COUNT	25236
SUBMISSION ID	1521067163		

PJN

ISSN 1680-5194

PAKISTAN JOURNAL OF
NUTRITION

ANSI*net*

308 Lasani Town, Sargodha Road, Faisalabad - Pakistan
Mob: +92 300 3008585, Fax: +92 41 8815544
E-mail: editorpjn@gmail.com



Research Article

Supplementation of Minerals (Phosphorus and Sulfur) and *Saccharomyces cerevisiae* in a Sheep Diet Based on a Cocoa By-product

¹Roni Pazla, ¹Mardiati Zain, ¹H. Irsan Ryanto and ²Afrini Dona

¹Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, Andalas University, Kampus Limau Manis, 25163 Padang, Indonesia

²Department of Animal Nutrition, STIPER Sijunjung, Sijunjung, Indonesia

8 Abstract

Objective: The objective of this research was to determine the productivity of sheep that were fed a complete diet consisting of ammoniated cocoa waste (leaves and pod husks) supplemented with minerals (phosphorus and sulfur) and *Saccharomyces cerevisiae* (*S. cerevisiae*). **Methodology:** Four local sheep aged 10 months to 1 year were used in this study and were divided into 4 feed treatment groups (four sheeps per/treatment) with four replications for each treatment as follows: Treatment A served as the control and sheep were fed a complete diet that consisted of cocoa waste (fiber source)+concentrate, treatment B consisted of treatment A+0.4% phosphorous (P) and 1% sulfur (S), treatment C consisted of treatment B+1% *S. cerevisiae* and treatment D consisted of treatment B+2% *S. cerevisiae*. Each treatment had a fiber source: concentrate ratio of 40:60. The data collected included feed intake, nutrient digestibility, average daily weight gain and feed efficiency. **Results:** Treatment C had highly significant effects ($p < 0.01$) on dry matter intake, organic matter intake, crude protein intake, digestibility, crude fiber intake, average daily gain and feed efficiency. Thus, the complete diet consisting of ammoniated cocoa waste increased feed intake, digestibility, daily weight gain and feed efficiency. The use of ammoniated cocoa waste-based complete feed with minerals (P and S) and 1% *S. cerevisiae* yielded 130 g/day of average daily weight gain. **Conclusion:** The diet consisting of ammoniated cocoa waste supplemented with minerals (P and S) and 1% *Saccharomyces cerevisiae* can be effectively used for fattening sheep.

Key words: Cocoa waste, digestibility, *Saccharomyces cerevisiae*, phosphorus, sulfur, sheep, daily gain

Received: February 01, 2018

Accepted: April 26, 2018

Published: June 15, 2018

Citation: Roni Pazla, Mardiati Zain, H. Irsan Ryanto and Afrini Dona, 2018. Supplementation of minerals (phosphorus and sulfur) and *Saccharomyces cerevisiae* in a sheep diet based on a cocoa by-product. Pak. J. Nutr., 17: 329-335.

Corresponding Author: Mardiati Zain, Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, Andalas University, Kampus Limau Manis, 25163 Padang, Indonesia

Copyright: © 2018 Roni Pazla *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

The leaves and pod husks of cocoa plants are plantation by-products that have poor nutritional value due to their high fiber content¹. This waste is highly available in Indonesia. Indonesia is one of the largest cocoa producers worldwide with a production capacity of 760,429 t from a total area of 1,7222,315 ha². Cocoa has the potential to be used as an alternative to grass feed, which has begun to diminish. Lignin is a polymer compound that binds to cellulose and hemicellulose in plant tissues. High lignin content in cocoa waste limits digestibility³. Ammoniation treatment is one way to loosen lingo-cellulose and lingo-hemicellulose that exist in the cocoa waste, thereby making it easier for rumen bacteria to digest fiber and supply nitrogen for growth⁴. Zain³ reported that ammoniated pod husks changes 100% native grass into sheep rations. The combination of 75% ammoniated cocoa leaves+25% ammoniated cocoa pod husks can be used as a fiber source for ruminant rations¹. The supplementation of rumen microbial growth factors such as minerals (P and S) and direct-fed microbials is an alternative to improve nutrient digestibility of cocoa waste. Phosphorus is important for the metabolism and maintenance of the cell wall and cell membrane and it is a component of nucleic acids and high energy molecules (ATP, ADP and AMP), sulfur supports amino acid and microbial protein synthesis⁵⁻⁷. Supplementation of the minerals P and S into cocoa waste-based ruminant rations increases the digestibility of nutrients *in vitro*^{8,9}. Direct-fed microbial products, such as a *S. cerevisiae*, have been used to improve fiber digestibility¹⁰ and animal production¹¹. The beneficial effects of these microbial compounds are associated with an increase in cellulolytic bacteria¹² and they have been considered as a potential feed to improve nutrient digestibility in goats¹³. Therefore, the objective of the present study was to determine the productivity of sheep fed a complete diet consisting of ammoniated cocoa waste (leaves and pod husks) supplemented with minerals (phosphorus and sulfur) and *S. cerevisiae*.

MATERIALS AND METHODS

This study was conducted using experimental sheep cages at the Laboratory Nutrition of Animal Food Science, Faculty of Animal Science, Andalas University.

Experimental design: Four local sheep aged 10 months to 1 year were used in this research. Sheep were divided into

Table 1: Ingredient composition and nutrition of experimental diet (% DM)

Items	Diet (% DM)			
	A	B	C	D
Cacao leaves ammoniated	30	29.79	29.49	29.19
Cacao pod husk ammoniated	10	9.93	9.83	9.73
Rice brand	32	31.78	31.46	31.14
Corn	15	14.90	14.75	14.60
Coconut cake	10	9.93	9.83	9.73
Salt	1	0.99	0.98	0.97
Cattle Mix	2	1.99	1.97	1.95
P		0.40	0.40	0.40
S		0.30	0.30	0.30
<i>S. cerevisiae</i>			1	2
Total	100	100	100	100
Nutrition (%)				
Protein	12.78	12.78	12.86	12.93
TDN	63.73	63.73	63.73	63.73
Ether extract	4.78	4.78	4.78	4.78

feed treatment groups with four replications for each treatment as follows: Treatment A served as the control and consisted of a complete diet of cocoa waste [fiber source+concentrate], treatment B consisted of treatment A+0.4% phosphorous (P) and 0.3% sulfur (S), treatment C consisted of treatment B+1% *Saccharomyces* sp. and treatment D consisted of treatment B+2% *Saccharomyces*. Each treatment had a fiber source:concentrate ratio of 40:60. The composition of feed including the concentrate is shown in Table 1.

Digestibility trials were conducted using four animals for each treatment and animals were separated in individual pens. Sheep were fed *ad libitum* during the adaption period (7 days). During the last 14 days of each preliminary period, animals were equipped with fitted bags for the total collection. During the collection period, accurate records were kept for each individual intake. The total fecal excretion was collected once daily and 10% representative samples were dried at 60°C overnight and kept in sealed bags until the analysis. Dry matter, organic matter, crude protein and crude fiber were analyzed by standard AOAC methods¹⁴. Neutral detergent fiber (NDF) and acid detergent fiber (ADF) were analyzed by standard methods according to Van Soest *et al.*¹⁵.

Statistical analysis: The data were analyzed using a one-way analysis of variance (ANOVA) for a 4x4 Latin square design and Duncan's Multiple Range Test (DMRT)¹⁶ was used to test the differences between treatments with confidence intervals of 1%.

RESULTS AND DISCUSSION

The effect of the minerals (P and S) and *S. cerevisiae* on the feed intake, nutrient digestibility of

Table 2: Feed intake of sheep with experimental diet

Parameters (g/day)	Treatments			
	A	B	C	D
Dry matter intake (DMI)	436.74 ^a	497.39 ^b	574.09 ^c	555.51 ^c
Dry matter intake (% BW)	2.72 ^a	3.09 ^b	3.44 ^c	3.32 ^{b,c}
Organic matter intake (OMI)	374.71 ^a	429.05 ^b	492.75 ^c	476.82 ^c
Crude protein intake (CPI)	55.52 ^a	63.57 ^b	73.00 ^c	70.64 ^c

^{a-c}Means in the same row with different letters are significantly different ($p < 0.01$)

Table 3: Nutrient digestibility of sheep with experimental diet

Parameters (%)	Treatments			
	A	B	C	D
Dry matter	43.01 ^a	52.68 ^b	59.75 ^c	36.65 ^c
Organic matter	44.32 ^a	53.05 ^b	62.78 ^c	44.80 ^c
Crude protein	42.80 ^a	52.60 ^b	60.03 ^c	57.28 ^c
Crude fiber	36.65 ^a	44.80 ^b	57.28 ^c	55.74 ^c

^{a-c}Means in the same row with different letters are significantly different ($p < 0.01$)

Table 4: Average daily gain and feed efficiency of sheep with experimental diet

Parameters	Treatments			
	A	B	C	D
Average daily gain (g/day)	75.01 ^a	100.04 ^b	130.00 ^c	125.03 ^c
Feed efficiency (%)	17.02 ^a	20.53 ^b	22.64 ^c	22.52 ^c

^{a-c}Means in the same row with different letters are significantly different ($p < 0.01$)

cocoa waste and average daily weight gain of sheep are presented in Table 2-4.

Feed intake of ammoniated cocoa waste: The palatability of a feed material can be accessed from the level of feed intake by livestock. The higher the value of feed intake indicates the ration has good palatability. The statistical analysis showed that treatment C had highly significant effects ($p < 0.01$) on DMI, OMI and CPI, suggesting that supplementation of minerals (P and S) and 1% *S. cerevisiae* with ammoniated cocoa waste can increase the intake of dry matter, organic matter and crude protein. The significant difference ($p < 0.01$) in treatment C was due to increased feed intake resulting from increased digestibility attributed to the combination of mineral supplementation (P and S) and 1% *S. cerevisiae*. This result is consistent with the opinion of Febrina *et al.*¹⁷, who showed that the level of feed intake is influenced by the digestibility of the feed.

The addition of P and S minerals to treatment C increased the feed intake of dry matter rations. Feed intake will increase if the digestibility increases and if the rumen fermentation process is optimal. Increased feed intake in treatment C was due to P and S mineral supplementation causing increased digestibility, resulting in

faster rumen content emptying, causing the livestock to consume more quickly. Microbial growth and metabolism in the rumen was optimal because P is an important component for microbial growth^{18,19}. The S is an important component of rumen bacteria for microbial protein synthesis because the digestibility of fibrous feed is highly dependent on the enzymes produced by the microbes. According to Komisarczuk and Durand²⁰, the main function of S is to support the formation of amino acids containing S and microbial protein synthesis and S is also important for the synthesis of some vitamins (thiamin and biotin) and coenzymes. The increasing population of microbes, especially cellulolytic microbes, requires increasing amounts of the resulting enzymes. Increased microbial activity causes optimal fermentation processes to occur in the rumen. For example, Zain *et al.*⁵ stated that P and S mineral supplementation in fiber feed rations improves fiber digestibility.

The combination of minerals (P and S) and the addition of 1% *S. cerevisiae* can improve the palatability and digestibility of feed to allow faster emptying of the rumen content, which causes cattle to consume more feed. Yoon and Kim¹² claimed that *S. cerevisiae* in the rumen utilize oxygen to ensure anaerobic conditions for rumen bacteria and stimulate a population of certain rumen bacteria, which is followed by increased utilization of ammonia and lactic acid, resulting in a stable pH of the rumen. Anaerobic conditions and stable pH in the rumen allow for better microbial protein synthesis, which increases the total population of rumen bacteria and the digestibility of crude fiber. Increased digestibility of crude fiber also increases feed intake and the supply of nutrients to the intestines, which ultimately improves the overall production response. For example, Paryad and Rashidi¹³ demonstrated that digestibility of goat livestock rations supplemented with *S. cerevisiae* is higher than non-supplemented rations and Dann *et al.*²¹ reported an increase in the intake of dry matter ration in dairy cattle due to the addition of *S. cerevisiae*.

Treatment D was not significantly different from treatment C but was lower than treatment C, suggesting that addition of 1% *S. cerevisiae* of the dry matter (DM) ration was optimum for the growth of rumen microbes. This result is due to the fact that the more *S. cerevisiae* inoculum is added, the more nutrients are required. When nutrients are not sufficient, then *S. cerevisiae* cannot multiply, thus inhibiting its metabolic activity and resulting in the death of *S. cerevisiae*. This is in accordance with research conducted

by Sukma *et al.*²², who found a decrease in the fermentation of *Aspergillus terreus* A with increasing doses of inoculum.

Table 2 shows that the feed intake of organic matter has a positive correlation with the feed intake of dry matter, which is due to the fact that substances contained in organic matter are part of the dry matter. The amount of dry matter intake will affect the intake of organic matter. Thus, increased dry matter intake results in increased feed intake of organic material and vice versa. The mean feed intake of crude protein in this study ranged from 55.22-73.00 g/day, showing that it met the standard estimate of the adequacy requirement of crude protein based on body weight to achieve a daily body weight gain of 100-56-58 g/day²³. Thus, protein feed intake in this study is sufficient for basic living needs and growth because protein feed intake is closely related to the increase in body weight of livestock. The value of rough protein intake in treatment C (73.00 g day⁻¹) in this study was similar to that of Suparjo *et al.*²³, who used cocoa pod-based rations fermented with *Phytophthora palmivora* bacteria and reported a protein intake of 72 g/day.

Nutrient digestibility of ammoniated cocoa waste: The digestibility of food substances from the ration determines the quality of the ration because a higher digestibility rate of a ration means the ration is a good quality for livestock feed intake and is used for the metabolism process of the body. Table 3 shows that the highest digestibility values of dry matter, organic matter, crude protein and crude fiber ($p < 0.01$) were present in treatment C. The addition of P and S minerals to treatment C enhanced digestibility. Microbial growth and metabolism in the rumen were optimal because P is an enzyme activator and is involved in reactions related to energy liberation to form ATP. Jamarun *et al.*¹⁹ stated that microbial populations and microbial protein synthesis will increase with the addition of mineral doses of P. S is an important component for rumen bacteria for microbial protein synthesis because the digestibility of fiber feed is highly dependent on the enzymes produced by these microbes. Large microbial populations, especially cellulolytic microbes, can produce more enzymes. According to Komisarczuk and Durand²⁰, the main function of S is to support the formation of amino acids containing S and microbial protein synthesis and S is also important for the synthesis of some vitamins (thiamin and biotin) and coenzymes. Increased microbial activity creates an optimal fermentation process in the rumen, which is in line with research conducted by Jamarun *et al.*¹⁹, who reported that *in vitro* P supplementation enhanced the digestibility of fiber fractions (NDF, ADF, cellulose and hemicellulose) in fermented palm fronds. Additionally, a study conducted by

Zain *et al.*⁷ proved that the addition of S minerals is able to improve the digestibility of the amber rice straw.

The protein digestibility value is related to the condition of rumen fluid populations, especially the proteolytic population. The proteolytic proportions of each treatment were also different. The presence of a combination of P and S minerals improved the activity of proteolytic bacteria. The combination of 0.4% P and 0.3% S supplementation in ammoniated cocoa waste has also been shown to increase the digestibility of dry matter, organic matter and crude protein *in vitro*⁸. Supplementation of P and S minerals is indispensable because the feed ingredients from agricultural/plantation waste in Indonesia are deficient for these important minerals and their supplementation produces positive results in cattle performance⁵.

Saccharomyces cerevisiae reduces methane production by increasing the digestibility of organic matter. Carbohydrates are the most influential component of organic matter in the determination of organic matter digestibility because carbohydrates as energy producers (volatile fatty acid) are the largest components in feed. By increasing the proportion of propionate, *S. cerevisiae* can reduce methane. This result is consistent with the results of a study by Lync and Martin²⁵, who found increased propionate production, thereby reducing methane production in yeast-supplemented rations. Propionate is a glucogenic VFA because it can be catabolized into glucose, whereas acetate and butyrate are non-glucogenic (ketogenic) VFAs. The rumen fermentation system that leads to propionate will result in lower methane production, resulting in more VFA produced as a major energy source for ruminants and for the synthesis of microbial proteins²⁶. Low methane production will increase the value of hexose conversion as less energy is wasted in methane. This is in accordance with the opinion of Krehbiel *et al.*²⁷, who stated that the rumen fermentation system that leads to the synthesis of propionic acid is more advantageous because the energy wasted as methane gas is reduced.

Average daily gain and feed efficiency of ammoniated cocoa waste: The increase in body weight is a benchmark of growth rate. Table 4 shows the average daily weight gain for the experimental animals ranging from 75.01-130 g. The highest daily weight gain ($p < 0.01$) was in treatment C. The body weight increases with increasing digestibility due to increasing intake of dry matter, which directly affects the intake of organic matter and the digestibility of food substances. Differences in daily weight gain associated with digestibility and the amount of digestible food substances are further utilized for the

growth of body tissues. Growth is related to the feed consumed because the level of feed intake directly affects muscle growth, especially crude protein and energy. Increased daily gain in treatment C was due to mineral supplementation (P and S) and *S. cerevisiae*. This supplementation increased the growth of rumen microbes, especially cellulolytic bacteria.

The P minerals are required for the degradation of cell wall fractions and the cellulolytic bacteria requirement of these P minerals is higher than hemicellulolytic or amylolytic bacteria⁶. S is indispensable in rumen microbes for the formation of amino acids containing sulfur. According to Komisarczuk and Durand²⁰, the main function of S is to support the formation of amino acids containing S and microbial protein synthesis and it is also important for the synthesis of some vitamins (thiamin and biotin) and coenzymes. Metabolic processes involving growth, weight gain and enzyme or hormone activity are largely determined by the availability of the essential amino acid methionine. Methionine is an amino acid containing S.

Increasing the microbial population by P and S mineral supplementation causes increased digestibility and nutrient availability for growth, which also increases the body weight gain of livestock. This is consistent with Febrina *et al.*^{17,18}, who reported that the addition of minerals both *in vivo* and *in vitro* has a positive effect on rumen microbial activity. In the case of mineral deficiency status, the microbial fermentation activity in the rumen does not optimally occur, which will lead to low ration feeding efficiency and will eventually decrease the growth of livestock.

Saccharomyces cerevisiae stimulates the growth of rumen bacteria, especially cellulolytic bacteria, which increases digestibility and feed intake. Callaway and Martin²⁸ demonstrated that the increased feed intake of dry matter and ration organic ingredients also increases the feed intake of nutrients, which increases the nutrients available in the body and eventually increases weight gain. Wallace and Newbold²⁹ suggested that one of the benefits of direct-fed microbials to ruminants, especially in adult cattle, is to increase meat production.

Feed efficiency is influenced by feed intake of dry matter/day and weight gain/day. Table 4 shows that the average feed efficiency ranges from 17-22.65%. The highest ration efficiency was found in treatment C (22.64%) and the lowest efficiency was found in treatment A (17.02%). A ration will be more efficiently used if the ration is consumed in small quantities and is able to provide large daily body weight replacement. The average daily weight gain for treatment C (130 g/day) was attributed to the protein intake that is

sufficient for livestock needs for basic life and growth. The mean protein feed intake during the study ranged from 55.22-73.00 g/day. The amount of protein intake met the estimated raw protein adequacy standard based on body weight to reach 100 g/day. For example, Suparjo *et al.*²³ fed fermented cocoa pods to goats and these rations were 56-58 g day⁻¹ of protein feed intake to reach the 100 g day⁻¹ body weight.

The ration efficiency value in this study was higher than that in the study by Suparjo³⁰, who obtained a ration efficiency value of 18% for a fermented pod-based diet fed to goats. The ration efficiency in the present study was better in the treatment with ammonia cocoa waste supplemented with *S. cerevisiae* and minerals (P and S) because the *Saccharomyces cerevisiae* supplementation improved the level of feed intake and digestibility of food substances, resulting in better livestock growth.

2 CONCLUSION

According to the results of this experiment, the supplementation of minerals (P and S) and *S. cerevisiae* into ammoniated cocoa waste-based rations could improve feed intake, nutrient digestibility and sheep performance.

5 SIGNIFICANCE STATEMENT

This study demonstrates that supplementation with direct-fed microbials, namely, *S. cerevisiae* and minerals (P and S) can increase the growth performance, nutrient digestibility and feed efficiency of sheep. This study will enable researchers to further investigate the effects of *S. cerevisiae* and the minerals P and S on sheep that are fed an ammoniated cocoa waste-based diet, which was previously unexplored. Thus, a new hypothesis regarding the effects of supplementation with *S. cerevisiae* and the minerals P and S may be developed.

ACKNOWLEDGMENTS

This study was supported by an MP3EI Research Grant from Directorate General Higher Education, Department of National Education Republic of Indonesia contract no 06/UN.16/PL-SN/2016. This study would not have been possible without the cooperation and technical assistance of the Laboratory of Ruminant Nutrition of Animal Science Faculty of Andalas University.

REFERENCES

1. Rahman, J., M. Zain and Erpomen, 2013. Increased ruminant livestock productivity through optimization of ration utilization based on local raw materials in West Sumatra. Superior Research Reports of Higher Education. Andalas University, Padang, Indonesia.
2. Directorate General of Estate Crops, 2015. Tree crop estate statistic of Indonesia 2014-2016 cocoa. Ministry of Agriculture, Jakarta, Indonesia.
3. Zain, M., 2009. Substitution of native grass with ammoniated cocoa pod in sheep diet. Media Peternakan, 32: 47-52.
4. Suryani, H., M. Zain, R.W.S. Ningrat and N. Jamarun, 2016. Supplementation of Direct Fed Microbial (DFM) on *in vitro* Fermentability and degradability of ammoniated palm frond. Pak. J. Nutr., 15: 90-95.
5. Zain, M., N. Jamarun and Zulkarnaini, 2010. Effect of phosphorus and sulphur supplementation in growing beef cattle diet based on rice straw ammoniated. Asian J. Scient. Res., 3: 184-188.
6. Zain, M., N. Jamarun and A.S. Tjakradidjaja, 2010. Phosphorus supplementation of ammoniated rice straw on rumen fermentability, synthesized microbial protein and degradability *in vitro*. World Acad. Sci. Eng. Technol., 41: 1040-1042.
7. Zain, M., N. Jamarun and Nurhaita, 2010. Effect of sulfur supplementation on *in vitro* fermentability and degradability of ammoniated rice straw. Pak. J. Nutr., 9: 413-415.
8. Bastari, R., 2014. Evaluation of fiber fraction *in vitro* digestibility (NDF, ADF, Cellulose and Hemicellulose) ammoniated cocoa wastes with mineral addition of phosphorus and sulfur. Undergraduate Thesis, Faculty of Animal Science, Andalas University, Padang, Indonesia.
9. Yulhamedi, 2014. *In vitro* digestibility of dry matter, organic matter and crude protein on ammoniated cocoa waste with addition of mineral sulfur and phosphorus. Ph.D. Thesis, Faculty of Animal Science, Andalas University, Padang, Indonesia.
10. Zain, M., N. Jamarun, A. Arnim, R.W.S. Ningrat and R. Herawati, 2011. Effect of yeast (*Saccharomyces cerevisiae*) on fermentability, microbial population and digestibility of low quality roughage *in vitro*. Arch. Zootech., 14: 51-58.
11. Tang, S.X., G.O. Tayo, Z.L. Tan, Z.H. Sun and L.X. Shen *et al.*, 2008. Effects of yeast culture and fibrolytic enzyme supplementation on *in vitro* fermentation characteristics of low-quality cereal straws. J. Anim. Sci., 86: 1164-1172.
12. Yoon, I.K. and M.D. Stern, 1995. Influence of direct-fed microbials on ruminal microbial fermentation and performance of ruminants: A review. Asian-Aust. J. Anim. Sci., 8: 533-555.
13. Paryad, A. and M. Rashidi, 2009. Effect of yeast (*Saccharomyces cerevisiae*) on apparent digestibility and nitrogen retention of tomato pomace in sheep. Pak. J. Nutr., 8: 273-278.
14. AOAC., 1995. The Official Method of Analysis of the Association of Official Analytical Chemists. 16th Edn., Association of Official Analytical Chemists, Washington DC.
15. Van Soest, P.J., J.B. Robertson and B.A. Lewis, 1991. Methods for dietary fiber, neutral detergent fiber and nonstarch polysaccharides in relation to animal nutrition. J. Dairy Sci., 74: 3583-3597.
16. Steel, R.G.D. and J.H. Torrie, 1980. Principles and Procedure of Statistics. McGraw-Hill Book Co. Inc., New York, USA.
17. Febrina, D., N. Jamarun, M. Zain and Khasrad, 2017. Effects of using different levels of oil palm fronds (FOPFS) fermented with *Phanerochaete chrysosporium* plus minerals (P, S and Mg) instead of Napier grass on nutrient consumption and the growth performance of goats. Pak. J. Nutr., 16: 612-617.
18. Febrina, D., N. Jamarun, M. Zain and Khasrad, 2016. The effects of P, S and Mg supplementation of oil palm fronds fermented by *Phanerochaete chrysosporium* on rumen fluid characteristics and microbial protein synthesis. Pak. J. Nutr., 15: 299-304.
19. Jamarun, N., M. Zain, Arief and R. Pazla, 2017. Effects of Calcium (Ca), Phosphorus (P) and Manganese (Mn) supplementation during oil palm frond fermentation by *Phanerochaete chrysosporium* on rumen fluid characteristics and microbial protein synthesis. Pak. J. Nutr., 16: 393-399.
20. Komizarczuk, S. and M. Durand, 1991. Effect of Mineral on Microbial Metabolism. In: Rumen Microbial Metabolism and Ruminant Digestion, Jouany, J.P. (Eds.). INRA Publication, Versailles, France.
21. Dann, H.M., J.K. Drackley, G.C. McCoy, M.F. Hutjens and J.E. Garrett, 2000. Effects of yeast culture (*Saccharomyces cerevisiae*) on prepartum intake and postpartum intake and milk production of jersey cows. J. Dairy Sci., 83: 123-127.
22. Sukma, L.N., Zackiyah and G.G. Gumilar, 2010. Enrichment of unsaturated fatty acids in bran with solid fermentation using *Aspergillus terreus*. J. Sci. Chem. Technol., 1: 66-72.
23. Supardjo, K.G. Wirawan, E.B. Laconi and D. Mangunwidjaja, 2011. Goat performance fed with fermented cocoa pod husk. Media Peternakan, 34: 35-41.
24. Jamarun, N., M. Zain, Arief and R. Pazla 2017. Effects of calcium, phosphorus and manganese supplementation during oil palm frond fermentation by *Phanerochaete chrysosporium* on laccase activity and *in vitro* digestibility. Pak. J. Nutr., 16: 119-124.
25. Lynch, H.A. and S.A. Martin, 2002. Effects of *Saccharomyces cerevisiae* culture and *Saccharomyces cerevisiae* live cells on *in vitro* mixed ruminal microorganism fermentation. J. Dairy Sci., 85: 2603-2608.

26. Jamarun, N. and Z. Mardiaty, 2013. Basic Ruminant Nutrition. 1st Edn., CV Jasa Surya, Padang, Indonesia.
27. Krehbiel, C.R., S.R. Rust, G. Zhang and S.E. Gilliland, 2003. Bacterial direct-fed Microbials in ruminant diets: Performance response and mode of action. *J. Anim. Sci.*, 81: E120-E132.
28. Callaway, E.S. and S.A. Martin, 1997. Effects of a *Saccharomyces cerevisiae* culture on ruminal bacteria that utilize lactate and digest cellulose. *J. Dairy Sci.*, 80: 2035-2044.
29. Wallace, R.J. and C.J. Newbold, 1992. Probiotics for Ruminants. In: Probiotics: The Scientific Basis, Fuller, R. (Ed.). Chapman and Hall, London, pp: 317-363.
30. Suparjo, 2010. Improving nutritive value of cocoa pod husk as feedstuff by bioprocesses with *Phanerochaete chrysosporium* with Mn^{2+} and Ca^{2+} . Ph.D. Thesis, Bogor Agricultural Institute, Bogor.

ORIGINALITY REPORT

% **18**
SIMILARITY INDEX

% **17**
INTERNET SOURCES

% **16**
PUBLICATIONS

% **6**
STUDENT PAPERS

PRIMARY SOURCES

1 nexusacademicpublishers.com %**4**
Internet Source

2 docplayer.net %**2**
Internet Source

3 repo.unand.ac.id %**2**
Internet Source

4 scialert.net %**2**
Internet Source

5 Heni Suryani, M. Zain, R.W.S. Ningrat, N. Jamarun. "Effect of Dietary Supplementation Based on an Ammoniated Palm Frond with Direct fed Microbials and Virgin Coconut Oil on the Growth Performance and Methane Production of Bali Cattle", Pakistan Journal of Nutrition, 2017 %**1**
Publication

6 journal.unisza.edu.my %**1**
Internet Source

7 June Kim, Ahyeong Yun, Sung Hwoan Cho, Chang Hoon Kim. " Effects of sulphur type and concentration on growth performance, %**1**

body composition and resistance of juvenile sea cucumber () subjected to air and low salinity exposure stressors ", Aquaculture Research, 2021

Publication

8

Arief ., Elihasridas ., Simel Somen, Elly Roza, Roni Pazla, Rizqan .. "Production and Quality of Etawa Raw Milk Using Palm Oil Industry Waste and Paitan Plants as an Early Feed", Pakistan Journal of Nutrition, 2018

Publication

% 1

9

Roni Pazla, Novirman Jamarun, Mardiaty Zain, Arief .. "Microbial Protein Synthesis and in vitro Fermentability of Fermented Oil Palm Fronds by Phanerochaete chrysosporium in Combination with Tithonia (Tithonia diversifolia) and Elephant Grass (Pennisetum purpureum)", Pakistan Journal of Nutrition, 2018

Publication

% 1

10

www.tandfonline.com

Internet Source

% 1

11

ansinet.com

Internet Source

% 1

12

Submitted to Universitas Diponegoro

Student Paper

% 1

13

ijeab.com

Internet Source

% 1

EXCLUDE QUOTES ON

EXCLUDE MATCHES < 1%

EXCLUDE
BIBLIOGRAPHY ON