

Cytotoxicity Study of Ethanol  
Extract of Bintangor Leaf  
(*Calophyllum soulattri* Burm.f)  
on T47D Breast Cancer Cell Line  
(Cytotoxicity Study with MTT  
Assay Method) Elidahanum  
Husni\*, Fatma Sri Wahyuni,  
*by Pj-13-2-152 Pharmacognocny*

---

**Submission date:** 04-Mar-2021 04:34PM (UTC+0800)

**Submission ID:** 1523948243

**File name:** PJ-13-2-152.pdf (1.25M)

**Word count:** 3284

**Character count:** 17525

# Cytotoxicity Study of Ethanol Extract of Bintangor Leaf (*Calophyllum soulattri* Burm.f) on T47D Breast Cancer Cell Line (Cytotoxicity Study with MTT Assay Method)

Elidahanum Husni\*, Fatma Sri Wahyuni, Hanifa Nurul Fitri, Elsa Badriyya

Elidahanum Husni\*, Fatma Sri Wahyuni, Hanifa Nurul Fitri, Elsa Badriyya

Faculty of Pharmacy, Andalas University, INDONESIA

## Correspondence

Elidahanum Husni

Faculty of Pharmacy, Andalas University, INDONESIA

E-mail: elidahanumhusni@phar.unand.ac.id

## History

- Submission Date: 31-10-2020;
- Review completed: 22-12-2020;
- Accepted Date: 30-12-2020.

DOI : 10.5530/pj.2021.13.

## Article Available online

<http://www.phcogj.com/v13/i2>

## Copyright

© 2021 Phcogj.Com. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.



## ABSTRACT

**Introduction:** The public has used Bintangor leaf (*Calophyllum soulattri* Burm.f) for various medical treatments, including treated inflamed eyes and gout. **Aim:** This research aimed to determine the cytotoxic effect of ethanol extract and fraction of *Calophyllum soulattri* Burm.f leaf toward T<sub>47</sub>D breast cancer cell. **Methods:** The test used T<sub>47</sub>D breast cancer cells, the 3-4,5-dimethylthiazol-2yl -2,5-diphenyltetrazolium bromide (MTT) test method, and ELISA Reader to determine the absorbance. This method's principle was the presence of tetrazolium salts by the reductase system in the mitochondria of living cells formed purple formazan crystals. The used parameter was the value of IC<sub>50</sub>. **Results:** The result showed that ethanol extract, n-hexane fraction, ethyl acetate fraction, and butanol fraction did not have a cytotoxic effect on T<sub>47</sub>D breast cancer cell. The values of IC<sub>50</sub> respectively are 585.31 µg/ml; 409.33 µg/ml; 534.08 µg/ml; and 563.22 µg/ml. **Conclusion:** Ethanol extract and *Calophyllum soulattri* Burm.f leaf fraction did not have a cytotoxic effect on T<sub>47</sub>D breast cancer cells.

**Key words:** Bintangor Leaf, Breast Cancer Line, *Calophyllum soulattri* Burm.f, Cytotoxicity, MTT Assay, T<sub>47</sub>D.

## INTRODUCTION

Indonesia is an island country located in the equatorial zone and known as a country that has natural wealth with a lot of diversity of plants. With the richness of this plant, it supports the development of new drugs from plants. Various types of plants have been widely used for generations to prevent and treat diseases, including cancer treatment. One cancer drug obtained from plants is vinblastine obtained from *Vinca rosea*, podophyllin obtained from the plant roots of *Podophyllum peltatum*, and paclitaxel that brought from the skin of the pine tree *Taxus breviflora*<sup>1</sup>.

The ideal anticancer drugs should be able to counteract cancer cells without damaging normal tissues. However, the current agents did not meet these criteria<sup>2</sup>. Therefore, the research should be continued to find the ideal cancer drug. One of the plants that have anticancer activity is bintangor (*Calophyllum soulattri* Burm.f). It is used to wash the inflamed eyes; a decoction of the bark is used to treat vaginal discharge and rheumatism. The seeds of Bintangor are used to treat scabies, ulcers, and hair growers<sup>3</sup>.

Some compounds that have been isolated from *Calophyllum soulattri* are terpenoid derivatives called soulattron A<sup>4</sup> and friedelin<sup>5</sup>. Xanthone derivatives include soulattrin, caloxanthone B, caloxanthone C, macluraxanthone philattrin, braxanthone, trapeziphloxanthone, and anaxanthone<sup>5</sup>. The xanton literature study shows pharmacological activity as an anticancer

characterized by cytotoxic activity and its ability as an antioxidant associated with heterocyclic forms of these compounds. Some heterocyclic compounds have been shown as an anticancer, especially in inducing apoptosis<sup>6,7</sup>.

In a previous study, cytotoxic tests of *Calophyllum soulattri* plants were carried out on 4 types of cancer cells, namely HeLa cells (cervical cancer), MDA-MB-231 cells (breast cancer), LS174T (colon cancer), T98G (glioblastoma) and also tested on normal human cells HEK293 (human embryonic kidney cells). The researcher used dichloromethane, ethyl acetate, and methanol as solvents, where the value of IC<sub>50</sub> on MDA-MB-231 breast cancer cells fulfills cytotoxic requirements with ethanol solvents.

This researcher also tested the pure compounds in *Calophyllum soulattri*, which is friedelin, showing IC value<sub>50</sub> good against HeLa cells and T98G that is smaller than 30 µg/ml<sup>8</sup>.

Based on the research, there was no cytotoxic assay study of ethanol extract of *Calophyllum soulattri* leaves against T<sub>47</sub>D breast cancer cells. Research on breast cancer cells is essential because breast cancer is one of the biggest cancer killers due to cancer each year<sup>9</sup>. Handling of cancer, in general, is still dependent on chemotherapy derived from synthetic chemicals. However, these chemical compounds can cause multidrug resistance effects, a phenomenon in which cancer cells treated with certain drugs will become resistant to other drugs that have almost the same structure and mechanism of action<sup>10</sup>. However, chemotherapy affecting cancer cells can also affect normal cells that grow rapidly<sup>1</sup>.

**Cite this article:** Husni E, Wahyuni FS, Fitri HN, Badriyya E. Cytotoxicity Study of Ethanol Extract of Bintangor Leaf (*Calophyllum soulattri* Burm.f) on T47D Breast Cancer Cell Line (Cytotoxicity Study with MTT Assay Method). Pharmacogn J. 2021;13(2):

In this study, cytotoxic ethanol extract, n-hexane fraction, ethyl acetate fraction and butanol fraction of *Calophyllum soulattri* Burm. f. leaf were carried out for T<sub>47</sub>D breast cancer cells using the MTT Assay method. The parameter observed is the value of I<sub>50</sub>. The purpose of this study was to determine the cytotoxic effects of ethanol extract and the leaf fraction of bintangor (*Calophyllum soulattri* Burm. f) on T<sub>47</sub>D breast cancer cells.

## MATERIAL AND METHOD

### Material

The materials used for the cytotoxicity test were ethanol extract, n-hexane fraction, ethyl acetate fraction, and butanol fraction, leaves of *Calophyllum soulattri* Burm.f obtained from previous researchers at the Faculty of Pharmacy, Universitas Andalas. Breast cancer cells obtained from the Laboratory of Parasitology, Faculty of Medicine, Universitas Gadjah Mada.

The chemicals used are methyl sulfoxide (DMSO) (Sigma<sup>®</sup>), 70% alcohol, the Rosewell Park Memorial Institute (RPMI) medium, trypsin-EDTA (Sigma<sup>®</sup>), phosphate buffer saline (PBS) (Sigma<sup>®</sup>), trypan blue and MTT reagent 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (Sigma<sup>®</sup>), reagent stopper.

### Cell sub-culture

The medium in the flask was discarded, then add 2 ml FBS and stir slowly for five minute. Then FBS is removed and observed under a microscope. Then add 2 ml of trypsin-EDTA then stir slowly, incubate for 5-10 minutes at 37°C, 5% CO<sub>2</sub>, then observe the cell under the microscope. Cells that are ready for use will float and separate from the colon. Then the trypsin-EDTA solution containing the cell was centrifuged at 3000 rpm for 5 minutes. Remove the supernatant, and the pellets were cultivated in 3 mediums. Put in a new flask, stirring gently. Incubation at 37°C, 5% CO<sub>2</sub>.

### Calculation of number of cell

The flask medium is discarded. A total of 2 ml of trypsin-EDTA is added to the flask, incubated for 5-10 minutes. Then the solution is centrifuged at 3000 rpm and the supernatant is removed. 5 ml of

medium was added, the solution is resuspended homogeneously. 10 µl of cell suspense is taken, place it on each of the squares cell calculation boxes then calculated under a microscope.

### Cell layout

The cell suspension is made in the medium (measured quantity and volume), mixed perfectly. Add 180 suspensions to each well except the well in the first and last columns. The first and last columns are blanks that only contain medium, while the second column is a control that contains cell suspension in the medium. Incubate at 37°C, 5% CO<sub>2</sub> for 24 hours

### Stock solution and dilution of the test solution

Weigh the sample to about 10 mg in Eppendorf. The sample is dissolved in 100 µl DMSO for men get the solution concentration of 0.1 mg/µl. Samples must be completely dissolved in DMSO. Create the main solution by piping 50 µl of the stock solution with 995 µl in the Eppendorf tube.

Perform a level dilution by moving the 500µl test solution from the first tube to the second tube. Do the same for the next tube so that a solution will be obtained with a concentration of 500 µg/ml, 250 µg/ml, 125 µg/ml, 62.5 µg/ml and 31.25 µg/ml in each well on the 96 plates well.

### Cell proliferation test

Test plates containing cell wells and incubated for 24 hours. Discard the media on the well 96 plates by turning the plate 180°. Samples are inserted on the plate where the test solution is started from the lowest concentration. Transfer 20 µl of the test solution to each test well into the control well enter 180 µl of media, and the blank is only filled with 200 µl of media. The plate is re-incubated for 24 hours in an incubator 37°C, 5% CO<sub>2</sub>.

Piped 10µl of MTT mg/ml solution into each well. Incubate for 3-4 hours at 37°C, CO<sub>2</sub>. After 3-4 hours, there will be a purple colour from the formal form of the call to prayer formed. Added 100µl reagent stopper in each well. The plate is wrapped using paper or aluminium foil, leave it for 24 hours at room temperature then read absorption used ELISA reader




Figure 1: Graph of the relationship between concentrations of ethanol extract of *Calophyllum soulattri* Burm.f with % viability of T<sub>47</sub>D breast cancer cell.




Figure 2: Graph of the relationship between the concentration of n-hexane fraction of *Calophyllum soulattri* Burm.f with % viability of T<sub>47</sub>D breast cancer cells.




Figure 3: Graph the relationship between the concentration of ethyl acetate fraction of *Calophyllum soulattri* Burm.f with % viability of T<sub>47</sub>D breast cancer cells.




Figure 4: The graph of the relationship between the concentration of butanol fraction of *Calophyllum soulattri* Burm.f with % viability of T<sub>47</sub>D breast cancer cells.

## Analysis of data

By using absorbance data obtained from measurements, it can be determined the percentage of cells inhibited using the following formula<sup>11</sup>:

$$\% \text{ of cell viability} = \frac{A_{\text{treatment}} - A_{\text{blank}}}{A_{\text{control}} - A_{\text{blank}}} \times 100\%$$

\*A = Absorbance

The relationship between the concentration of the test solution and cell viability can be displayed in the form of a concentration chart with live cell presentation. From the graph, the price of IC<sub>50</sub> can be determined from each solution.

## RESULTS AND DISCUSSION

The cytotoxic effect of (*Calophyllum soulattri* Burm. f) fraction was conducted on T<sub>47</sub>D breast cancer cells using the MTT assay method obtain IC<sub>50</sub> values of ethanol extract 585.31 µg/ml, a fraction of n-hexane 409.33 µg/ml, ethyl acetate fraction 534.08 µg/ml and butanol fraction 563.22 µg/ml.

## DISCUSSION

The MTT method is one of the colourimetric methods using a colour reaction in the measurement of a number of living cells based on changes in solution 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyltetrazolium bromide (MTT) which is a formazan crystal which purple by mitochondria which are active in living cells. MTT is absorbed into living cells and broken down through a reduction reaction by the reductase enzyme in the mitochondrial respiration chain to formazan which is insoluble in water<sup>12</sup>.

The parameters used for cytotoxic tests are IC<sub>50</sub> values. The value of IC<sub>50</sub> is the concentration of a substance where the substance can inhibit half of the activity or cell growth tested. The IC<sub>50</sub> value indicates a concentration value that can inhibit 50% of cell proliferation and points to the potential for toxicity of a compound to cells. The greater the price of IC<sub>50</sub>, the more the compound has no potential as a cytotoxic compound.

According to The American National Cancer Institute, the compound is said to have cytotoxic activity if the IC value is  $IC_{50} < 30 \mu\text{g/ml}$ <sup>13</sup>. The MMT assay has been done and the result showed that the IC<sub>50</sub> of ethanol extract was 585.31 µg/ml, n-hexane fraction was 409.33 µg/ml, ethyl acetate fraction was 534.08 µg/ml and butanol fraction was 563.22 µg/ml. Thus, the ethanol extract, n-hexane fraction, ethyl acetate fraction and butanol fraction have no cytotoxic activity because of the value of IC<sub>50</sub> was more than 100 µg/ml.

In a previous study, cytotoxic testing of *Calophyllum soulattri* Burm.f on MDA-MB-231 breast cancer cells showed a cytotoxic effect. The extract used in the study used ethanol and methanol solvents<sup>8</sup>. In these studies, conducted using T<sub>47</sub>D breast cancer cells showed a non-toxic effect. It occurs because of the expression of receptors in different cells. Where prescription expression on MDA-MB-231 cells is epidermal growth factor (EGF) which can change growth factor alpha (TGF), whereas in T<sub>47</sub>D cells expression of receptors can express positive estrogen and progesterone receptors<sup>13,14</sup>. Progesterone receptors are receptors that can stimulate growth, differentiation, and development of breast cancer. Where excessive exposure to these receptors can cause breast cancer<sup>15</sup>. It is suspected that T<sub>47</sub>D breast cancer cells given treatment with *Calophyllum soulattri* Burm.f extract cannot occupy and interact with estrogen and progesterone receptors so that it does not cause any effect.

Based on the results of the research that has been done, it can be concluded that the ethanol extract, n-hexane fraction, ethyl acetate and butanol do not have cytotoxic activity. Therefore, further research is still needed by carrying out cytotoxic testing of bintangor leaf extract on other cancer cells.

## CONCLUSION

The IC<sub>50</sub> of ethanol extract was 585.31 µg/ml, n-hexane fraction was 409.33 µg/ml, ethyl acetate fraction was 534.08 µg/ml and butanol fraction was 563.22 µg/ml. Thus, the ethanol extract, n-hexane fraction, ethyl acetate fraction and butanol fraction have no cytotoxic activity because of the value of IC<sub>50</sub> was more than 100 µg/ml. Based on that results ethanol extract and bintangor leaf fraction (*Calophyllum soulattri* Burm. f) do not have a cytotoxic effect on T<sub>47</sub>D breast cancer cells. This solutions should be tested on another cancer cell to determine the cytotoxic effect of extract of this plant.

## ACKNOWLEDGEMENT

The authors would like to deliver a special appreciation to Research Institutions and Community Service (LPPM) of Andalas University for the financial support on professor acceleration research scheme.

## CONFLICTS OF INTEREST

The authors declare that there is no conflicts of interest regarding the publication of this article.

## ABBREVIATIONS


EGF: Epidermal growth factor; HeLa Cell: One of the cells grown from the cervical cancer of a young African-American woman, The designation "HeLa" was taken from the name of Henrietta Lacks; IC<sub>50</sub>: Inhibitory Concentration; MDA-MB: stands for "M.D. Anderson and MB stands for Metastasis Breast cancer; MTT Assay: Microculture tetrazolium; PBS: Phosphate Buffer Saline; TGF: Tumor Growth Factors, SDS: Sodium dodecylsulphate.

## REFERENCES

1. Tjay, TH. & Rahardja, K. Obat-obat Penting. Ed. 6. Jakarta: Elex Media Computindo; 2007.
2. Katzung, Bertram G. Farmakologi Dasar dan Klinik. Ed. 12. Jakarta: Penerbit EGC; 2016.
3. Heyne K. Tumbuhan Berguna Indonesia I. Jakarta: Badan Penelitian dan Pengembangan Kehutanan. Departemen Kehutanan Republik Indonesia; 1998.
4. Nigam KS, Banerji S, Rebuffat S, Cesario M, Pascard C and Bodo B. Soulattrone A, A C24 terpenoid from *Calophyllum soulattri*. *Phytochemistry*. 1988;27(2).
5. Mah SH, Gwendoline CLE, Mawardi R, Yun HTY, Moh AS, Soek, ST. A New Pyranoxanthone from *Calophyllum soulattri*: Department of Chemistry Faculty of Science, Universiti Putra Malaysia *Molecules*. 2011;16.
6. Kamal A, Mallareddy A, Suresh P. Heterocycles as Inducers of Apoptosis. *Journal Apoptotic Regulators*. 2012;45-92.
7. Pinto MMM, Sousa ME, Nascimento MSJ. Xanthone Derivates: New Insights in Biological Activities. 2005;2517-38.
8. Mulyadi T, Rachmadiarti F, Prameswari A, Agyani VUW, Saputri RD, Tjahjandarie TS, et al. Airlangins AB, two new isoprenylated benzofuran-3-ones from the stem bark of *Calophyllum soulattri*. *Natural Product Research*. 2018;32(13):1493-8.
9. Kementerian Kesehatan Republik Indonesia. Pusat data dan informasi. Jakarta Selatan: Kementerian Kesehatan RI; 2015
10. Baguley BC. Multidrug Resistance Mechanism in Cancer: *Molecular Biotechnology*; 2010;46(3):308-16.

11. Park, Chung-Min, and Ming Xian. Use of phosphorodithioate-based compounds as hydrogen sulfide donors. In *Methods in Enzymology*. vol. 554, pp. 127-142. Academic Press, 2015.
12. Stockert JC, Blázquez-Castro A, Cañete M, Horobin RW, Villanueva Á. MTT assay for cell viability: Intracellular localization of the formazan product is in lipid droplets. *Acta histochemica*. 2012;114(8):785-96.
13. Suffness M, Pezzuto JM. *Methods in plant biochemistry: assays for bioactivity*. Academic Press, London 1990:71-133.
14. Clarke. RB., Howell, A., Anderson, E. Type 1 IGF Receptor Gene Expression in Normal Breast Tissue Treated with Estrogen and Progesterone. *Br J Cancer*. 1997;75:2776-8.
15. CCRC Line. *Regulasi siklus sel*. Yogyakarta: Universitas Gadjah Mada; 2012.

## GRAPHICAL ABSTRACT



## ABOUT AUTHORS



**Dr. Elidahanum Husni, Apt.:** Dr. Elidahanum Husni, M.Si, Apt.: Currently as a lecturer at the Faculty of Pharmacy, University Andalas. Graduated from Faculty of Pharmacy Universitas Andalas in 1986, then Master Program in 1995 at School of Pharmacy Bandung Institute of Technology and Doctoral Program in Department Biomedical, Faculty of Medicine, University Andalas in 2015. The research and expertise is Pharmacognosy



**Prof. Dr. Fatma SriWahyuni, Apt.:** Currently a lecturer at the Faculty of Pharmacy, Andalas University. Got her undergraduate degrees from Andalas University in 1998 and finished her PhD from University Putra Malaysia in 2010. Her research is in activity studies of Sumatran Plants.



**Elsa Badriyya, M. Si. Apt.:** Currently a lecturer at the Faculty of Pharmacy, Andalas University. Graduated from the School of Pharmacy, Bandung Institute of Technology in 2016, then the Master's Program in 2017 at the Bandung Institute of Technology, School of Pharmacy. Currently working as a lecturer in Pharmacology and Clinical Pharmacy at the Faculty of Pharmacy, Andalas University.



**Hanifa Nurul Fitri:** She is a student at Faculty of Pharmacy, Andalas University who has been involved in research that studies Cytotoxicity Study of Ethanol Extract of Bintangor Leaf (*Calophyllum soulattri* Burm.f) on T47D Breast Cancer Cell Line (Cytotoxicity Study with MTT Assay Method).

**Cite this article:** Husni E, Wahyuni FS, Fitri HN, Badriyya E. <sup>1</sup> Cytotoxicity Study of Ethanol Extract of Bintangor Leaf (*Calophyllum soulattri* Burm.f) on T47D Breast Cancer Cell Line (Cytotoxicity Study with MTT Assay Method). Pharmacog J. 2021;13(2):

# Cytotoxicity Study of Ethanol Extract of Bintangor Leaf (Calophyllum soulattri Burm.f) on T47D Breast Cancer Cell Line (Cytotoxicity Study with MTT Assay Method) Elidahanum Husni\*, Fatma Sri Wahyuni,

## ORIGINALITY REPORT

17%

SIMILARITY INDEX

18%

INTERNET SOURCES

11%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

1

[scholar.unand.ac.id](http://scholar.unand.ac.id)

Internet Source

13%

2

[repo.unand.ac.id](http://repo.unand.ac.id)

Internet Source

4%

Exclude quotes On

Exclude matches < 3%

Exclude bibliography On