

Isolation and Characterization of Lactic Acid Bacteria Proteases from Bekasam for use as a Beef Tenderizer

by Yuherman Yuherman

Submission date: 25-Jan-2022 12:09PM (UTC+0800)

Submission ID: 1747607183

File name: 361-367_PJN_2018,Isolation.pdf (571.62K)

Word count: 4471

Character count: 23404

PJN

ISSN 1680-5194

PAKISTAN JOURNAL OF
NUTRITION

ANSI*net*

308 Lasani Town, Sargodha Road, Faisalabad - Pakistan

Mob: +92 300 3008585, Fax: +92 41 8815544

E-mail: editorpjn@gmail.com



Research Article

Isolation and Characterization of Lactic Acid Bacteria Proteases from Bekasam for use as a Beef Tenderizer

¹Afriani, ²Arnim, ³Yetti Marlida and ²Yuherman

¹Department of Animal Production, Faculty of Animal Science, Jambi University, Indonesia

²Department of Animal Production, Faculty of Animal Science, Andalas University, Padang, Indonesia

³Department of Animal Nutrition, Faculty of Animal Science, Andalas University, Padang, Indonesia

Abstract

Background and Objective: Proteases are important enzymes and have high economic value due to their wide applications in the food industry as a meat tenderizer. Protease use in the food industry necessitates an understanding of the capabilities and influencing factors of these enzymes to accelerate enzymatic reactions. This study aimed to isolate and characterize the proteases of lactic acid bacteria (LAB) from Bekasam. **Methodology:** The samples were obtained from the third, fifth, seventh, ninth and eleventh day of fermentation to isolate the proteolytic LAB. Characterization of proteases includes the incubation time, casein substrate concentration, optimum temperature and pH, metal ion contents and stability. The LAB with the highest protease activity is identified molecularly and isolated through 16S rDNA sequencing and phylogenetic analysis based on the Neighbor Joining method. **Results:** The results showed that the best isolate was BAF-715 because it had the highest protease activity (18.84 U mL⁻¹) at 40 h of incubation. The optimum activity of this protease on a casein substrate at 2.5% occurred at an incubation temperature of 40 °C at pH 7 and in the presence of Mg²⁺ and Mn²⁺ (5 mM) as activators. Based on molecular DNA identification, the BAF-715 isolate is determined to be *Pediococcus pentosaceus*. **Conclusion:** A protease produced by *Pediococcus pentosaceus* showed the highest proteolytic activity, making it the best protease for application as a beef tenderizer.

Key words: Lactic acid bacteria, protease, bekasam

Received: September 27, 2017

Accepted: May 04, 2018

Published: July 15, 2018

Citation: Afriani, Arnim, Yetti Marlida and Yuherman, 2018. Isolation and characterization of lactic acid bacteria proteases from bekasam for use as a beef tenderizer. Pak. J. Nutr., 17: 361-367.

Corresponding Author: Afriani, Department of Animal Production, Faculty of Animal Science, Jambi University, Indonesia

Copyright: © 2018 Afriani *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Bekasam is a traditional salt-fermented product from fish, which are found in many areas in Indonesia. Many microorganisms contain lactic acid bacteria (LAB)¹. LAB can produce several metabolites; they can also produce proteolytic enzymes around cell walls, in cytoplasmic membranes and in cells². Protease is an enzyme that breaks the peptide bonds in proteins to produce simpler compounds, such as small peptides and amino acids.

Protease is a very important enzyme in food and non-food industries. Some uses of proteases in the food industry include reducing turbidity in the beer industry, reducing gluten in the bread industry, agglomerating milk in the cheese industry and tenderizing meat in the meat processing industry^{3,4}. Proteases can be easily isolated from various sources of plants, animals and microbes through fermentation processes⁵.

Microbes used as an enzyme source are more advantageous than plant and animal sources because microbes can grow rapidly on inexpensive substrates and can be easily enhanced by manipulating the growth conditions and applying genetic engineering⁶ and biotechnology⁷. The existence of a superior microbe is an important factor in the production of enzymes.

Several types of LAB have proteolytic activity, such as *L. plantarum*, *L. brevis*, *L. pentosus*, *Pediococcus acidilactici*, *Pediococcus pentosaceus* and *Lactobacillus* spp⁸⁻¹⁰. LAB are safe microbes and enzymes produced from LAB can be used directly on food.

The ability of proteases to accelerate reactions is influenced by several factors, such as pH, temperature and metals serving as activators or inhibitors, that change enzymatic characteristics. Proteases from *Pediococcus acidilactici* have an optimum activity at pH 4, an incubation temperature of 28°C and a 2% casein concentration¹¹. The protease activity from *Lactobacillus plantarum* was highest at 4.43 U mg⁻¹, pH 7, an incubation temperature of 47°C and with a Mg²⁺ metal activator¹². The 220 proteases from *P. pentosaceus* stabilized at pH 4-7, an incubation temperature of 20-40°C and with metal activators/inhibitors of Ca²⁺, Zn²⁺, Mg²⁺ and Fe³⁺; the protease activity was inhibited by EDTA and sodium dodecyl sulfate¹⁰.

From the description above, this study was conducted to determine the optimum conditions for the production of protease enzymes by LAB.

MATERIALS AND METHODS

Isolation of lactic acid bacteria: LAB was isolated from Bekasam samples on the third, fifth, seventh, ninth and

eleventh fermentation days of 2 replications and the bacteria were grown on MRS agar medium. From each sample, five different colonies were grown on MRS agar medium+CaCO₃ using the pour plate method. The isolate, which formed a clear zone, was then purified by the streak plate method on the same medium.

Test of proteolytic lactic acid bacteria: A LAB proteolytic test was used with an agar skim milk (ASM) medium with 3% skim and 3% agar composition. The proteolytic isolates were characterized by the formation of a clear zone on the agar skim medium and then, the clear zone was measured. The proteolytic index was calculated by measuring the ratio of the diameter of the clear zone/colony diameter (R).

Production and isolation of protease enzymes: The production and isolation of protease enzymes are performed by adding one dose of isolate to 50 mL MRS broth and incubating the sample in a shaker incubator with an agitation speed of 150 rpm at 37.5°C for 24 h. The inoculum was then inoculated on protease production media with a composition of 2% skim milk, 0.5% peptone, 0.1% yeast extract, 2% glucose, 0.1% NaCl, 0.008% KH₂PO₄, 0.01% MgSO₄·7H₂O and 0.04% (NH₄)₂SO₄. The sample was incubated in a shaker incubator with an agitation speed of 150 rpm at 37.5°C with various production times, namely, 8, 16, 24, 32, 40, 48 and 56 h. The sample was centrifuged at a rate of 12000 rpm for 15 min at 4°C to separate the supernatant and its sediment. The supernatant is a crude extract of the protease and is used for testing the protease activity.

Protease activity measurements: Protease activity was measured by the method of Bergmeyer *et al.*¹⁴ using casein substrate Hammerstein 2% (b/v). A unit of protease activity is defined as the amount of enzyme that can produce one μmol of tyrosine product per minute under the measurement conditions. Isolates with the highest proteolytic activity were tested further.

Protease characterization: Characterization of proteases includes the effect of the casein substrate concentration on enzyme activity, which was tested by reacting an enzyme solution with a casein concentration of 0.5, 1, 1.5, 2, 2.5 and 3%.

The pH effect on the enzyme activity was tested by using different buffer treatments. The universal buffers used had pH values of 3, 4, 5, 6, 7, 8 and 9.

The effect of temperature on enzyme activity was tested on the first incubation carried out at different temperatures, i.e., 30, 40, 50, 60, 70, 80 and 90°C.

The effect of the metal ions Mg^{2+} , Cu^{2+} , Ca^{2+} , Fe^{2+} , Zn^{2+} and Mn^{2+} in sulfate salts with a concentration of 5 mM as an activator or inhibitor on protease activity was tested using an optimum pH buffer and optimum temperature. As a benchmark, the treatment used buffer without the addition of metal ions (nonmetal).

Molecular identification of selected LAB isolates: LAB isolates with the highest proteolytic activity were molecularly identified by extracting DNA encoding 16S rRNA. DNA sequencing of the encoded 16S rRNA was performed by 1st Base, PT Genetics of Indonesia. An analysis of the sequencing results was performed using the BLASTing sequence of nucleotides from the sequencing results of 16S rRNA with the database available at www.ncbi.nlm.nih.gov. Phylogenetic analysis was performed with the Neighbor Joining method using the program Molecular Evolutionary Genetics Analysis (MEGA) version 6.0 with Multiple Sequence Comparison by Log Expectation (MUSCLE).

Amplification of DNA encoding 16S rRNA with PCR (polymerase chain reaction): The amplification reaction of the DNA sample was carried out in 0.2 mL PCR tubes. To each PCR reaction tube was added 0.25 μ L RBC Taq (5 U mL^{-1}), 10 \times 5 μ L Taq buffer (containing Mg^{2+}), 4 μ L total of 2.5 mM dNTP and universal primer 63F (5'-CAGGCCTAACACATGCAAGTC-3') and universal primer 1387R (5'-GGGCGWGTGTACAAGGC-3') at 1.25 μ L (20 pmol) and 1.25 μ L (20 pmol), respectively; 2.5 μ L (100 ng) of genomic extract and ddH₂O up to 50 μ L were also added. PCR amplification was performed using a PTC 100 PCR instrument (MJ Research, Inc.) at an initial denaturation temperature of 95°C for 5 min, followed by 30 cycles of annealing at 94°C for 30 sec, as well as an extension at 50°C for 1 min, 72°C for 2 min and a final stage of 72°C for 5 min. The PCR products were stored at 4°C for further examination using 1% agarose electrophoresis in 1 \times TAE, 100 V for 30 min¹⁵.

Analysis of the DNA sequence of the genes encoding 16S rRNA: The DNA sequencing of the genes encoding 16S rRNA was performed by using 1st Base, PT, Indonesian Genetics. The sequencing analysis was performed by BLAST to obtain the nucleotide sequence from the 16S rRNA sequencing results with the available database at www.ncbi.nlm.nih.gov. Then, a multiple alignment was performed by using the Clustal W Program. Furthermore, visualization of kinship was performed using a phylogenetic tree combination via the Neighbor Joining method¹⁶ with the MEGA 6.0 program¹⁷.

RESULTS AND DISCUSSION

Isolation of lactic acid bacteria and proteolytic test of protease-producing bacteria: The isolation process obtained 50 bacterial colonies. Then, from these 50 colonies, which provided clear zones in the MRS and CaCO₃ media, 25 isolates showed the characteristics of LAB. The formation of clear zones indicated that the bacteria produced an excess of secondary metabolites (lactic acid) so that an excess of lactic acid was shown by the clear zones around the bacterial colonies.

The 25 isolates of LAB were tested for proteolytic activity, as shown by their ability to form a clear zone on the agar skim medium. When viewed from the protease activity of the clear zone/colony diameter (R) size, the number of isolates exhibiting high protease activity ($R \geq 2$) was less than that of isolates with $R \leq 2$ (Table 1). Three isolates of LAB with a high proteolytic index ($R \geq 2$) were BAF-514, BAF-715 and BAF-1121 (Table 2).

Protease enzyme activity: A qualitative test of protease activity showed that the BAF-514, BAF-715 and BAF-1121 isolates could produce extracellular protease by forming a large clear zone around the cell colony. The larger clear zone that was formed indicated that the microbes were highly capable of changing the substrate that was contained in the medium¹⁸.

The proteolytic index of isolates BAF-514, BAF-715 and BAF-1121 was 2.06, 2.29 and 2.02, respectively. The protease activity of the BAF715 isolate in the skim milk agar medium was demonstrated by the presence of clear zones around the colony (Fig. 1). The results reported by Wikandari *et al.*⁹ indicated that the LAB with a proteolytic index ≥ 2 were *L. plantarum* N2352, *L. plantarum* B1765, *L. plantarum* T2565, *L. plantarum*

Table 1: Number of isolates of lactic acid bacteria and proteolytic lactic acid bacteria from *Bekasam*

Isolates	Total isolates
Acid-forming bacteria	50
Lactic acid bacteria (LAB)	25
LAB proteolytic	17
LAB proteolytic $R \geq 2$	3
LAB proteolytic $R \leq 2$	14
LAB non-proteolytic	8

Table 2: Chosen isolates of lactic acid bacteria and the proteolytic index

Isolates	Proteolytic index
BAF-514	2.06
BAF-715	2.29
BAF-1121	2.02




Fig. 1: Clear zone as a qualitative indicator of LAB proteases




Fig. 2: Activity of isolate enzymes BAF-514, BAF-715 and BAF-1121 for different incubation times




Fig. 3: Effect of the casein substrate concentration on the protease activity of the BAF-715 isolate

B1465, *L. pentosus* B2555 and *Pediococcus pentosaceus* B1666. Interestingly, research by Yusmarini *et al.*⁸ proved that LAB from the genus *Lactobacillus* R.1.3.2 and R.11.1.2 has a proteolytic zone of 1.725 and 1.650 cm, respectively.




Fig. 4: Effect of pH on the protease activity of lactic acid bacteria BAF-715

The results for the protease activity as a function of incubation time (Fig. 2) showed that the optimum activity of the BAF-514 and BAF-1121 isolate proteases at 48 h of incubation time was 11.32 and 9.65 U mL⁻¹, respectively. This result indicates that the proteases were produced at a stationary phase, namely, the peak cell growth phase. Meanwhile, the isolate protease activity of BAF-715 at 40 h of incubation time was 18,844 U mL⁻¹, revealing that the protease was produced along with the cell growth before reaching the stationary phase. The value of this protease activity was higher than that of Putranto¹⁹. The *Lactobacillus acidophilus* bacteria were at the highest activity when the incubation time was 18 hours (0.752 U mg⁻¹). Because isolate BAF-715 showed the highest proteolytic activity, this isolate was selected for further research.

Characterization of the crude extract protease of the selected LAB isolate (BAF-715)

Effect of the casein substrate concentration: The effect of the casein substrate concentration on the protease activity is shown in Fig. 3. The highest protease activity was obtained at a concentration of 2.5% at 88.4 U mL⁻¹. Meanwhile, the protease activity decreased when the concentration of the casein substrate was 3%. Comparing the results of Oke and Onilude¹¹ study using *P. acidilactici* and the results of Akinkugbe and Onilude²⁰ study using *Lactobacillus acidophilus*, the highest protease activity was achieved at a casein concentration of 2%.

Effect of pH: The pH activity profile of an enzyme describes the pH at the time that the proton should be donated or received on the catalytic side of the enzyme to be at the desired ionization level. Figure 4 shows that the isolate protease activity of LAB has an optimum pH of 7 with an enzyme activity of 19.89 U mL⁻¹. Changes in alkaline or acidic

pH conditions result in a decrease in enzyme activity. It is estimated that the enzyme activation change is caused by ionization of the enzyme's ionic group, on the active side or on the other side, thus indirectly affecting the active side. The ionic group plays a role in maintaining the active side conformation for binding the substrate and converting it into the product. These enzyme activity results were higher than that reported in the research of Putranto¹⁹, in which *Lactobacillus acidophilus* bacteria had an optimum pH of 6 with an enzyme activity of 13.5 units mg⁻¹; the protease product of *P. acidilactici* had an optimum pH of 4.0. The research results by Oke and Onilude¹¹ using the protease from *Lactobacillus plantarum* had an optimum pH of 7, which was in agreement with the research results by Sulthoniyah *et al.*¹². Ionization can also be experienced by the substrate or substrate-enzyme complex, which also affects enzyme activity²¹.

Effect of temperature: Temperature is a vital factor affecting protease activity. According to Baehaki and Budiman²², in general, each enzyme has a maximum activity at a certain temperature, meaning that the enzyme activity will increase as the temperature increases until the optimum temperature is reached. The increase in temperature affects the substrate conformation change so that the substrate's active side encounters barriers while entering the enzyme's active site, causing a decrease in enzyme activity. In Fig. 5, the optimum activity at 40°C was 29.19 U mL⁻¹ and the enzyme activity decreased as the temperature increased. At a temperature of 30°C, there was an increase in protease activity. At 50-90°C, a decrease in protease activation was observed. The stability of an enzyme depends on several things, namely, hydrogen bonding, hydrophobic bonding, ionic interactions and disulfide bridges. The stability of the enzyme in response to temperature can be maintained when the protein's amino acids form a certain conformation that is resistant to the effects of denaturation; in general, proteins are denatured at high temperatures²³. Compared with the *Lactobacillus plantarum* protease, the optimum temperature is 47°C with an activity of 9.76 U mL⁻¹; the research results by Sulthoniyah *et al.*¹² on the optimum activity of the proteases of *Streptococcus lactis* and *Lactococcus lactis* revealed an optimum temperature of 28°C, in agreement with the research results by Akinkugbe and Onilude²⁰.

Effect of metal ions: The results showed that the metal ions that act as activators are Mg²⁺ and Mn²⁺ ions at a concentration of 5 mM with an activity of 23.16 and




Fig. 5: Effect of temperature on the protease activity of lactic acid bacteria BAF-715




Fig. 6: Metal ion effect on the protease activity of lactic acid bacteria BAF-715

39.39 U mL⁻¹, respectively. While with Cu²⁺, Ca²⁺, Fe²⁺ and Zn²⁺, the protease activity values were 11.54, 10.03, 12.19 and 12.21 U mL⁻¹, respectively, showing a reduction compared with the activity of the treatment without adding metal ions in which the activity was only 12.67 U mL⁻¹, as shown in Fig. 6. Cofactor bonds or enzyme inhibitors may alter the enzyme's ability to bind to the substrate and alter the enzyme's catalytic capacity because the enzyme structure in the presence of an inhibitor or cofactor undergoes physical and chemical changes such that the activity changes²⁴. The research results by Xu *et al.*¹⁰ demonstrated that *P. pentosaceus* had 220 proteases, the metal activator was the Ca²⁺ ion, the inhibitors were Zn²⁺, Mg²⁺ and Fe³⁺ and the activity was inhibited by the metal-chelating EDTA and sodium dodecyl sulfate. Regarding the protease of *Lactobacillus plantarum*, the metal activator was Mg²⁺ with an enzyme activity of 7.5 UI mg⁻¹, as shown in the research results by Sulthoniyah *et al.*¹². According to Suhartono²⁴, the presence of cofactors or inhibitors may alter the enzyme's ability to bind the substrate and may alter the enzyme's catalytic capacity because the enzyme structure undergoes physical and chemical changes such that the activity changes.




Fig. 7: Phylogenetic trees of isolate BAF-175 based on the 16S rRNA gene with maximum likelihood (ML), neighbor joining (NJ) and maximum parsimony (MP) analysis

The 0.005 scale shows the evolutionary distance at the branch length, whereas the number on the branch shows the bootstrap value 1000 (ML/NJ/MP)

Molecular identification of isolate BAF-715: Genotypic characterization of the LAB isolate was performed based on DNA sequencing of the gene encoding 16S rRNA to determine the genus and species. The DNA encoding the 16S rRNA can be used as a molecular marker for the definition of a species because these molecules exist in every organism and they perform identical functions in all organisms. DNA from isolate BAF-715 was amplified using 63F and 1387R primers. These primers do not form a duplex structure with an enzyme-recognizable 5' edge 5'-3'exonuclease and no nucleotides are cut off at the 5' end to affect the primary annealing temperature²⁵.

The DNA sequence analysis of the 16S rRNA BAF-715 isolate obtained sequences with a length of 1200 bp. The sequencing of the 16S rRNA BAF715 isolates compared with sequences in the GenBank NCBI database using the BLAST algorithm. The BLAST result for isolate BAF-715 showed 99% homology/similarity to *Pediococcus pentosaceus* (KP 18922.8.1) (Fig. 7). The classification of isolate BAF-715 is as follows: Kingdom: Bacteria; Division: Firmicutes; Class: Bacilli; Order: Lactobacillales; Family: Lactobacillaceae; Genus: *Pediococcus*; Species: *Pediococcus pentosaceus*.

CONCLUSION

Isolate BAF-715 has a higher protease activity than that of isolate BAF-514 and BAF-1121 (18.48, 11.32 and 9.65 U mL⁻¹, respectively). Isolate BAF-715 shows the highest proteolytic index and activity, with protease activity at an incubation time of 40 h, 2.5% casein substrate, 40 °C, pH 7, metal ions Mg²⁺ and Mn²⁺ (5 Mm) as activators and metal ions Fe²⁺, Cu²⁺, Ca²⁺ and Zn²⁺ as inhibitors. The result of the molecular identification with 16S rRNA sequencing and phylogenetic analysis based on the Neighbor Joining method for isolate BAF-715 reveals

that it is *Pediococcus pentosaceus*. A protease produced by *Pediococcus pentosaceus* from Bekasam showed a high proteolytic activity, making it the best protease for application as a beef tenderizer.

SIGNIFICANCE STATEMENTS

This study discovers the optimum conditions for the production of the protease enzyme by *Pediococcus pentosaceus* bacteria; the findings can be beneficial for the food industry to improve meat tenderizers. This study will help researchers uncover the ability of *Pediococcus pentosaceus* to produce a protease enzyme that can be explored by other researchers. Thus, a new theory on the optimum conditions for *Pediococcus pentosaceus* to produce the protease enzyme may be developed.

ACKNOWLEDGMENTS

This study was funded and supported by the Directorate General of Higher Education of the Ministry of Research, Technology and Higher Education through the Hibah Doktor scheme 2017, No: 48/UN21.17/PP/2017. We are very grateful to the Minister of Research, Technology and Higher Education for allowing us to conduct this study.

REFERENCES

- Desniar, I.R., A. Suwanto and N.R. Mubarik, 2012. Antimicrobial compounds produced by lactic acid bacteria from *bekasam*. *J. Akuatika*, 3: 135-145.
- Rao, M.B., A.M. Tanksale, M.S. Ghatge and V.V. Deshpande, 1998. Molecular and biotechnological aspects of microbial proteases. *Microbiol. Mol. Biol. Rev.*, 62: 597-635.

3. Takagi, H., M. Kondou, T. Hisatsuka, S. Nakamori, Y.C. Tsai and M. Yamasaki, 1992. Effects of an alkaline elastase from an alkalophilic *Bacillus* strain on the tenderization of beef meat. *J. Agric. Food Chem.*, 40: 2364-2368.
4. Wilson S.A., O.A Young, T. Coolbear and R.M. Daniel, 1992. The use of protease from extreme thermophilic for meat tenderization. *Meat Sci.*, 32: 93-103.
5. Oyeleke, S.B., E.C. Egwim and S.H. Auta, 2010. Screening of *Aspergillus flavus* and *Aspergillus fumigatus* strains for extracellular protease enzyme production. *J. Microbiol. Antimicrob.*, 2: 83-87.
6. Sandhya, C., A. Sumantha, G. Szakacs and A. Pandey, 2005. Comparative evaluation of neutral protease production by *Aspergillus oryzae* in submerged and solid-state fermentation. *Biochemistry*, 40: 2689-2694.
7. Kumar, D. and T.C. Bhalla, 2004. Purification and characterization of a small size protease from *Bacillus* sp. APR-4. *Indian J. Exp. Biol.*, 42: 515-521.
8. Yusmarini, R. Indrati and Y. Marsono, 2010. Proteolytic activity of lactic acid bacteria in soy milk fermentation. *J. Technol. Food Ind.*, 21: 129-134.
9. Wikandari, P.R., Suparmo, M. Yustinus and S.R. Endang, 2012. Characterization of proteolytic lactic acid bacteria in the Bekasam. *J. Natur Indonesia*, 14: 120-125.
10. Xu, Y., M. Dai, J. Zang, Q. Jiang and W. Xia, 2015. Purification and characterization of an extracellular acidic protease of *Pediococcus pentosaceus* isolated from fermented fish. *Food Sci. Technol. Res.*, 21: 739-744.
11. Oke, M.A. and A.A. Onilude, 2014. Partial purification and characterization of extracellular protease from *Pediococcus acidilactici*. *Niger. J. Basic Applied Sci.*, 22: 19-25.
12. Sulthoniyah, S.T.M., Hardoko and H. Nursyam, 2015. Characterization of extracellular protease lactic acid bacteria from shrimp paste. *J. Life Sci. Biomed.*, 5: 1-5.
13. El-Safey, E.M. and A.U.M. Raouf, 2004. Production, purification and characterization of protease enzyme from *Bacillus subtilis*. Proceeding International Conference for Development and the Environment in the Arab World, March 23-25, 2004, Assiut, Egypt, pp: 23-25.
14. Bergmeyer, H.U., J. Bergmeyer and M. Graal, 1983. *Methods of Enzymatic Analysis*. Vol. 2, Weinheim Deer Field Beach, Florida.
15. Suharsono and U. Widyastuti, 2008. *Practicum guide. Introduction to molecular genetics*. Department of Biology-FMIPA, Bogor Agricultural Institute, Bogor.
16. Tamura, K., M. Nei and S. Kumar, 2004. Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc. Natl. Acad. Sci. USA.*, 101: 11030-11035.
17. Tamura, K., G. Stecher, D. Peterson, A. Filipski and S. Kumar, 2013. MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.*, 30: 2725-2729.
18. Nurmalinga, A., Periadnadi and Nurmia, 2013. Isolation and partial characterization of indigenous bacteria of fermentation of durian fruit (*Durio zibethinus*). *J. Biol. Univ. Andalas*, 2: 8-13.
19. Putranto, W.S., 2006. Purification and characterization of proteases produced by *L. acidophilus* in milk dairy fermentation. Proceedings of the National Seminar on Biotechnology Capturing Opportunities Through Biotechnology, (BCOTB'06), Center for Biotechnology Research, LIPI.
20. Akinkugbe, A.O. and A.A. Onilude, 2013. Selective comparability and physiological studies of lactic acid bacteria protease and *Calotropis procera* (Linn) extracts. *Peer J. Pre*, Vol. 1.
21. Muchtadi, D., N.S Palupi and M. Astawan, 1996. *Enzymes in the food industry*. Inter-University Center for Food and Nutrition, Agricultural Institute, Bogor.
22. Baehaki, A. and R.A. Budiman, 2011. Isolation and characterization of proteases from Indralaya marshland bacteria. *J. Teknol. Food Ind.*, 21: 37-42.
23. Seftiono, H., 2017. Penentuan aktivitas enzim mananase dari berbagai mikroorganisme di Indonesia dan Peranannya Dalam Bidang Pangan: Kajian Pustaka. *Agrointek*, 11: 14-20.
24. Suhartono, M.T., 1989. *Enzymes and biotechnology*. Ministry of Education and Culture of Directorate General of Higher Education between Biotechnology University, Bogor Agricultural Institute, Bogor.
25. Marchesi, J.R., T. Sato, A.J. Weightman, T.A. Martin, J.C. Fry, S.J. Hiom and W.G. Wade, 1998. Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding for bacterial 16S rRNA. *Applied Environ. Microbiol.*, 64: 795-799.

Isolation and Characterization of Lactic Acid Bacteria Proteases from Bekasam for use as a Beef Tenderizer

ORIGINALITY REPORT

6%

SIMILARITY INDEX

%

INTERNET SOURCES

6%

PUBLICATIONS

3%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

4%

★ Zurmiati ., Wizna ., M. Hafil Abbas, Maria Endo Mahata. "Production of Extracellular β -mannanase by *Bacillus amyloliquefaciens* on a Coconut Waste Substrate", Pakistan Journal of Nutrition, 2017

Publication

Exclude quotes On

Exclude bibliography On

Exclude matches < 3%