

Turnitin Originality Report

Processed on: 24-Jul-2020 10:42 PM +08

ID: 1361605643

Word Count: 3337

Submitted: 1

LLRD 2020_Gita Ciptaan By
Gita Ciptaan

6% match
(publications)

[Mirawati., Gita Ciptaan, Ade Djulardi. "The Combined Effects of Fungi Phanerochaete chrysosporium and Neurospora crassa and Fermentation](#)

[Time to Improve the Quality and Nutrient Content of Palm Oil Sludge", Pakistan Journal of Nutrition, 2019](#)

Similarity Index	Similarity by Source	
14%	Internet Sources:	N/A
	Publications:	14%
	Student Papers:	0%

4% match (publications)

[Mirawati., A. Djulardi, G. Ciptaan. "Effect of Fermented Palm Oil Sludge with Neurospora crassa Added to Rations on Broiler Production Performance", Pakistan Journal of Nutrition, 2018](#)

3% match (publications)

[Mirawati., Ade Djulardi, Gita Ciptaan. "Utilization of Fermented Palm Kernel Cake with Sclerotium rolfsii in Broiler Ration", International Journal of Poultry Science, 2018](#)

Livestock Research for Rural Development 32 (8) 2020 LRRD Mission Guide for preparation of papers LRRD Newsletter Citation of this paper Performance of quails fed palm oil sludge fermented with Phanerochaete chrysosporium and Neurospora crassa Gita Ciptaan, Mirawati and Ade Djulardi Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, University of Andalas, Padang 25163, Indonesia gciptaan@ansci.unand.ac.id Abstract This study aims to determine what percentage of the use of fermented palm oil sludge (FPOS) with Phanerochaete chrysosporium and Neurospora crassa in the ration on the performance and quality of quail eggs. This study used 200 quails ages 8 weeks, 20 units of 45x20x30 cm battery cages, each unit occupied by 10 quails, feeding and drinking containers. This study uses a completely randomized design experimental method with 5 treatment rations and 4 replications. The treatment rations using 0, 15, 20, 25 and 30% FPOS, respectively. The rations were arranged based on 20% isoprotein and 2800 kcal/kg isocaloric. The variables were observed: feed consumption, egg production (quail day production), egg mass, feed conversion, egg weight, eggshell thickness and yellow egg color. The results of the variance analysis showed that the use of FPOS with Neurospora crassa and Phanerochaete chrysosporium gave no significant effect ($P>0.05$) on the feed consumption, egg weight, eggshell thickness and egg yolk color but were significantly effected ($P<0.05$) on egg production, egg mass and feed conversion. Based on the results of the study it can be concluded that the optimal used of fermented palm oil sludge with Phanerochaete

chryso sporium and [Neurospora crassa](#) is 25% [in](#) laying quail rations. Keywords: Neurospora crassa, Phanerochaete chryso sporium Introduction One of the animal that have great potential to be developed is quail. Quail breeding does not require large tracts of land because of its small body, quail has the ability to grow and multiply quickly unlike other birds. From the production, cycle quail are able to lay eggs at the age of 35-42 days and can reach 200-300 eggs per year. In animal husbandry business, feed is a big enough cost, reaching 70-80% of production costs, the high cost of feed is due to some of the ingredients of feed ingredients are still imported such as corn, soybean meal and fish meal. From this problem, we need to find alternative feed ingredients that can reduce feed costs. One that can be used is palm oil sludge (POS), palm oil sludge (POS) is waste generated in the extraction and extortion process of palm fruit taken from Crude Palm Oil (CPO). Palm oil sludge is a potential plantation waste, the price is cheap, does not compete with human needs and does not interfere with animal health. Based on data released by the Directorate General of Plantations (2016) [the area of oil palm plants in Indonesia is](#) around 11,914,499 hectares, with a production of 33,229,381 tons. [Each hectare of oil palm can produce](#) 4 tons of oil per year, which is obtained from around 16 tons of fresh fruit bunches (FFB). Every ton of bunches of fresh palm fruit can produce 250 kg of palm oil, 294 kg of palm oil sludge, 35 kg of oil palm flowers and 180 kg of oil palm juice (Mathius et al 2003). Palm oil sludge has high nutritional content such as: [11.1% crude protein, 17% crude fiber, 12% crude lipid](#) (Noferdiman [and](#) Yani 2013), 7.27% hemicellulose and 14.21% lignin (Noferdiman 2004). Mirnawati et al (2017) states that palm oil sludge contains 90.47% dry matter, 13% crude protein, 12.31% crude fat and 32.07% crude fiber with a metabolic energy of 1105.87 kcal / kg. Although the nutrient content of palm oil sludge is quite good as a feed ingredient, it is constrained because crude fiber is so high. This is certainly a problem if it is given to poultry because poultry cannot hydrolyze crude fiber. Sinurat et al (2000) stated that palm oil sludge could only be used 5% in poultry rations. Efforts to [reduce the content of crude fiber and increase the use of palm](#) oil sludge in poultry feed need to be processed microbially through fermentation. Fermentation is processing through the process of decomposition or overhaul of food substances from complex forms into simple food substances with the help of enzymes produced by microbes (Mirnawati et al 2019a). Where microbes can degrade lignin and crude fiber so that it is easier to digest and add flavor and aroma and result in an increase the quality and nutritional content of feed ingredients (Mirnawati et al 2019b). Ligninase is a lignin degrading enzyme produced by microorganisms that are lignolytic. Where lignocellulose bonds can be broken by ligninases such as lignin peroxidase (LiP), manganese peroxidase (MnP) and laccase (Takano et al 2004). LiP and MnP enzymes are produced by several microorganisms, one of which is Phanerochaete chryso sporium. Howard et al (2003) states that Phanerochaete chryso sporium fungi can produce high ligninase and cellulase where Phanerochaete chryso sporium can degrade lignin and its derivatives effectively by producing extracellular peroxidase enzymes in the form of lignin peroxidase and manganese peroxidase. Fermentation with Phanerochaete chryso sporium in a solid substrate allows a change in components of material [that are difficult to digest to be easily digested](#) such as cellulose and hemicellulose into simple sugars thereby increasing the nutritional value of the product. Noferdiman (2009) fermented palm oil sludge (POS) with 6% Phanerochaete chryso sporium inoculum for 8 days can reduce crude fiber from 20.16% to 12.22%, lignin from 14.21% to 8.94% and increase in protein roughly from 10.57% to 14.10% but its utilization in broiler rations is only 15%. [Mirnawati et al](#) (2015) [conducted a study](#) by selecting [three types](#) of Neurospora [crassa, Neurospora sitophila and Neurospora sp](#) molds grown on palm oil sludge (POS). [The best results in fermented palm oil sludge](#) are Neurospora crassa [compared to](#) the [other](#) two [types of](#) mold, where there is an increase in crude

protein from 13% to 20.42%, reducing crude fiber from 32.07% to 23.02%, and metabolic energy from 1105 kcal/kg to 2024 kcal/kg. But its utilization in broiler rations is only 22% (Mirnawati et al 2018). To increase [the nutrient content and](#) the use of [palm oil sludge](#) in poultry rations, the two molds are combined. Mirnawati et al (2019c) have fermented [palm oil sludge by combining Phanerochaete chrysosporium and Neurospora crassa with](#) a ratio (4: 1) fermented for 13 days and giving the best results crude protein 26.20%, crude fiber 14.49%, Lignin 14.54%, Caratenoid 2020 g/100g, Nitrogen retention 58.20%, Crude fiber digestibility 57.66%. From the data above there has been an increase in the nutritional content of fermented POS even this FPOS can be used 22% in broiler rations (Mirnawati et al 2018). Based on the description, it is necessary to do research to find out what percentage (%) of [fermented palm oil sludge with](#) Phanerochaete chrysosporium and [Neurospora crassa can be used](#) in quail rations, it is hoped that FPOS can match the quail control ration so as to provide a favorable performance. Methodology Purpose of the experiment This experiment is purposed [to study the effect of use several levels of fermented palm oil sludge](#) (FPOS) with Phanerochaete chrysosporium and Neurospora crassa in the ration on the performance and egg quality of laying quails (feed consumption (g/head/day), egg production (quail day production/%), egg mass (g/head/day), feed conversion, egg weight (g/grain), eggshell thickness (mm) and egg yolk colour). [Experimental animal and diet composition Two hundred](#) laying quails (Coturnix-coturnix japonica) aged 8 weeks with an average production of 35%. [The experiment was performed in a completely randomized design \(CRD\) with five treatments](#) in the ration (0%, 10%, 15%, 20%, and 25% FPOS) [and four replications. There were](#) ten quails [per unit](#) of [experiment. These](#) laying quails [were kept in](#) battery cages (45 x 20 x 30cm). [The diets were formulated in iso protein 20% and iso caloric 2800 kcal/kg ration](#) (Djulardi 1995). Feed ingredients used was consisted of yellow maize, soybean meal, rice bran, fermented palm oil sludge (FPOS), fish meal, coconut oil, bone flour, mineral B12 and top mix. [Diet and drinking water were provided ad-libitum.](#) The composition of the feed ingredients of treatment ration is shown in Table 1 and the nutrient of the treatment ration is shown in Table 2. The [procedure of preparing](#) FP OS [Fermented palm](#) oil sludge [was the product of 80% POS plus 20% rice bran that was fermented with](#) Phanerochaete chrysosporium and Neurospora crassa as much as 10% with the ratio of each inoculum which is 4: 1. Subsequently incubated in an incubator for 13 days. Furthermore, the FPOS is harvested and roasted at a temperature of 600C to dry. After dry, FPOS is ground and ready to be given in quail ration. Based on the results of Mirnawati et al (2019c), found that the nutritional content of FPOS with combination of Phanerochaete chrysosporium and Neurospora crassa can be seen in table 3. Data collection Data measured are feed consumption, egg production (quail day production), egg mass, feed conversion, egg weight, eggshell thickness and quail egg color Data analysis Datas obtained were processed statistically by analysis of variance of a completely randomized design, followed by DMRT test for the difference between treatments, according to Steel and Torrie (1995). Table 1. Composition of diets Feed Ingredients FPOS in diet (%) 0 15 20 25 30 Yellow Maize Soybean Meal Rice Bran Fish meal FPOS Coconut oil Bone Flour Mineral B12 Top Mix Total (%) 50.00 45.55 23.00 17.75 7.25 7.00 16.00 16.00 0.00 15.00 1.00 0.95 1.25 1.25 1.00 1.00 0.50 0.50 100 100 44.00 43.80 14.85 12.00 6.65 5.25 16.00 16.00 20.00 25.00 0.75 0.20 1.25 1.25 1.00 1.00 0.50 0.50 100 100 43.50 9.00 3.75 16.00 30.00 0.00 1.25 1.00 0.50 100 Table 2. The nutrient content (%) and metabolic energy (kcal/kg) of the treatment diet Feed nutrition FPOS in diet (%) 0 15 20 Crude protein 20.03 20.16 20.13 Crude Fat 3.13 3.07 2.86 Crude Fiber 3.53 4.72 5.30 Ca 2.08 2.06 2.04 P Available 0.95 0.98 1.00 ME 2814 2817 2816 25 30 20.12 20.03 2.29 2.07 5.81 6.31 2.03 2.01 1.02 1.04 2816 2837 Table 3. The nutrient content (%) and metabolic energy

(kcal/kg) of fermented [palm oil sludge](#) with Phanerochaete [chryso sporium and Neurospora crassa](#) Ingredient Crude Protein Crude Fiber Crude Fat Ca P ME FPOS 26.20 14.49 2.22 0.28 0.65 2788 Result Mean of feed consumption, egg production (quail day production), egg mass, feed conversion, egg weight, egg shell thickness and egg color of quail that received a mixture [of fermented palm oil sludge with](#) Phanerochaete chryso sporium and [Neurospora crassa](#) can be seen in Table 4. [The](#) results of the diversity analysis showed that the use of FPOS with Phanerochaete [chryso sporium and Neurospora crassa](#) up [to the](#) level of 25% in ration had no significant effect ($P>0.05$) on the feed consumption, egg weight, shell thickness and egg yolk color of laying quail. However, result showed significant effect on egg production, egg mass and feed conversion ($P<0.05$). Table 4. Mean content of feed consumption, egg production (quail day production), egg mass, feed conversion, egg weight, egg shell thickness and quail egg color Parameter FPOS in diet (%) SEM p 0% Consumption(g/head/day) Feed 22.13 Egg Production (%) 60.39a Egg mass(g/head/day) 6.15a Feed Conversion 3.60a Egg 10.18 Weight (g) 15% 22.13 60.31a 6.13a 3.62a 10.17 Shell Thickness(mm) 0.28 0.28 Egg Yolk 6.00 6.25 Note: Different superscripts shows significant effect ($P<0.05$) 20% 25% 22.15 22.14 60.26a 60.21a 6.13a 6.11a 3.63a 3.66a 10.16 10.15 0.27 0.27 6.50 6.75 30% 22,84 0.36 58.98b 0.32 5.98b 0.04 3.82b 0.05 10.14 0.01 0.26 0.003 7.00 0.25 0.05 0.01 0.01 0.01 0.05 0.05 0.05 Discussion Results of the research showed that [palm oil sludge fermented with](#) Phanerochaete chryso sporium and [Neurospora crassa can be used up to](#) 25% in laying quail [rations](#), as seen from feed consumption, egg production, feed conversion, egg weight, eggshell thickness, and egg yolk color from laying quail that are matching with the control ration. [An increase in the use of fermented](#) palm sludge in quail [rations](#) is caused by [the](#) palm sludge having undergone fermentation where the fermented product will increase palatability due to physical changes in the fermentation process such as taste, odor, texture and easier to digest from the original ingredients. This is consistent with the opinion of Mirnawati et al (2019b) that the fermented product can produce a better aroma and taste than the original material that are more preferred by the quail. Hidayat (2007) also found that the fermentation process can provide beneficial physical changes such as aroma, taste, texture, which are better than the origin. Murugesan et al (2005) found that fermented products can produce flavors that are preferred by livestock and have several vitamins (B1, B2, B12) so that livestock is more preferred than the original ingredients. Fermentation can [improve the quality of palm oil sludge](#) (POS) [and](#) increase digestibility of [the](#) ration so that the protein contained in the ration can be utilized for egg production. This is consistent with the opinion of Setiarto et al (2016) that fermentation can increase protein digestibility and amino acid levels, thus with the increase in digestibility of rations and amino acids, egg production will also increase. This is due to the decreasing amount of use of maize, soybean meal and bran and the increasing use of fermented palm oil sludge products so that it still gives the same egg yolk color. FPOS products used in the ration contained 20.25 $\mu\text{g}/100\text{g}$ carotenoids (Mirnawati et al 2018). Carotenoids contained in FPOS products come from Neurospora crassa which functioned to give the color of egg yolk. According to Hausman and Sandman (2000) β -carotene is a carotenoid group compound that is unstable because it is easily oxidized and will turn into xanthofil. The xanthofil pigment will give the yolk a yellow color. Poultry that consume rations containing higher carotenoid pigments (β -carotene and xanthofil) will produce eggs with a high egg yolk score. If the use of fermented palm oil sludge in the ration is increased, egg production, egg mass and feed conversion will decrease. The decrease in this parameter is due to increased of crude fiber ration (6.31%). According to Djulardi (1995) that crude fiber in quail feeds is below 5%. For more details, this parameter reduction can be seen in Figures 1, 2 and 3. Figure

1. Egg production Figure 2. Egg mass Figure 3. Feed conversion

Conclusion From the results of the study, it can be concluded that the optimal use of fermented palm oil sludge with Phanerochaete chrysosporium and Neurospora crassa is 25% in laying quail rations, it can even reduce the use of soybean meal by about 50% in quail rations.

Acknowledgement This study was financially supported by funds provided by the National Strategic Individual from the Directorate General of Higher Education, Ministry of Research, Technology and Higher Education, Republic of Indonesia: 50/SP2H/LT/DRPM/ 2019, fiscal year 2019.

References Direktorat Jendral Perkebunan 2016 Buku Statistik Perkebunan. <http://ditjenbun.pertanian.go.id/?publikasi=buku-publikasi-statistik-2016-2018> Djulardi A 1995 Respon burung puyuh petelur (Coturnix-coturnix japonica) terhadap pemberian ransum dengan berbagai kandungan fosfor imbalanced protein. Disertasi. Program Pasca Sarjana Padjajaran. Bandung. Hausman A and Sandman G 2000 A single five-step desaturase is involved in the carotenoid biosynthesis pathway to beta-carotene and torulene in Neurospora crassa. J. Genet. Biol. 30: 147-53. <https://doi.org/10.1006/fgbi.2000.1212> Hidayat N 2007 Teknologi Pertanian Pangan. <http://pikiranrakyat./indek.html>. (diakses tanggal 14 Januari 2018) Howard R T E, Abotsi E L, Jansen V R and Howard S 2003 Linicellulose biotechnology: issue of bioconversion and enzyme production. African Journal of Biotech. (2); 602-619. <http://dx.doi.org/10.5897/AJB2003.000-1115> Mathius W, Sitompul D, Manurung B P and Asmi W 2003 Produk sampingan tanaman dan pengolahan buah kelapa sawit sebagai bahan dasar pakan komplit; suatu tinjauan. Prosiding. Loka Karya Nasional: Sistem Integrasi Kelapa Sawit-sapi. Bengkulu, 9-10 September 2003. P. 120-128. <https://docplayer.info/47747315-> Mirnawati, Djulardi A and Ciptaan G 2015 Peningkatan kualitas bungkil inti sawit dan lumpur sawit melalui aplikasi bioteknologi sebagai bahan pakan unggas rendah kolesterol. Laporan Penelitian Unggulan Universitas Andalas, Padang. Kontrak No.30/SP2H/PL/DIT.LITABMAS/II/2016. Universitas Andalas. Mirnawati, Djulardi A and Ciptaan G 2017 Role of humic acid in improving the nutrient content and quality of fermented palm oil sludge. Pakistan Journal of Nutrition. 16 (7): 538-543. DOI: 10.3923/pjn.2017.538.543. Mirnawati, Djulardi A and Ciptaan G 2018 Effect of fermented palm oil sludge with Neurospora crassa added to rations on broiler production performance. Pakistan Journal of Nutrition. 17 (10): 487-491. DOI: 10.3923/pjn.2018.487.491. Mirnawati, Ciptaan G and Ferawati 2019a The effect of Bacillus subtilis inoculum doses and fermentation time on enzyme activity of fermented palm kernel cake. J. World Poult. Res. 9 (4): 211-216. DOI: https://dx.doi.org/10.36380/jwpr.2019.26 Mirnawati, Ciptaan G and Ferawati 2019b Improving the quality and nutrient content of palm kernel cake through fermentation with Bacillus subtilis. Livestock Research for Rural Development. 31 (7): 2019. www.lrrd.org/lrrd31/7/mirna31098.html Mirnawati, Ciptaan G and Djulardi A 2019c The Combined effects of fungi Phanerochaete chrysosporium and Neurospora crassa and fermentation time to improve the quality and nutrient content of palm oil sludge. Pakistan Journal of Nutrition. 18 (5): 437-442. DOI: 10.3923/pjn.2019.437.442. Murugesan G S, Sathiskumar M and Swarnnathan K 2005 Supplementation of waste tea fungal biomass as a dietary ingredient for broiler chicken. Bioresource Technology 96:1743-1748. <https://doi.org/10.1016/j.biortech.2005.01.006> Noferdiman 2004 Uji coba limbah sawit dalam ransum ayam broiler. Majalah Ilmiah Angsana. 08(1); 17-26. Noferdiman 2009 Pengaruh penggunaan lumpur sawit fermentasi dengan jamur P. chrysosporium dalam ransum terhadap performans ayam broiler. Jurnal Ilmiah Ilmu-Ilmu Peternakan. 12 (4): 176-185 <https://online-journal.unja.ac.id/jiip/article/view/166/148> Noferdiman and Yani A 2013 Nutrition content of fermented palm oil sludge with Phanerochaete chrysosporium. Agripet. 13: 47-52.

<http://www.jurnal.unsyiah.ac.id/agripet/article/download/820/758> Setiarto R H B, Widhyastuti N and I Saskiawan 2016 Pengaruh fermentasi fungi bakteri asam laktat khamir terhadap kualitas nutrisi tepung sorgum. Agritech. 36(4). <https://doi.org/10.22146/agritech.16769> [Sinurat A P,](#) Purwadaria [I,](#) Ketaren [P P,](#) Zainuddin [D and](#) Kompiang [I P](#) 2000 Pemanfaatan lumpur sawit untuk ransum unggas: 1. Lumpur sawit kering dan produk fermentasinya sebagai bahan pakan ayam pedaging. JITV 5: 107-112. <https://docplayer.info/51111708> [Steel R G D and](#) Torrie [J H](#) 1995 [Prinsip Prosedur](#) Statistika. Penerjemah Bambang Sumantri. Gramedia Pustaka, Jakarta Takano M, Nakamura M, Nishida A and Ishihara M 2004 Manganase peroxidase from Phanerochaete crassa. WD1694. Bull. <https://www.ffpri.affrc.go.jp/pubs/bulletin/351/documents/390-2.pdf>
Received 10 March 2020; Accepted 25 June 2020; Published 1 August 2020 Go to top