

Jurnal 3 132

by Harnentis1 Harnentis1

Submission date: 13-Nov-2022 10:59PM (UTC+0800)

Submission ID: 1952411105

File name: Jurnal 3.132.Polish Potential probiotic, 2021.pdf (3.14M)

Word count: 169

Character count: 980

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/351570060>

Potential probiotic yeast isolated from an Indonesian indigenous fermented fish (Ikan Budu)

Article in *Potravinarstvo* · May 2021

DOI: 10.5219/1544

CITATIONS

3

READS

247

7 authors, including:



Yeti Marlida
Universitas Andalas

72 PUBLICATIONS 469 CITATIONS

[SEE PROFILE](#)



Nurul Huda
Universiti Malaysia Sabah (UMS)

274 PUBLICATIONS 3,817 CITATIONS

[SEE PROFILE](#)



Frederick Adzitey
University for Development Studies

124 PUBLICATIONS 1,721 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Assessment of the Presence of Selected Heavy Metals and their Concentration Levels in Fresh and Grilled Beef/Guinea Fowl Meat in the Tamale Metropolis, Ghana [View project](#)



Prevalence and antibiotic susceptibility of *Salmonella enterica* isolated from cow milk, milk products and hands of sellers in the Tamale Metropolis of Ghana [View project](#)



POTENTIAL PROBIOTIC YEAST ISOLATED FROM AN INDONESIAN INDIGENOUS FERMENTED FISH (IKAN BUDU)

Yetti Marlida, Nurul Huda, Harnentis, Yuliaty Shafan Nur, Nuri Mekar Lestari,
Frederick Adzitey, Mohd Rosni Sulaiman

ABSTRACT

Budu is a fermented food resulting from the activities of microorganisms like lactic acid bacteria and yeast. Budu, therefore, serves as a source of probiotics that can have beneficial effects on livestock and humans. Nonetheless, their selection has to be done with caution. The current study purposed to find out whether budu has desirable probiotic properties. This was done by determining its pH, bile acid tolerance, hydrophobicity, and inhibition of pathogens such as *Staphylococcus aureus*, *Salmonella enteritidis*, and *Escherichia coli*. An *in vitro* experiment was conducted using three *Saccharomyces cerevisiae* (coded as SC 11, SC 12, and SC 21) in the preparation of budu. The whole experiment was repeated four times. The budus were tested for their probiotic properties (low pH, bile salts, hydrophobicity, and inhibition of pathogenic bacteria). The results showed that the three *Saccharomyces cerevisiae* survived in gastric juice and bile acid, exhibited good hydrophobicity, and could inhibit pathogenic bacteria, both gram-positive and negative pathogens. They were able to survive at pH 2 for 3 h (40.70 to 55.1%), at pH 2 for 5 h (35.25 to 46.88%), in 0.3% bile acid incubated for 3 h (69.69 to 86.56%), in 0.3% bile acid incubated for 5 h (82.22 to 88.18%) and hydrophobicity ability of 97.0 to 98.1%. The inhibition activity against pathogenic bacteria, that is, *Escherichia coli* was 2.50 to 3.81 mm, *Staphylococcus aureus* was 1.66 to 3.71 mm, and *Salmonella enteritidis* was 1.20 to 2.64 mm.

Keywords: probiotic properties; inhibition of pathogenic bacteria; Indonesian fermented fish; Ikan Budu; poultry infection

INTRODUCTION

In recent times, the use of antibiotic growth promoters (AGP) in livestock business or broiler business, in particular, has been banned. This is due to consumer concerns about the presence of AGP residues in products such as meat, milk, and eggs, because of the potential risk of drug resistance they pose to humans. Farmers always try to find a substitute for AGP with organic compounds such as plant extracts, prebiotics in the form of MOS (mannan oligosaccharides) and FOS (fructooligosaccharides), and probiotics such as giving live microorganisms to livestock (Davari et al., 2019). Microorganisms of the lactobacillus genera are mostly used for the commercial production of probiotics, especially in fermented milk worldwide (Sharif et al., 2017). Probiotics promote the growth of healthy microflora in the gastrointestinal tract (Rajoka et al., 2018).

Probiotics can be from bacteria, fungi, and yeast. Bacteria that are widely used as probiotics are lactic acid bacteria, from fungi are *Rhizopus oligosporus*, while from yeast are *Saccharomyces cerevisiae* and *Saccharomyces boulardii*. The source of microorganisms used as probiotics is essential, which is usually obtained from the digestive tract of livestock because they are already adapted to the

intestine. There are not many reports showing that probiotics are isolated from fermented foods such as budu.

Traditional fermented fish, also known as budu is produced mainly in West Sumatra, Indonesia. Budu is usually made from leather skin (*Chorinemus* spp.) and Spanish mackerel (*Scomberomorus* spp.) known as Ikan Talangand Ikan Tenggiri, respectively, in the Indonesian language (Huda, 2012). Budu processing starts with the hanging of fresh fish by the tail fin for four hours under room temperature. It is then eviscerated, washed, and covered with a layer of salt in a traditional container. After which, it is stored for one day at room temperature and sun-dried for five days. Garlic and white pepper can be sprinkled on the fish during the drying process to improve the quality of the budu (Huda and Ahmad, 2006).

Anggraini et al. (2019) isolated lactic acid bacteria from budu and found that the LAB produced Gamma-aminobutyric acid (GABA), which served as anti-heat stress for broilers. They also found that yeast undergoes symbiosis with lactic acid bacteria in budu. Stadie et al. (2013) reported a symbiotic relationship between yeast and lactic acid bacteria of water kefir origin. Symbiosis

(commensalism or mutualism) widely occurs in fermented foods such as sourdough, milk kefir, and yogurt.

As a probiotic, yeast must be able to withstand gastric pH, bile acids, and pathogenic bacteria. **Brandão et al. (2014)** found that acidic pH was not affected by the fatty acid composition of *S. boulardii*. Yeast is capable of maintaining its internal pH by consuming H⁺ through a metabolic pathway and by using cell buffer systems. **Ogunremi et al. (2015)** added that *Pichia kudriavzevii* ROM 11, that is, yeast from Ogi, which is a cereal pudding fermented food from Nigeria usually made from corn, sorghum, or millet, had a resistance of 86.36% against bile acids with a concentration of 0.3%.

The purpose of the research was to find yeast present in budu that can serve as a probiotic for potential application for poultry production.

Scientific hypothesis

As the yeast-derived from fermented foods from West Sumatra such as fermented fish (budu) has not been exploited, especially as a candidate for probiotics, the study hypothesizes that probiotic yeast exists in fermented fish Ikan Budu.

MATERIAL AND METHODOLOGY

Samples

A sample of budu was purchased from a traditional producer at Pariaman Regency, West Sumatra, Indonesia. Budu was made from coral reef fish such as red Kakap (*L. campechanus*) and Tenggiri fish (*Scomberomorini*) as shown in Figure 1.

Chemicals

Chemicals used in this study were NaCl (Merck, Germany), glycerol (Merck, Germany), HCl (Merck, Germany), oxgall (synthetic bile salts) (Merck, Germany), phosphate buffer (Merck, Germany), and lactic acid (Merck, Germany). The media used in this study were yeast universal agar, nutrient broth, and nutrient agar. All media used were also purchased from Merck, Germany.

Biological Material

Biological material involved in this study was isolated of *Salmonella enteritidis*, *Staphylococcus aureus*, *Saccharomyces cerevisiae*, and *Escherichia coli*.

Equipment

The equipment used in this study were microscope (Merck, Germany) spectrophotometer (Mettler Toledo UV Vis, Ingggris), incubator (Thomas Scientific, USA),

autoclaved (Systec autoclaved company, Japan), and caliper (Misumi, Indonesia).

Description of Experiments

Isolation of yeast was conducted according to the method of **Bajwa and Sharma (2018)**. The budu (1 g) was added to 9 mL of 0.9% NaCl (saline) solution and mixed thoroughly for 60 s. Serial dilution was then carried out in saline solution and spread plated onto yeast universal agar. The yeast universal agar was composed of 3.0 g.L⁻¹ malt extract, 3.0 g.L⁻¹ yeast extract, 10.0 g.L⁻¹ glucose, 5.0 g.L⁻¹ peptone, and 15.0 g.L⁻¹ agar. The spread plated yeast universal agar was incubated for 72 h at 28 °C. Presumptive yeast showed white-to-yellow colonies under the microscope. Such isolates were randomly selected and further purified on yeast universal agar. Yeasts showing the typical appearance of *Saccharomyces* (white-to-yellow colonies) were selected. The selected yeast strains were further purified by successive streaking on yeast universal media. Three isolates were maintained at -80 °C in 20% (v/v) glycerol (Hi-Media).

pH resistivity test was done using a modified nutrient broth in which 0.1 N HCl was added to achieve a pH of 2 which corresponds with gastric pH as described by **Ogunremi et al. (2015)**. The yeast extract (1 mL) was inoculated in modified HCl nutrient broth and incubated at 37 °C for 3 and 5 h. After which, the absorbance was read at a wavelength of 600 nm. This research was conducted with three replications. Isolates resistance was also expressed as a percentage, according to **Ogunremi et al. (2015)**.

Bile salt resistivity test was conducted using mixed modified HCl nutrient broth with 0.3% oxygall (synthetic bile salts) and incubated for 3 and 5 h (**Ogunremi et al., 2015**). The culture was streaked onto a modified nutrient broth and incubated at 37 °C. 1 mL of yeast isolate was streaked onto the bile salt modified nutrient broth medium. Incubation was done at 37 °C (**Ogunremi et al., 2015**). The results were obtained qualitatively by comparison of the yeast absorbancy of the control (not streaked with yeast) with the streaked modified nutrient broth (0.3% bile salts).

Hydrophobicity test or attachment was carried out by the method of **Vinderola et al. (2004)** using stainless steel plates. Clean and dried stainless-steel plates were marked on one side. One hundred (100) mL of distilled water was used to dissolve 0.8 g of nutrient broth. Growth media and stainless steel were autoclaved at 121 °C for 15 min.



Figure 1 Fermented fish (budu) made from Tenggiri fish (*Scomberomorini*).

The stainless steel plate was placed in 25 mL nutrient broth inoculated with 1 mL of LAB in an Erlenmeyer and incubated for 24 h at 29 °C. Furthermore, the surface of the stainless steel was evenly wiped with a swab. The swab was put in a tube containing 10 mL of phosphate buffer and homogenized. It was then measured by looking at the absorbance at a wavelength of 600 nm (A). To measure the growth in the liquid phase, 1 mL of liquid was taken from nutrient broth media and diluted in 9 mL of phosphate buffer solution. After which, it was measured at a wavelength of 600 nm (Ao). The percentage hydrophobicity was calculated using the formula of Fadda et al. (2017).

The antimicrobial activity test of 3 yeast isolates against *Salmonella enteritidis*, *Staphylococcus aureus*, and *Escherichia coli* was carried out based on a modification from Diosma et al. (2013). Nutrient agar (10 grams) was added to 500 mL of distilled water, homogenized, heated in a water bath, and autoclaved. The media was allowed to cool (± 45 °C), and 0.2% of test bacteria was added into ± 10 mL Petri dishes and allowed to solidify. Meanwhile, a blank antibiotic disk was soaked in the lactic acid bacteria solution for approximately 10 min and was placed on the surface of the nutrient agar medium containing the pathogenic bacterial isolates. It was then incubated aerobically at 37 °C for 24 h. After incubation, the diameter of the inhibition zones was measured using a caliper.

Statistical Analysis

Data were subjected to a one-way analysis of variance (ANOVA), and Tukey's test was used for comparison of means using SPSS version 20.0 Software (SPSS Inc., Chicago, IL, USA). A significant difference was defined at $p < 0.05$.

RESULTS AND DISCUSSION

Resistance test of yeast isolates to gastric pH

The resistance of yeast to gastric pH was tested at pH 2 because the pH in the proventriculus and gizzard is 2.0 – 3.5 (Rougie' re and Carre, 2010). The gastric pH was tested for 3 h and 5 h, the results for which can be seen in Table 1. The results of the study showed that all yeast isolates could survive at pH 2 with resistance $>30\%$. The resistance of the yeast isolates to pH did not differ significantly ($p > 0.05$) from each other. The three yeast isolates of budu origin grew at pH 2 with the viability of 55.1% for isolate SC 11, 43.70% for isolate SC 12, and 40.70% for isolate SC 21, which were incubated for 3 h. When the time incubation time was increased to 5 h, the percentage viability decreased.

The pH of 2.0 – 3.5 is pH in the proventriculus, where HCl is produced. The probiotic yeast work in the gastrointestinal tract (GIT) by providing nutrients that aid in the digestion of food and inhibition of harmful bacteria. Probiotics are also mixed or added to feed to increase the rate of feed and nutrient utilization (Markowiak and Śliżewska, 2018).

The results in Table 1 show that the yeast isolate that had the highest resistance to gastric juice was SC 11, with resistance $>50\%$. Nurnaafi, Setyaningsih and Desniar (2015) explained that good probiotic isolates are those with a survival rate of more than 50% under low pH conditions and are resistant to bile salts. The resistance of isolate SC

11 at 3 h incubation time was 55.10% and decreased at 5 h incubation time to 46.88%, a difference of 8.22%. The results of this study were similar to those of Kumura et al. (2004), who found that yeast (*Kluyveromyces lactis* S25) isolated from commercial blue cheese and kefir had a resistance of 54.7%. The results of this study for SC 11 were higher than those of Diosma et al. (2013). They examined yeast isolates from kefir (tested at pH 2.5 with an incubation time of 3 hours) and reported that *Kluyveromyces marxianus* 8116 had 45.5% resistance, *Saccharomyces cerevisiae* 8115 had 40.5% resistance, and *Saccharomyces boulardii* had 45.5% resistance.

Tovar et al. (2002) reported that when yeast isolates enter the digestive tract of poultry, they must be able to survive at low pH because the proventriculus and gizzard have a pH of 2.0 – 4.5. Zubaidy and Khanda (2014) added that *Saccharomyces cerevisiae* var *boulardii* (S.b32) was able to survive at low pH. Glucomannan, chitin, mannoprotein, and beta-glucan make up the cell component of *Candida* sp. Drabikova et al. (2009). However, beta-glucan forms the largest (50 – 60%) component of the inner layer of the cell wall, while chitin forms 1 – 10%. Mannoproteins form mainly 30 – 40% of the outer layer of the cell wall. They play a major role in interactions with the host, determine the nature of the cell surface and cell-to-cell recognition (Vickova et al., 2004).

Resistance test of yeast isolates to bile salts

Resistance of the yeast isolates to bile salt was not significantly different ($p > 0.05$). The results in Table 1 shows that isolate SC 11 had a resistance ability of 69.69% at an incubation time of 3 hours and increased to 82.02% at an incubation time of 5 hours, a difference of 12.33% increase. The results of this study are comparable to those of Chen et al. (2010), who examined yeast isolated from fresh milk on Beijing and Heilongjiang farms against 0.3% bile salts and reported that *Pichia fermentans* HJ15 isolate had 79% resistance, *Pichia kudriavzevii* BY10 isolate had a resistance of 25.9%, and *Yarrowia lipolytica* HJ6 isolate had a resistance of 62.9%.

Yeast can survive in bile salt solutions because of its extreme environmental resistance (Chen et al., 2010). Chen et al. (2010) explained that yeast develops resistance properties in stressful environments (salt, acids, and sugars), and in competition with other microbial yeasts, they can live a normal life. The difference in the results of this study can be due to the differences in the type of yeast tested against 0.3% bile salts. Yeast cell walls are mostly composed of beta-glucan (Lee et al., 2001). Ooi and Liu (2000) reported that beta-glucan is a linear polysaccharide that contains monomers of glucose that are linked by glycosidic bonds. Beta-glucan is water-soluble, and a small concentration will produce high viscosity (Suzuki et al., 2001) and will form a gel in the digestive tract to increase the excretion of bile acids. By this, the fat is not emulsified and absorbed in the stomach.

Inhibition test against pathogenic bacteria

The ability of yeast to act antagonistically is due to changes in medium pH, competition for nutrients, secretion of antimicrobial agents, and production of ethanol in high concentrations. *Staphylococcus aureus*, *Escherichia coli*, and *Salmonella enteritidis* were chosen because they are among the pathogenic bacteria associated with poultry and other animals. The results of the study (Table 2) showed that the inhibitory zone produced by isolate SC 11 against *Escherichia coli* was 3.81 mm, and greater than the inhibition zone of 3.71 mm produced against *Staphylococcus aureus*. However, the inhibition of the yeast isolates to *Staphylococcus aureus*, and *Escherichia coli* did not differ significantly ($p > 0.05$). The differences in the bacterial cell walls could not cause significant differences in their resistance to yeast. *Staphylococcus aureus* is gram-positive bacteria, while *Escherichia coli* is gram-negative bacteria. **Saidi et al. (2019)** reported that gram-negative bacteria have a thinner layer of peptidoglycan (5 – 10 cm), while gram-positive bacteria have a thicker layer of peptidoglycan (20 – 80 cm). Therefore, it was more difficult for yeast isolates to penetrate the cell wall of *Staphylococcus aureus* bacteria than the cell wall of *Escherichia coli* bacteria.

Table 2 shows that the inhibition of *Salmonella enteritidis* by yeast was numerically lower than that of *Staphylococcus aureus* and *Escherichia coli*. This might be influenced by antigens present in *Salmonella*. According to **Wang et al. (2020)**, *Salmonella* has three main antigens, namely: somatic antigens or O antigens, flagellate antigens or

H antigens, and capsule antigens or Vi antigens; which produce enterotoxins and cytotoxins, making it difficult for yeast to inhibit their growth.

The inhibitory strength possessed by yeast varies; therefore, different yeast species will produce different inhibition and metabolite activities during fermentation. **Freimoser et al. (2019)** studied the antagonistic activity of *Kloeckera* and *Kluyveromyces* species against bacteria and found that they produce intracellular and extracellular antimicrobial compounds that inhibit the growth of gram-negative and gram-positive bacteria. Research by **Suzuki et al. (2001)** and **Marquina et al. (2002)** found yeast to produce antagonistic activity due to the production of killer toxins or mycotoxins. These toxins are extracellular proteins or glycoproteins that can damage the cell membrane. The antimicrobial activity of yeast through the secretion of organic acids and antimicrobial peptides has been reported (**Boirivant and Strober, 2007; Vanderpool, Yan and Polk, 2008; Ciorba, 2012**). *S. boulardii* secretes mainly capric acid, a medium-chain fatty acid which showed bioactivity against *Candida albicans* and formation of biofilms (**Krasowska et al., 2009; Murzyn et al., 2010**). *S. cerevisiae* secretes antimicrobial peptides (saccharomycin), which inhibits the growth of pathogenic bacteria (**Hammami et al., 2013**). Antimicrobial peptides inhibit bacteria growth by absorbing the cell membrane receptors, destructing cell membrane permeability and alteration of intracellular pH (**Rizk et al., 2018**).

Table 1 The resistance of yeast isolates towards acid and bile salt conditions.

Isolates yeast	Time (3 h)(%)	Time (5 h)(%)
Acid condition		
<i>Saccharomyces cerevisiae</i> (SC) 11	55.10 ±2.19	46.88 ±1.82
<i>Saccharomyces cerevisiae</i> (SC) 12	43.70 ±1.35	39.36 ±0.80
<i>Saccharomyces cerevisiae</i> (SC) 21	40.70 ±0.87	35.26 ±0.38
Bile salt condition		
<i>Saccharomyces cerevisiae</i> (SC) 11	69.69 ±0.14	82.02 ±0.53
<i>Saccharomyces cerevisiae</i> (SC) 12	84.54 ±1.37	87.43 ±1.91
<i>Saccharomyces cerevisiae</i> (SC) 21	86.56 ±1.71	88.18 ±1.72

Note: values were reported as means ±SD of triplicate groups.

Table 2 The resistance of yeast isolates towards pathogenic bacteria associated with poultry.

Yeast isolates	Diameter inhibition zone (mm)		
	<i>Escherichia coli</i>	<i>Staphylococcus aureus</i>	<i>Salmonella enteritidis</i>
<i>Saccharomyces cerevisiae</i> (SC) 11	3.81	3.71	2.64
<i>Saccharomyces cerevisiae</i> (SC) 12	2.50	2.56	2.52
<i>Saccharomyces cerevisiae</i> (SC) 21	1.63	1.66	1.20

Note: values were reported as means ±SD of triplicate groups; mean values in the same column with different lowercase were significantly different ($p < 0.05$).

Table 3 The hydrophobicity of yeast on stainless steel plates.

Yeast Isolates	Hdrophobicity (%)
<i>Saccharomyces cerevisiae</i> (SC) 11	97.00 ±0.24
<i>Saccharomyces cerevisiae</i> (SC) 12	97.96 ±0.72
<i>Saccharomyces cerevisiae</i> (SC) 21	98.71 ±0.19

Note: values were reported as means ±SD of triplicate groups.

Hydrophobicity using stainless steel plates

Table 3 shows the hydrophobicity of the yeast isolates. From Table 3, all the yeast isolates had a hydrophobicity percentage >90%. The hydrophobicity of yeast isolates was not significantly different ($p > 0.05$).

The results of this study were higher than those of Sourabh et al. (2011), who showed that *Saccharomyces cerevisiae* isolated from traditional West Himalayan fermentation food, Sc01 had a hydrophobicity percentage of 59.65%. Fadda et al. (2017) found that *Saccharomyces boulardii* isolated from codex had a hydrophobicity ability of 55.9%, *Kluyveromyces lactis* isolated from kefir had a hydrophobicity ability of 74.1 – 79.4% and *Kluyveromyces marxianus* had a hydrophobicity ability of 75.9%.

The ability of microorganisms to attach to the digestive tract becomes one of the selection criteria for probiotics. The formation of colonies in the digestive tract is influenced by the ability of microorganisms to attach to the digestive tract that is specific to the host (Suzuki et al., 2001). Tovar et al. (2002) reported that some yeast species can synthesize and secrete polyamine molecules which can stimulate the development of the digestive tract and the production of digestive enzymes.

CONCLUSIONS

The results indicated that all the isolates identified were *Saccharomyces* sp., (SC 11; SC 12 and SC 21) and showed notable potential probiotic properties. They exhibited a better survival in gastric juice and bile acid, showed high hydrophobicity, and the ability to inhibit pathogenic bacteria (gram-positive and negative pathogens) associated with poultry. They were able to live at pH 2 for 3 h (40.70 to 55.1%), pH 2 at 5 h (35.25 to 46.88%), in bile acid 0.3% incubated for 3 h (69.69 to 86.56%), and at 5 h (82.22 to 88.18%), and hydrophobicity ability of 97.0 to 98.1%. The inhibition zones produced by *Saccharomyces* sp. against *Escherichia coli* was 2.50 to 3.81 mm, *Staphylococcus aureus* was 1.66 to 3.71 mm, and *Salmonella enteritidis* was 1.20 to 2.64 mm.

REFERENCES

Anggraini, L., Marlida, Y., Wizna, W., Jamsari, J., Mirzah, M., Adzitey, F., Huda, N. 2019. Molecular identification and phylogenetic analysis of gaba-producing lactic acid bacteria isolated from indigenous dadih of West Sumatera, Indonesia. *F1000 research*, vol. 7, p. 1663. <https://doi.org/10.12688/f1000research.16224.3>

Bajwa, J., Sharma, N. 2018. Evaluation of probiotic properties of yeasts isolated from sidra - an ethnic fermented fish product of North East India. *International Journal of Current Microbiology and Applied Sciences*, vol. 7, no. 2, p. 2632-2643. <https://doi.org/10.20546/ijemas.2018.702.320>

Boirivant, M., Strober, W. 2007. The mechanism of action of probiotics. *Current Opinion in Gastroenterology*, vol. 23, p. 679-692. <https://doi.org/10.1097/MOG.0b013e3282f0cffe>

Brandão, R. L., Rosa, J. C. C., Nicoli, J. R., Almeida, M. V. S., Carmo, A. P., Queiros, H. T., Castro, I. M. 2014. Investigating acid stress response in different *Saccharomyces* strains. *Journal of Mycology*, vol. 2014, p. 1-9. <https://doi.org/10.1155/2014/178274>

Chen, L.-S., Ma, Y., Maubois, J.-L., He, S.-H., Chen, L.-J., Li, H.-M. 2010. Screening for the potential probiotic yeast

strains from raw milk to assimilate cholesterol. *Dairy Science & Technology*, vol. 90, p. 537-548. <https://doi.org/10.1051/dst/2010001>

Ciorba, M. A. 2012. A gastroenterologist's guide to probiotics. *Clinical Gastroenterology and Hepatology*, vol. 10, no. 9, p. 960-968. <https://doi.org/10.1016/j.cgh.2012.03.024>

Davari, D. D., Negahdaripour, M., Karimzadeh, I., Seifan, M., Mohkam, M., Masoumi, S. J., Berenjian, A., Ghasemi, Y. 2019. Prebiotics: Definition, types, sources, mechanisms, and clinical applications. *Foods*, vol. 8, no. 92, p.1-27. <https://doi.org/10.3390/foods8030092>

Diosma, G., Romanin, D. E., Rey-Burusco, M. F., Londero, A., Garrote, G. L. 2013. Yeasts from kefir grains: Isolation, identification, and probiotic characterization. *World Journal of Microbiology and Biotechnology*, vol. 30, p. 43-53. <https://doi.org/10.1007/s11274-013-1419-9>

Drabikova, K., Perecko, T., Nosal, R., Bauerova, K., Ponist, S., Mihalova, D., Kogan, G., Jancinova, V. 2009. Glucmannan reduces neutrophil free radical production in vitro and in rats with adjuvant arthritis. *Pharmacological Research*, vol. 59, no. 6, p. 399-403. <https://doi.org/10.1016/j.phrs.2009.02.003>

Fadda, M. E., Valentina, M., Maura, D., Maira, B. P., Sofia, C. 2017. In vitro screening of *Kluyveromyces* strains isolated from fiore sardo cheese for potential use as probiotics. *Food Science and Technology*, vol. 75, p. 100-106. <https://doi.org/10.1016/j.lwt.2016.08.020>

Freimoser, F. M., Mejia, M. P. R., Tilocca, R., Migheli, Q. 2019. Biocontrol yeasts: Mechanisms and applications. *World Journal of Microbiology and Biotechnology*, vol. 35, p. 154. <https://doi.org/10.1007/s11274-019-2728-4>

Hammami, R., Fernandez, B., Lacroix, C., Fliss, I. 2013. Anti-infective properties of bacteriocins: An update. *Cellular and Molecular Life Sciences*, vol. 70, p. 2947-2967. <https://doi.org/10.1007/s00018-012-1202-3>

Huda, N. 2012. Indonesian Fermented Fish Products. In Hui, Y. H., Özgül Evranuz, E. *Handbook of Animal-Based Fermented Foods and Beverage Technology*. 2nd ed. Boca Raton, US : CRC Press, p. 717-737. ISBN9780429107528.

Huda, N., Ahmad, R. 2006. Budu and tukai: endemic fermented fish products from WestSumatra. *INFOFISH International*, vol. 3, p. 49-51.

Krasowska, A., Murzyn, A., Dyjankiewicz, A., Lukaszewicz, M., Dziadkowiec, D. 2009. The antagonistic effect of *Saccharomyces boulardii* on *Candida albicans* filamentation, adhesion and biofilm formation. *FEMS Yeast Research*, vol. 9, no. 8, p. 1312-1321. <https://doi.org/10.1111/j.1567-1364.2009.00559.x>

Kumura, H., Tanoue, Y., Tsukahara, M., Tanaka, T., and Shimazaki, K. 2004. Screening of dairy yeast strains for probiotic applications. *Journal. Dairy Science*, vol. 87, no. 12, p. 4050-4056. [https://doi.org/10.3168/jds.S0022-0302\(04\)73546-8](https://doi.org/10.3168/jds.S0022-0302(04)73546-8)

Lee, J. N., Lee, D. Y., In-Hye, J., Gi-Eun, K., Kim, H. N. 2001. Purification of soluble β -Glucan with immunoenhancing activity from the cell wall of yeast. *Bioscience Technology and Biochemistry*, vol. 65, no. 4, p. 837-841. <https://doi.org/10.1271/bbb.65.837>

Markowiak, P., Śliżewska, K. 2018. The role of probiotics, prebiotics and synbiotics in animal nutrition. *Gut Pathology*, vol. 10, p. 21. <https://doi.org/10.1186/s13099-018-0250-0>

Marquina, D., Santos, A., Peinado, J. M. 2002. Biology of killer yeasts. *International Microbiology*, vol. 5, p. 65-71. <https://doi.org/10.1007/s10123-002-0066-z>

- Murzyn, A., Krasowska, A., Stefanowicz, P., Dziadkowiec, D., Lukaszewicz, M. 2010. Capric acid secreted by *S. boulardii* inhibits *C. albicans* filamentous growth, adhesion and biofilm formation. *PLoS One*, vol. 5, p. e12050. <https://doi.org/10.1371/journal.pone.0012050>
- Nurnaafi, A., Setyaningsih, I., Desniar. 2015. Potensi probiotik bakteri asam laktat asal bekasam ikan nila (Probiotic potential of lactic acid bacteria from bekasam nila fish). *Jurnal Teknologi dan Industri Pangan*, vol. 26, no. 1, p. 109-114. <https://doi.org/10.6066/jtip.2015.26.1.109> (in Indonesian)
- Ogunremi, O. R., Sanni, A. I., Agarwal, R. 2015. Probiotic potential of yeasts isolated from some cereal-based Nigeria traditional fermented food products. *Journal of Applied Microbiology*, vol. 119, no. 3, p. 797-808. <https://doi.org/10.1111/jam.12875>
- Ooi, V. E., Liu, F. 2000. Immunomodulation and anti-cancer activity of polysaccharide protein complexes. *Current Medicinal Chemistry*, vol. 7, no. 7, p. 715-729. <https://doi.org/10.2174/0929867003374705>
- Rajoka, M. Sh. R., Hayat, H. F., Sarwar, S., Mehwish, H. M., Ahmad, F., Hussain, N., Shah, S. Z. H., Khurshid, M., Siddiqui, M., Shi, J. 2018. Isolation and evaluation of probiotic potential of lactic acid bacteria isolated from poultry intestine. *Microbiology*, vol. 87, no. 1, p. 116-126. <https://doi.org/10.1134/S0026261718010150>
- Rizk, Z., El Rayess, Y., Ghanem, C., Mathieu, F., Taillandier, P., Nehme, N. 2018. Identification of multiple-derived peptides produced by *Saccharomyces cerevisiae* involved in malolactic fermentation inhibition. *FEMS Yeast Research*, vol. 18, no. 7, p. foy080. <https://doi.org/10.1093/femsyr/foy080>
- Rougie' re, N., Carre, B. 2010. Comparison of gastrointestinal transit times between chickens from D1 and D2 genetic lines selected for divergent digestion efficiency. *Animal & The Animal Consortium*, vol. 4, no. 11, p. 1861-1872. <https://doi.org/10.1017/S1751731110001266>
- Saidi, N., Owlia, P., Mahmoud, S., Marashi, A., Sadari, H. 2019. Inhibitory effect of probiotic yeast *Saccharomyces cerevisiae* on biofilm formation and expression of α -hemolysin and enterotoxin A genes of *Staphylococcus aureus*. *Iranian Journal of Microbiology*, vol. 11, no. 3, p. 246-254. <https://doi.org/10.18502/ijm.v11i3.1331>
- Sharif, M., Moridnia, A., Mortazavi, D., Salehi, M., Bagheri, M., Sheikhi, M. 2017. Kefir: a powerful probiotics with anticancer properties. *Medical Oncology*, vol. 34, p. 183. <https://doi.org/10.1007/s12032-017-1044-9>
- Sourabh, A., Kanwar, S. S., Sharma, O. P. 2011. Screening of indigenous yeast isolates obtained from traditional fermented foods of Western Himalayas for probiotic attributes. *Journal of Yeast and Fungal Research*, vol. 2, no. 8, p. 117-126. <https://doi.org/10.5897/JYFR.9000045>
- Stadie, J., Gultiz, A., Ehrmann, M. A., Vogel, R. F. 2013. Metabolic activity and symbiotic interactions of lactic acid bacteria and yeasts isolated from water kefir. *Food Microbiology*, vol. 35, no. 2, p. 92-98. <https://doi.org/10.1016/j.fm.2013.03.009>
- Suzuki, C., Ando, Y., Machida, S. 2001. Interaction of SMKT, a killer toxin produced by *Pichia farinosa*, with the yeast cell membranes. *Yeast*, vol. 18, no. 16, p. 1471-1478. <https://doi.org/10.1002/yea.791>
- Tovar, D., Zambonino, J., Cahu, C., Gatesoupe, F. J., Vazquez-Ju'arez, R., Lesel, R. 2002. Effect of live yeast incorporation in compound diet on digestive enzyme activity in sea bass (*Dicentrarchus labrax*) larvae. *Aquaculture*, vol. 204, no. 1-2, p. 113-123. [https://doi.org/10.1016/S0044-8486\(01\)00650-0](https://doi.org/10.1016/S0044-8486(01)00650-0)
- Vanderpool, C., Yan, F., Polk, D. B. 2008. Mechanisms of probiotic action: Implications for therapeutic applications in inflammatory bowel diseases. *Inflammatory Bowel Diseases*, vol. 14, no. 1, p. 1585-1596. <https://doi.org/10.1002/ibd.20525>
- Vickova, V., Duhova, V., Svidova, S., Farkassova, A., Kamasova, S., Vlcek, D., Kogan, G., Rauko, P., Miadokova, E. 2004. Antigenotoxic potential of glucomannan on four model test systems. *Cell Biology and Toxicology*, vol. 20, no. 6, p. 325-332. <https://doi.org/10.1007/s10565-004-0089-7>
- Vinderola, C. G., Medici, M., Perdigon, G. 2004. Relationship between interaction sites in the gut, hydrophobicity, mucosal immunomodulating capacities and cell wall protein profiles in indigenous and exogenous bacteria. *Journal of Applied Microbiology*, vol. 96, no. 2, p. 230-243. <https://doi.org/10.1046/j.1365-2672.2004.02158.x>
- Wang, M., Qazi, I. H., Wang, L., Zhou, G., Han, H. 2020. Review salmonella virulence and immune escape. *Microorganisms*, vol. 8, no. 407, p. 1-23. <https://doi.org/10.3390/microorganisms8030407>
- Zubaidy, Z. M. A., Khanda, O. K. 2014. Isolation and identification of *Saccharomyces cerevisiae* var *boulardii* and its uses as a probiotic (in vitro). *Rafidain Journal of Science*, vol. 25, no.1, p. 1-11. <https://doi.org/10.33899/rjs.2014.86051>

Funds:

The authors are grateful to the Ministry of Research, Technology and Higher Education of Indonesia for funding the BOPTN Andalas University Grants through Research Cluster Grant No: 57/ UN.16.17/HGB/LPPM/2019. Grant for publication fee was by Universiti Malaysia Sabah, Jalan UMS, 88400, Kota Kinabalu, Sabah, Malaysia.

Acknowledgments:

We would like to thank the University of Andalas, University for Development Studies and Universiti Malaysia Sabah for the support to conduct research in the area of food science and technology.

Conflict of Interest:

The authors declare no conflict of interest.

Contact Address:

*Yetti Marlida, University of Andalas, Faculty of Animal Science, Department of Animal Nutrition and Feed Technology, Padang 25163, Indonesia,

E-mail: yettimarlida@ansci.unand.ac.id

ORCID: <https://orcid.org/0000-0001-9134-3954>

*Nurul Huda, Universiti Malaysia Sabah, Faculty of Food Science and Nutrition, Department of Food Science and Nutrition, 88400, Kota Kinabalu, Sabah, Malaysia; Universitas Sebelas Maret, Faculty of Agriculture, Department of Food Science and Technology, Jln. Ir. Sutami 36 A, Surakarta, Central Java, 57126, Indonesia,

E-mail: drnurulhuda@ums.edu.my

ORCID: <https://orcid.org/0000-0001-9867-6401>

Harmintis, University of Andalas, Faculty of Animal Science, Department of Animal Nutrition and Feed Technology, Padang 25163, Indonesia,

E-mail: harmintis@ansci.unand.ac.id

ORCID: <https://orcid.org/0000-0001-7580-5637>

Yuliaty Shafan Nur, University of Andalas, Faculty of Animal Science, Department of Animal Nutrition and Feed Technology, Padang 25163, Indonesia,

E-mail: yuliaty@ansci.unand.ac.id

ORCID: <https://orcid.org/0000-0002-4689-9213>

Nuri Mekar Lestari, University of Andalas, Faculty of Animal Science, Department of Animal Nutrition and Feed Technology, Padang 25163, Indonesia,

E-mail: nrlhd_usm@yahoo.com.my

ORCID: <https://orcid.org/0000-0002-2452-5640>

Frederick Adzitey, University for Development Studies, Faculty of Agriculture, Department of Food Science, P.O. Box TL 1882, Tamale, Ghana,

E-mail: adzitey@yahoo.co.uk

ORCID: <https://orcid.org/0000-0002-8814-0272>

Mohd Rosni Sulaiman, Universiti Malaysia Sabah, Faculty of Food Science and Nutrition, Department of Food Science and Nutrition, 88400, Kota Kinabalu, Sabah, Malaysia,

E-mail: rossulma@ums.edu.my

ORCID: <https://orcid.org/0000-0003-1753-7093>

Jurnal 3 132

ORIGINALITY REPORT

0%

SIMILARITY INDEX

0%

INTERNET SOURCES

0%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

Exclude quotes On

Exclude bibliography On

Exclude matches < 3%

