

Evaluation of Formulated Diets Enriched by Spinach Extracts for the Broodstock Females, *Portunus pelagicus* (Linnaeus, 1758)

by

Submission date: 31-Mar-2023 03:23PM (UTC+0800)

Submission ID: 2051845318

File name: 283-290.pdf (615.98K)

Word count: 5610

Character count: 29008

<http://www.pjbs.org>

PJBS

ISSN 1028-8880

**Pakistan
Journal of Biological Sciences**

ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan



Research Article

Evaluation of Formulated Diets Enriched by Spinach Extracts for the Broodstock Females, *Portunus pelagicus* (Linnaeus, 1758)

¹Efrizal, ²Rusnam, ³Suryati, ¹Nofa Yolanda, ⁴Ferry Lismanto Syaiful and ⁵Ainul Mardiah

²⁰
¹Department of Biology, Faculty of Mathematics and Natural Sciences, Andalas University, Kampus Limau Manis, Padang, West Sumatera, 25163, Indonesia

¹⁰
²Department of Agricultural Engineering, Faculty of Agriculture, Andalas University, Kampus Unand Limau Manis, Padang, West Sumatera, 25163, Indonesia

³Faculty of Pharmacy, Andalas University, Kampus Unand Limau Manis, Padang, West Sumatera, 25163, Indonesia

⁴Faculty of Animal Science, Andalas University, Kampus Unand Limau Manis, Padang, West Sumatera, 25163, Indonesia

⁵Faculty of Science, Nahdlatul Ulama University of West Sumatera, Padang, Indonesia

Abstract

Background and Objective: Spinach extract (*Amaranthacea tricolor*) is a very prospective feed raw materials to stimulate the molting and growth of crab female broodstock. The purpose of this study was to evaluate the quality of broodstock females, *Portunus pelagicus* (Linnaeus, 1758) at different levels of spinach extract based on physical, organoleptic and chemical tests. **Materials and Methods:** Four different treatments of spinach extract (0, 250, 500 and 750 ng g⁻¹ crab, respectively) were used in this study. The female crab samples were collected from coastal region of Padang, West Sumatera and placed randomly in four concrete tanks (200 × 100 × 100 cm). Each concrete tank consist of five units of plastic box (45.5 × 32.5 × 16.5 cm) with the maximum density was one crab per box. **Results:** The organoleptic test showed that formulated diets enriched with spinach extract (0, 250 and 500 ng g⁻¹ crab) had a smooth texture, pungent aroma and brown. Whereas from the physics test results obtained good water stability (rupture velocity ranged from 89.20-105.40 min and solids dispersion ranged from 4.97-7.17%), hardness (92.66-98.07%) and sinking velocity (3.88-5.88 cm sec⁻¹) (p<0.05). The results also showed that formulated diet enriched with spinach extract doses of 250-750 ng g⁻¹ crab gave a value of delicacy of feed (0.195-0.386 g crab⁻¹ weight/day) which was significantly different (p<0.05) with 0 ng g⁻¹ crab (0.445 g crab⁻¹ weight/day). The chemical test shows the moisture content were 11.60%, ash 9.31%, protein 44.38%, fat 7.64% and carbohydrate 14.46%. **Conclusion:** It was concluded that there was a linear relationship between dose of spinach extracts in formulated diet and rupture velocity and a quadratic relationship between dose of spinach extracts in formulated diet and solid disperse, hardness and sinking velocity and delicacy of feed.

Key words: *Amaranthacea tricolor*, crab, formulated diet, *Portunus pelagicus*, spinach extract, broodstock females, rupture velocity, Linnaeus, 1758

Citation: Efrizal, Rusnam, Suryati, Nofa Yolanda, Ferry Lismanto Syaiful and Ainul Mardiah, 2019. Evaluation of formulated diets enriched by spinach extracts for the broodstock females, *Portunus pelagicus* (Linnaeus, 1758). Pak. J. Biol. Sci., 22: 283-290.

³⁶
Corresponding Author: Efrizal, Department of Biology, Faculty of Mathematics and Natural Sciences, Andalas University, Kampus Limau Manis, Padang, West Sumatera, 25163, Indonesia Tel: +6281378155502

²
Copyright: © 2019 Efrizal *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

³⁹
Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Commodity demand for crab increases every year, while the supply of this commodity is still depending on nature fishing^{1,4}. The efforts are needed towards controlled cultivation, but the constraints of this effort are still difficult to obtain the right amount and time, because seeding technology is needed. To anticipate this problem, it needs a technique to ripen the gonads of the mother crab which is the initial determinant of hatchery success. To stimulate the process of gonadal maturation, it is usually carried out by hormonal manipulation, feed and environmental manipulation^{5,6}. Improvement of nutrients in artificial feed for the gonadal maturation process has been carried out on various types of fish⁷. In mangrove crabs, *Scylla serrata* has also been carried out by several researchers^{8,9}. According to Millamena and Quinito⁸, for the maturation process of gonadal crabs, the balance diet of Omega-3 and Omega-6 is needed. Feed tested on crustaceans (*Penaeus japonicus*) containing a mixture of soybean oil (Omega-6 source) and squid liver oil (Omega-3 source) with a ratio of 3:4% is better for the gonad maturation process compared to feed containing soybean oil or squid liver oil only¹⁰. Likewise, the conclusions of several other researchers stated that feed which is quality is feed with a protein content, fat, carbohydrates, minerals and vitamins that are balanced^{8,9,11}.

The development of the next crab cultivation technology is quite promising with the emergence of soft shell production, with selling prices reaching twice as high as hard-shelled ones¹². The introduction of soft shell production technology to induce molting using spinach extract (*Amaranthus tricolor*) was discovered by Aslamyah and Fujaya¹³. The artificial feed enriched spinach extract (*vitomolt*) with protein content of 30.62 and 41.72% carbohydrate effectively stimulated the molting of mangrove crabs, with a percentage of molting¹⁴ (100%).

Recently, there is no research which is related to application of spinach extract (*A. tricolor*) to improve the nutritional quality of formulated diets for molting and growth of crab female broodstock. Therefore, this study was aimed to investigate the formulated diets enriched with spinach extract to find the best formula for molting and growth of crab female broodstock.

57 MATERIALS AND METHODS

Time and site: The study was conducted at Balai Benih Ikan Pantai (BBIP) Teluk Buo, Balai Benih Ikan (BBI) Bungus,

Padang and the Laboratory of Animal Physiology, Department of Biology, Padang, West Sumatera from April-July, 2018.

43 Experimental design and formulated diet: The method used in this study is completely randomized design (CRD) with four treatments and five replications of dietary with spinach extracts. They were treatment with Fdiet 1, formulated diet without spinach extract 0 ng g⁻¹ crab; Fdiet 2, formulated diet enriched with spinach extract 250 ng g⁻¹ crab; Fdiet 3, formulated diet enriched with spinach extract 500 ng g⁻¹ crab; and Fdiet 4, formulated diet enriched with spinach extract 750 ng g⁻¹ crab. Formulated diet¹⁵⁻¹⁷ is a modified formulation (Table 1) for the broodstock of mud crab, *Scylla serrata*⁸. Spinach extract was dissolved with 80% ethanol with the ratio 1:1 then homogenized. The solution was added 80% ethanol to 20 mL kg⁻¹ of feed and sprayed evenly into the test feed, then the feed was dried. The test feed is stored until it is ready for use¹³.

Organoleptic test: Organoleptic testing includes texture, odour and color of feed. The texture of the feed can be seen from the surface of the feed is smooth, fibrous or perforated. The odour of feed determines the quality of the feed as it is closely related to the recipient or the allure of the crab to the feed. The odour of the feed is determined by making the category not pungent, pungent and quite pungent. Feed color depends on the type of raw materials used¹⁸.

Physical test: Physical tests include the stability of the feed in water, the level of hardness, the sinking velocity and the delicacy of feed¹⁸.

Stability of feed in water: The stability test in water is a test of the endurance level of feed in the water or how long it takes until the feed is mushy and crumble. The stability of

Table 1: Modified composition of broodstock formulated diet for females blue swimming crab, *P. pelagicus* in g/100 g dry diet

Ingredients	Diet (g/100 g)
Fish flour	20
Clam flour	20
squid flour	20
wheat flour	5
Seaweed	4
Cod liver oil	5
Lecithin (lecithin)	3
Cholesterol	1
Vitamin mix	3
Mineral mix	4
Dicalcium phosphate	3
Vit. E (IU kg ⁻¹ feed)	600

feed in water including rupture velocity and dispersion of solids. The crash test measures how long it takes until the feed is destroyed in the water. This rupture test was observed visually. Included feed 10 bars into a beaker filled with 1 L of water, observations made every 5 min to know the feed was soft or not. Observation continues until the feed is broken/destroyed. The solid dispersion is carried out by feeding 10 g inserted into a 6×6×6 cm gauze box with pores of about 1 mm, then immersed in the aquarium for 2 h then lifted and dried under the sun. After that, it is weighed to a constant weight. The solid dispersion is calculated using the formula:

$$\text{Solid dispersion (\%)} = \frac{\text{Dry weight of initial feed} - \text{Dry weight of final feed}}{\text{Dry weight of initial feed}} \times 100$$

Hardness level test: The feed hardness rate test was measured by feeding 5 g of feed into a paralon pipe with a height of 1 m. Then the feed is loaded with a weight of a weighing 500 g. Feed that has been loaded then sieved using a sieve of 0.5 mm. The level of hardness is calculated in the percentage of non-destructive feed using a sieve.

Sinking velocity test: The sinking velocity test is performed by measuring the length of time it takes for feed to move from the water surface to the bottom of the maintenance medium. Feed five pellets put into a beaker glass with a bottom height of the container 20 cm from the surface of the water. The stopwatch is executed just as the feed is dropped into the water surface. The sinking speed is the distance in the feeding time until it is based on the measuring cup.

Delicacy feed test: The delicacy feed test was measured from the amount of feed consumed per crab body weight¹⁹. The female crab samples were collected from coastal region of Padang, West Sumatera and placed randomly in four concrete tanks (200×100×100 cm) which each put five units in a plastic box (45.5×32.5×6.5 cm) at maximum density of one crabs per box. Tanks were provided with substrate of around 15 cm thick layer of sand and adequate aeration^{2,11,20}. The crabs were maintained in monitored water depth of 25-30 cm with the salinity of 30-32 ppt, pH 7.77-7.96, temperature 26-27°C and DO 7.00-7.30 ppm. Each crab was provided with a shelter made of PVC pipe: 13 cm in diameter and 40 cm in length to serve as a refuge during molting. Formulated diets made from food wastes enriched by spinach extracts were fed daily at 3% biomass (1700-1800 h) and uneaten food was removed every morning.

Chemical test: Chemical test of feed includes determination of quantity and quality of nutrition in feed which is determination of proximate composition of protein, fat, ash, water, coarse fiber and NFE content in test feed²¹. This method refers to the test method of BSN²².

Data analysis: The Organoleptic data were analyzed descriptively by comparing the results obtained between treatments with supporting literature. While the data physical and chemical tests were analyzed by using one way ANOVA and then Duncan's test to know the difference between treatment²³. The standard deviation of each parameter and treatment was determined and expressed as the Mean±SE. The treatment effects were considered to be significant at p<0.05. Arcsine transformation is performed before analyzing data in percentage form. The relationship between dose of spinach extracts in formulated diet and physical tests were analysed using statistical computer software (version 19.0 for Windows, SPSS Inc., Chicago, IL), for significant correlation between them.

RESULTS

Organoleptic test: Based on the organoleptic test, the results have showed that different texture of diet treatment. The texture treatment of Fdiet 1 has a smooth texture, while the treatment of Fdiet 2, 3 and 4 has a fibrous texture. Organoleptic observations of feed odour also showed differences. The odour of treatment Fdiet 1 has a strong odour, Fdiet 2 and 3 smell quite stinging, while Fdiet 4 treatment has a non-stinging odour. Furthermore, the 3rd organoleptic test is the color of the feeds test. The results showed that the same result for all samples which the color is brown (Table 2).

Physical test: Observations of physical tests include: (a) The stabilization of feed in the water (rupture velocity and solid dispersion), (b) Level of feed hardness, (c) Velocity of feed sinking and (d) Delicacy of feed.

Stabilization of feed in the water

Rupture velocity: The rupture velocity at different doses of spinach extract was generally longest ranging from 89.20±3.63 to 105.40±3.55 min. The rupture velocity (min) decreased with increasing dose of spinach extract (p<0.05). The longest rupture velocity was achieved at 0 and 250 ng g⁻¹ crab, whereas the lowest (p<0.05) rupture velocity occurred at 750 ng g⁻¹ crab with 89.20±3.63 min (Table 2). The relationship between dose of spinach extracts in

Table 2: Organoleptic and physical tests of formulated diets made from food wastes enriched by spinach extracts for the broodstock females blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758)

Parameters of the test	Treatments (n = 5)				Regression equation	R ²
	Fdiet 1	Fdiet 2	Fdiet 3	Fdiet 4		
Organoleptic test						
Texture	Smooth	Fibrous	Fibrous	Fibrous		
Smell	Pungent	Quite pungent	Quite pungent	No pungent		
Color	Light brown	Light brown	Light brown	Light brown		
Physical test						
Stabilization of feed in water						
Rupture velocity (min)	105.40±3.55 ^a	100.20±2.95 ^{ab}	94.20±1.64 ^b	89.20±3.63 ^c	RV = -0.0218DSE+105.44 (Linear)	0.5563
Solids disperse (%)	6.69±0.37 ^a	6.01±0.40 ^{ab}	4.97±0.22 ^b	7.17±0.52 ^c	DS = 1×10 ⁻⁵ DSX ² -0.0085DSX+6.8732 (Quadratic)	0.4431
Hardness (%)	92.66±1.25 ^a	95.54±0.57 ^b	98.07±0.36 ^c	93.04±1.06 ^a	H = -3×10 ⁵ DSE ² +0.0252DSE+92.299 (Quadratic)	0.5636
Sinking velocity (cm sec ⁻¹)	3.88±0.36 ^a	5.20±0.19 ^b	5.88±0.16 ^c	4.97±0.17 ^b	SV = -9×10 ⁻⁶ DSE ² +0.0083DSE+3.8288 (Quadratic)	0.7313
Delicacy of feed (g crab ⁻¹ weight/days)	0.445±0.273 ^b	0.195±0.075 ^a	0.129±0.024 ^a	0.386±0.141 ^a	DF = 2×10 ⁻⁶ DSE ² -0.0016DSE+0.4518 (Quadratic)	0.1724

Means within a given column with different superscripts were significantly different ($p < 0.05$). Values are means ± standard errors (SE). n = Replication, Fdiet 1: Formulated diet without spinach extract 0 ng g⁻¹ crab, Fdiet 2: Formulated diet enriched with spinach extract 250 ng g⁻¹ crab, Fdiet 3: Formulated diet enriched with spinach extract 500 ng g⁻¹ crab and Fdiet 4: Formulated diet enriched with spinach extract 750 ng g⁻¹ crab

Table 3: Proximate composition of formulated broodstock diet for females blue swimming crab, *P. pelagicus* without spinach extract on a dry matter basis

Proximate composition (%)	Formulated diet (%)
Moisture	11.60
Ash	9.31
Crude protein	44.38
Fat	7.64
Carbohydrate	14.46

formulated diet and rupture velocity was linear with an equation of $RV = -0.0218DSE + 105.44$. This relationship was found to be highly significant ($R^2 = 0.5563$; $p < 0.05$).

Solids disperse: The doses of spinach extract (0, 250 and 500 ng g⁻¹ crab) significantly influenced ($p < 0.05$) the percentage of solid dispersion in formulated diet. The highest percentage of solid dispersion was achieved at spinach extract levels of 0, 250 and 750 ng g⁻¹ crab (6.01±0.40 to 7.17±0.52%), whereas the lowest percentage of solid dispersion is found at 500 ng g⁻¹ crab (4.97±0.22%). The relationship between dose of spinach extracts in formulated diet and solid disperse was found to be quadratic ($DS = 1 \times 10^{-5} DSX^2 - 0.0085DSX + 6.8732$; $R^2 = 0.4431$; $p < 0.05$) (Table 2).

Hardness level: Treatment of different doses of spinach extract (0, 250, 500 and 750 ng g⁻¹ crab) showed different effects on hardness level in formulated diet (Table 2). The highest ($p < 0.05$) percentage of hardness level was achieved by spinach extracts in formulated diet at 500 ng g⁻¹ crab (98.07±0.36%) compared to hardness level at 0 ng g⁻¹ crab

(92.66±1.25%), 250 ng g⁻¹ crab (95.54±0.57%) and 750 ng g⁻¹ crab (93.04±1.06%). A quadratic relationship of $H = -3 \times 10^5 DSE^2 + 0.0252DSE + 92.299$; $R^2 = 0.5636$; $p < 0.05$) was found between dose of spinach extracts in formulated diet and hardness.

Sinking velocity: The sinking velocity of Fdiet 3 (5.88±0.16 cm sec⁻¹) revealed the increase ($p < 0.05$) at doses of spinach extract 500 ng g⁻¹ crab, whereas the lowest ($p < 0.05$) sinking velocity occurred at 0 ng g⁻¹ crab with 3.88±0.36 cm sec⁻¹ (Table 2). The relationship between dose of spinach extracts in formulated diet and sinking velocity was found to be quadratic ($SV = -9 \times 10^{-6} DSE^2 + 0.0083DSE + 3.8288$; $R^2 = 0.7313$; $p < 0.05$).

Delicacy of feed: Data (Table 2) showed that doses of spinach extract in formulated diets ($p < 0.05$) affected delicacy of feed. The lowest dose of spinach extract (0 ng g⁻¹ crab) resulted in the highest feed delicacy (0.445±0.273 g crab⁻¹ weight/days). Regression analysis showed the relationship to be quadratic ($DF = 2 \times 10^{-6} DSE^2 - 0.0016DSE + 0.4518$; $R^2 = 0.1724$; $p > 0.05$).

Chemical test: Based on the results of chemical testing with the proximate analysis of the test feed in Table 3. The results showed that the nutritional composition of the test feed were containing 11.60% moisture content, 9.31% ash, 44.38% protein, 7.64% fat, 14.46% carbohydrate and which is a fairly balanced percentage with the needs of blue swimming crabs.

DISCUSSION

The application of spinach extract on formulated diets have an effect on the textures and odour of diet. Then, the composition of the raw materials and appearance also have an effect on the texture of feed. The surface texture of feed showed smooth, fibrous or perforated²⁴. Furthermore, the fineness of the raw material, the amount of fiber and the binder used in the composition have an effect on the texture of formulated diet. The texture of feed and the level of fineness of the raw material have affected the compactness and flatness²². The different of feed odour has related to the composition of raw materials such as; spinach extract doses. According to previous report, the type and amount of spinach extract added to the process making of feed has effect to odour²⁴. Furthermore, the good formulated diet has natural scent odour and taste^{7,25}. The colour level of the formulated diets (Table 2) was within the acceptable limits (light brown). There was no different colour in each diet, because, the different of spinach doses extracts has no effect on feed composition. Similar results with previous study, the colour of formulated diet has related or depends on the type of raw material²⁴.

The rupture velocity of formulated diet is strongly affected by the fineness of raw materials²⁶. Furthermore, the rupture velocity of pellets can be greatly improved by proper selection of feed ingredients, processing techniques and the use of proper processing equipment²⁷. The lowest value of rupture velocity (Fdiet 4) is found at doses spinach extract 750 ng g⁻¹ crab. The lower rupture velocity at high doses spinach extract may be due to the feed raw material to be less bound or less compact which results in accounting for the difference in water stability. The rupture velocity of feed is influenced by several factors, such as the amount and type of raw material²⁸. The results from this study showed spinach extract to be water stable pellet than those by Solomon *et al.*²⁹ studies who produced a water stability of around 56-57 min on feed made from local carbohydrate sources and added with *Saccharomyces cerevisiae* yeast. However, it is still lower than the general requirements of crustacean feed. According to a report, the stability of feed in water ranges²⁸ from 3-5 h.

The high doses of spinach extract 750 ng g⁻¹ crab provided the highest percentage of solid dispersion (7.17±0.52%), while the lowest value (4.97±0.22%) was observed at doses of spinach extract 500 ng g⁻¹ crab. The level of solid dispersion should be no more than 10%, because it greatly affected the quality and quantity of nutrients contained in the feed³⁰. In general, all the formulated diets in this study already have good solid dispersion values

(4.97±0.22-7.17±0.52%) compared to the Aslamyah and Karim²⁴ studies who produced solid dispersions above 10% (11.14-11.87%).

In the present study, the results indicated that the highest level of hardness was found in the Fdiet 3, which was 98.07% because its caused of the fineness of the feed and the composition used were balanced. The level of hardness of feed is influenced by the fineness of the particles and the level of density of particles as often as the constituent ingredients of feed¹⁹. The lowest percentage level of hardness in the Fdiet 1 was 92.66%. The difference may be due to feeding of spinach added which made the feed a little more compact. Therefore, it can be concluded that low percentage was included in the level of good hardness as the level of hardness of treatment above 80% was definitely high²⁴. Meanwhile, other studies showed that good pellets are harder than usual pellets²⁸. The range of levels of feed hardness obtained in this study was almost same as that obtained from tiger shrimp feed containing several types of seaweed flour as adhesive¹⁹, namely 92.27-94.67%.

Throughout the monitoring period, the treatment of Fdiet 3 with the addition of spinach extract as much as 500 ng g⁻¹ crab had the highest and best sinking velocity of 5.88 cm sec⁻¹, followed by Fdiet 2 and 4, respectively. Spinach extract may directly inclusion increased the specific gravity of the feed so that the feed will sink faster. Specific gravity of feed has been reported by Mudjiman²⁸, feed buoyancy has to do with it, the greater specific gravity of the feed compared to the specific gravity of water the faster it will sink. It was also reported that shrimp, catfish and demersal fish pellets should sink faster than usual. The Fdiet 1 treatment has the longest sinking velocity of 3.88 cm sec⁻¹. Therefore, it can be concluded that effect of spinach are increase in volume of feed composition²⁸.

In the short term study reported that the delicacy test of feed is increased by attractants and odours by it contains¹⁹. The findings of current study were also supported by previous reports that good quality food has a distinctive odour that is favored by culture. The low tasty power in the Fdiet 1 treatment with a value of 0.445 g was thought to be without additives which spinach extract had a pungent odour not liked by the crab, so that it had low tasty power^{25,26}.

Current study showed the result of proximate composition of the formulated diet. The result showed that the nutritional composition of the test feed contains a fairly balanced percentage with the needs of crabs. The formulated diet has lower moisture content. Moisture content is within the recommended value (<12%)^{18,31}. Moisture content of formulated diet is an important parameters of their

susceptibility to microbial spoilage. When the moisture content is on the high side, it encourages the growth of micro-organisms⁷. Ash content of the formulated diet was relatively low (9.31%). The BSN recommended (\leq 13%) ash contents for the formulated diet¹⁸. In addition, Piedecausa *et al.*³² and Ayuba and Iorkohol³³ recommended ash content of fish formulated diets are lower than recommended value by BSN¹⁸. This might be due to the ash effect on digestibility and growth³⁴⁻³⁶.

High protein content of diet is thought to be due to the high protein content of the raw materials for the formulated diet used, such as fish meal, squid and shellfish^{8,37}. Protein content of diet is within the recommended (40-50% in dry form for crab growth)^{2,8,17,38}. The fat content of the diet was obtained 7.64%, this contents are within the recommended value¹⁸ of BSN of \geq 5%. Low fat content of formulated diet may be necessary to prolong its shelf life also by avoiding lipids peroxidation. Furthermore, the high fat content will cause side effects that decrease food consumption, growth and liver degeneration¹⁰. Thus, the optimal fat requirement of omnivorous cultivation group³⁹ is 6-8%. Carbohydrates, just like fats are needed for energy. Carbohydrates content of the formulated diet was relatively lower than recommended value by Alava and Pascual⁴⁰ (25-30%). The carnivorous group cultivation needs around 10-20% carbohydrates³⁹.

CONCLUSION

Based on organoleptic, physical and chemical tests on evaluated feed, it was showed that the enrichment of spinach extract in different formulated diets have produced different feed quality. The proximate analysis showed that the nutrients of feed tests were within the range requirement blue swimming crab. There was a linear relationship between dose of spinach extracts in formulated diet and rupture velocity and a quadratic relationship between dose of spinach extracts in formulated diet and solid disperse, hardness, sinking velocity and delicacy of feed.

SIGNIFICANCE STATEMENT

Formulated diet is an alternative media application of spinach extract (SE) which is known containing molting stimulant. However, the utilization of the formulated diet is an expensive fed with a very high protein concentration and it's mainly produced from fish based materials. Therefore, it is

needed further study regarding formulated diet which substitute with vegetable materials that could be used for stimulated the molting and growth of blue swimming crabs.

ACKNOWLEDGMENTS

This study was supported by research grant Research and Community Service Institution, Andalas University (No. 38/UN.16.17/PP.PGB/LPPM/2018). The authors wish to thank all the staff at Balai Benih Ikan Pantai (BBIP), Teluk Buo and Balai Benih Ikan (BBI) Bungus, Padang, Department of Maritime and Fisheries Affairs and the Laboratory of Animal Physiology Department of Biology, Faculty of Math and Science, Andalas University, Padang, West Sumatera for technical assistance rendered.

REFERENCES

1. FAO, 2000. FAO Yearbook: Fisheries Statistic. Food and Agriculture Organization (FAO), Rome, Italy, pp: 713.
2. Efrizal, A.A., M.S. Kamarudin, C.R. Saad and S.M.N. Amin, 2015. Some aspects of reproductive biology of blue swimming crab (*Portunus pelagicus* (Linnaeus, 1758)) under laboratory conditions. *J. Fish. Aquat. Sci.*, 10: 77-91.
3. Hassan, A., T.N. Hai, A. Chatterji and M. Sukumaran, 2011. Preliminary study on the feeding regime of laboratory reared mud crab larva, *Scylla serrata* (Forsskal, 1775). *World Applied J.*, 14: 1651-1654.
4. Romano, N., M.A. Safee, M. Ebrahimi and A. Arshad, 2016. Fatty acid compositional changes during the embryonic development of the swimming crab, *Portunus pelagicus* (Tunidae: Decapoda). *Invertebr. Reprod. Dev.*, 60: 112-117.
5. Lockwood, A.P.M., 1967. Aspect of the physiology of crustacean. The Aberdeen University Press.
6. Primavera, J.H., 1985. A review of maturation and reproduction in closed thelycum penaeids. Proceedings of the 1st International Conference on the Culture of Penaeid Prawns/Shrimps, December 4-7, 1984, Iloilo City, Philippines, 4: 47-64.
7. Watanabe, W.O., M.S. Alam, A.D. Ostrowski, F.A. Montgomery, J.E. Gabel, J.A. Morris Jr. and P.J. Seaton, 2016. Live prey enrichment and artificial microdiets for larviculture of atlantic porgy *Pagrus pagrus*. *Aquacult. Rep.*, 3: 93-107.
8. Millamena, O.M. and E.T. Qunitio, 2000. The effects of diets on reproductive performance of eyestalk ablated and intact mud crab *Scylla serrata*. *Aquaculture*, 181: 81-90.
9. Aslamyah, S. and Y. Fujaya, 2009. Formulasi pakan buatan khusus kepiting yang berkualitas murah dan ramah lingkungan. *Sains Teknol.*, 9: 133-141.

10. Alava, V.R., A. Kanazawa, S.I. Teshima and S. Koshio, 1993. Effect of dietary phospholipids and n-3 highly unsaturated fatty acids on ovarian development of kuruma prawn. *Nippon Suisan Gakkaishi*, 59: 345-349.
11. Efrizal and Rusnam, 2017. The influence of dietary vitamin E on the gonad development of blue swimming crab *Portunus pelagicus* (Linnaeus, 1758). *Aust. J. Basic Applied Sci.*, 11: 7-15.
12. Fujaya, Y., 2011. Growth and molting of mud crab administered by different doses of vitomolt. *J. Akuakultur Indones.*, 10: 24-27.
13. Aslamyah, S. and Y. Fujaya, 2011. Effectiveness of artificial diet enriched by spinach extract on molting stimulation to induce soft shell crab. *J. Akuakultur Indones.*, 10: 8-15.
14. Aslamyah, S. and Y. Fujaya, 2012. Stimulasi molting dan pertumbuhan kepiting bakau (*Scylla* sp.) melalui aplikasi pakan buatan berbahan dasar limbah pangan yang diperkaya dengan ekstrak bayam. *ILMU KELAUTAN: Indones. J. Mar. Sci.*, 15: 170-178.
15. Efrizal, 2013. Perkembang gonad induk rajungan, *Portunus pelagicus* (Linnaeus, 1758), dengan manipulasi pakan alami dan buatan. *Proceedings of Seminar Nasional Biodiversitas dan Ekologi Tropika Indonesia (BioETI) Universitas Andalas*, September 14, 2013, Padang, pp: 45-51.
16. Efrizal, E., 2017. Effects of stocking density on survival rate and larval development of blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758) under laboratory conditions. *Aquacult. Aquarium Conserv. Legislation-Int. J. Bioflux Soc. (AACL Bioflux)*, 10: 217-226.
17. Efrizal, E., I.J. Zakaria, R. Rusnam, S. Suryati and N. Yolanda, 2018. Studies on biological test formulated diets supplementation of vitamin E for the broodstock of females blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758). *F1000 Res.*, Vol. 7 10.12688/f1000research.15885.2.
18. Badan Standarisasi Nasional, 2006. SNI 01-4266-2006 Tepung ikan/bahan baku pakan. Badan Standarisasi Nasional, Jakarta. <https://www.scribd.com/document/372992348/259098711-Pakan-Buatan-Bagi-Ikan-Mas-Cyprinus-Carpoi-L-Pada-Idaya-Intensif-pdf>
19. Saade, E. and S. Aslamyah, 2009. Uji fisik dan kimiawi pakan buatan untuk udang windu *Penaeus monodon* fab. yang menggunakan berbagai jenis rumput laut sebagai bahan perekat. *Tora (Ilmu Kelautan Perikanan)*, 19: 107-115.
20. Efrizal, 2015. Effects of salinity on survival rate and larval development of blue swimming crab *Portunus pelagicus* (Linnaeus, 1758) under laboratory conditions. *Aust. J. Basic Applied Sci.*, 9: 557-563.
21. Balazs, G.H., E. Ross and C.C. Brooks, 1973. Preliminary studies on the preparation and feeding of crustacean diets. *Aquaculture*, 2: 369-377.
22. Badan Standarisasi Nasional, 2002. SNI 01-2891-1992 cara perlakuan makanan dan minuman. Badan Standar Nasional, Jakarta. <https://www.slideshare.net/Fitrijasmineandriani/sni-01-28911992-cara-uji-makanan-minuman>
23. Steel, R.G.D. and J.H. Torrie, 1980. *Principles and Procedures of Statistics: A Biometrical Approach*. 2nd Edn., McGraw Hill Book Co., New York, USA., ISBN-13: 9780070609266, pages: 633.
24. Aslamyah, S. and M.Y. Karim, 2013. Organoleptic, physical and chemical tests of artificial feed for milk fish substituted by earthworm meal (*Lumbricus* sp.). *J. Akuakultur Indones.*, 12: 124-131.
25. Saenz de Rodriganez, M.A., B. Gander, M. Alaiz and F.J. Moyano, 2011. Physico chemical characterization and *in vitro* digestibility of commercial feeds used in weaning of marine fish. *Aquacult. Nutr.*, 17: 429-440.
26. Murdinah, T., S. Suwarno and P. Sumpeno, 1991. Mempelajari jenis bahan pemikat untuk pakan udang. *J. Penelitian Capanen Perikanan*, 70: 29-36.
27. Lim, C. and G. Cuzon, 1994. Water stability of shrimp pellet: A review. *Asian Fish. Sci.*, 7: 115-127.
28. Djiman, A., 2009. *Makanan ikan*. Penebar swadaya. Jakarta.
29. Solomon, S.G., G.A. Ataguba and A. Abeje, 2011. Water stability and floatation test of fish pellets using local starch sources and yeast (*Saccharomyces cerevisiae*). *Int. J. Latest Trends Agri Food Sci.*, 1: 1-5.
30. Lim, C. and G.W. Dominy, 1991. Performance of binders in pelleted shrimps diet. *Proceedings of the Aquaculture Feed Processing and Nutrition Workshop, September 19-25, 2005, Bantul and Indonesia*, pp: 149-157.
31. ADCP., 1983. *Fish Feeds and Feeding in Developing Countries*. Aquaculture Development and Coordination Program, FAO, Rome, pp: 97.
32. Piedecausa, M.A., F. Aguado-Gimenez, B. Garcia-Garcia, G. Ballester and T. Telfer, 2009. Settling velocity and total ammonia nitrogen leaching from commercial feed and faecal pellets of gilthead seabream (*Sparus aurata* L. 1758) and seabass (*Dicentrarchus labrax* L. 1758). *Aquacult. Res.*, 10: 1703-1714.
33. Ayuba, V.O. and E.K. Iorkohol, 2013. Proximate composition of some commercial fish feeds sold in Nigeria. *J. Fish. Aquat. Sci.*, 37: 248-252.
34. Dale, N., 1997. Metabolizable energy of meat and bone meal. *Applied Poult. Res.*, 6: 169-173.
35. Mendez, A. and N. Dale, 1998. Rapid assay to estimate calcium and phosphorus in meat and bone meal. *J. Applied Poult. Sci.*, 7: 309-312.
36. Wang, X. and C.M. Parsons, 1998. Effect of raw material source, processing systems and processing temperatures on amino acid digestibility of meat and bone meals. *Poult. Sci.*, 77: 834-841.

37. Oniam, V., L. Chuchit and W. Arkronrat, 2012. Reproductive performance and larval quality of blue swimming crab (*Portunus pelagicus*) broodstock, fed with different feeds. Songklanakarin J. Sci. Technol., 34: 381-386.
38. Ding, L.Y., M. Jin, P. Sun, Y. Lu and H.N. Ma *et al.*, 2017. Cloning, tissue expression of the fatty acid-binding protein (Pt-FABP1) gene and effects of dietary phospholipid levels on *fabp* and vitellogenin gene expression in the female swimming crab *Portunus trituberculatus*. Aquaculture, 474: 57-65.
39. Creswell, R.L., 1993. Feeding and Nutrition. In: Aquaculture Desk Reference. Creswell, R.L., (Eds.). Springer, Boston, MA., New York, ISBN: 978-1-4684-7115-1, pp: 102-146.
40. Alava, V.R. and F. Pascual, 1987. Carbohydrate requirements of *Penaeus monodon* (Fabricius) juveniles. Aquaculture, 61:211-217.

Evaluation of Formulated Diets Enriched by Spinach Extracts for the Broodstock Females, *Portunus pelagicus* (Linnaeus, 1758)

ORIGINALITY REPORT

23%

SIMILARITY INDEX

20%

INTERNET SOURCES

21%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Timóteo T. Watanabe, Laura S. López-Greco, Fernando J. Zara. "Seminal fluid and spermatophore production in a western Atlantic invasive swimming crab, *Charybdis hellerii*, reveals a different pattern to *Portunoidea*", *Arthropod Structure & Development*, 2022
Publication 1%
- 2 oap-journals.org
Internet Source 1%
- 3 Carlos Cortés, Germán E. Merino. "Settling velocity and particle size bioengineering parameters gathered from solids generated by wild-caught premature punctuated snake-eels (*Ophichthus remiger*) reared in a marine prototype recirculating aquaculture system", *Aquacultural Engineering*, 2020
Publication 1%
- 4 uncw.edu
Internet Source 1%
- 5 www.fao.org

Internet Source

1 %

6

www.frontiersin.org

Internet Source

1 %

7

docsdrive.com

Internet Source

1 %

8

S Prasetyo, S Anggoro, T R Soeprbowati. "Potential of water hyacinth (*Eichhornia crassipes* (Mart.) Solms) in Rawapening lake as raw material for fish feed", *Journal of Physics: Conference Series*, 2021

Publication

1 %

9

idosi.org

Internet Source

1 %

10

www.ajol.info

Internet Source

1 %

11

Mat Ghani Hilm, Mhd Ikhwanuddi. "Salinity and Preservation Conditions of Lipofuscin Extracts in Blue Swimming Crab, *Portunus pelagicus*", *Pakistan Journal of Biological Sciences*, 2020

Publication

1 %

12

Adedokun, S. A., and O. Adeola. "Apparent metabolizable energy value of meat and bone meal for white pekin ducks", *Poultry Science*, 2005.

Publication

1 %

13	Yushinta Fujaya, Letty Fudjaja, Juanita Mandagi, Erwan Kow, Wahyudi. "Pilot project for the application of "Vitomolt" which combines probiotic and phytobiotic properties in commercial shrimp culture; Production studies and economic valuation", IOP Conference Series: Earth and Environmental Science, 2021 Publication	1 %
14	Submitted to Kampala International University Student Paper	1 %
15	journal.ugm.ac.id Internet Source	1 %
16	www.scialert.net Internet Source	1 %
17	123dok.com Internet Source	1 %
18	eprints.cmfri.org.in Internet Source	<1 %
19	Carlos I Perez-Rostro. "Quantitative genetic parameter estimates for size and growth rate traits in Pacific white shrimp, Penaeus vannamei (Boone 1931) when reared indoors", Aquaculture Research, 6/2003 Publication	<1 %
20	oamjms.eu Internet Source	<1 %

21 Bamisaye, F.A., Emmanuel Oladipo Ajani, I.O. Nurain, and J.B. Minari. "Nutritional Evaluation of Some Fortified Traditional Weaning Meals from Oriade Local Government Area of Osun State, Nigeria", International Journal of Applied Sciences and Biotechnology, 2014.
Publication

22 Jirawat Saetan, Thanyaporn Senarai, Montakan Tamtin, Wattana Weerachatyanukul et al. "Histological organization of the central nervous system and distribution of a gonadotropin-releasing hormone-like peptide in the blue crab, Portunus pelagicus", Cell and Tissue Research, 2013
Publication

23 scholarsjunction.msstate.edu
Internet Source

24 www26.us.archive.org
Internet Source

25 ugspace.ug.edu.gh
Internet Source

26 Submitted to Universitas Diponegoro
Student Paper

27 Yushinta Fujaya, Nita Rukminasari, Nur Alam, Muhammad Rusdi, Hanafiah Fazhan, Khor Waiho. "Is limb autotomy really efficient compared to traditional rearing in

soft-shell crab (*Scylla olivacea*) production?", Aquaculture Reports, 2020

Publication

28

law.resource.org

Internet Source

<1 %

29

Andi Aliah Hidayani, Yushinta Fujaya, Dody Dharmawan Trijuno, Moh. Tauhid Umar et al. "REPRODUCTIVE PERFORMANCE OF INTRASPECIFIC HYBRIDS OF BLUE SWIMMING CRAB (*Portunus pelagicus*)", Indonesian Aquaculture Journal, 2022

Publication

<1 %

30

www.slideshare.net

Internet Source

<1 %

31

Submitted to Monash University

Student Paper

<1 %

32

www.updatepublishing.com

Internet Source

<1 %

33

Khor Waiho, Mhd Ikhwanuddin, Juliana C. Baylon, Mohamad Jalilah, Nita Rukminasari, Yushinta Fujaya, Hanafiah Fazhan. "Moult induction methods in soft - shell crab production", Aquaculture Research, 2021

Publication

<1 %

34

Warren Dominy. "A Comparative Study of the Physical and Biological Properties of Commercially-Available Binders for Shrimp Feeds", Journal of Applied Aquaculture, 01/14/2004

<1 %

35	mail.scialert.net Internet Source	<1 %
36	www.entomologyjournals.com Internet Source	<1 %
37	Aberra Melesse, Herbert Steingass, Margit Schollenbe, Markus Rodehutsco. "Chemical Compositions and in vitro Methane Production Potentials of Some Agro-industrial By-products and Crop Residues", <i>Journal of Applied Sciences</i> , 2019 Publication	<1 %
38	journal.unhas.ac.id Internet Source	<1 %
39	journals.plos.org Internet Source	<1 %
40	nanopdf.com Internet Source	<1 %
41	www.scribd.com Internet Source	<1 %
42	jifro.ir Internet Source	<1 %
43	repository.lppm.unila.ac.id Internet Source	<1 %
44	Khaleg Manerii, Amin Oujifard, Ahmad Ghasemi, Mansour Torfi Mozanzadeh. "Reproductive performance and vitellogenin	<1 %

mRNA transcript abundance in the hepatopancreas of female *Litopenaeus vannamei* fed diets with different soy lecithin content", *Animal Reproduction Science*, 2019

Publication

45 Peng Sun, Min Jin, Liyun Ding, You Lu, Hongna Ma, Ye Yuan, Qicun Zhou. "Dietary lipid levels could improve growth and intestinal microbiota of juvenile swimming crab, *Portunus trituberculatus*", *Aquaculture*, 2018

Publication

46 complete.bioone.org

Internet Source

47 ejournal.almaata.ac.id

Internet Source

48 C.R.L. Amaral, V. Maciel, R. Goldenberg-Barbosa, D.A. Silva, A. Amorim, E.F. Carvalho. "Tuna fish identification using mtDNA markers", *Forensic Science International: Genetics Supplement Series*, 2017

Publication

49 E Saade, A Solicha, I R Fadillah. "Effect of seaweed, *Kappaphycus alvarezii* fermentation by various fermenters combinations as thickener on gel strength, attractiveness and palatability of gel diet in *Tilapia*, *Oreochromis niloticus*", IOP

Conference Series: Earth and Environmental Science, 2020

Publication

-
- | | | |
|----|--|------|
| 50 | staff.unand.ac.id
Internet Source | <1 % |
| 51 | Mhd Ikhwanuddi, Nur Salwani Ah, Syahnnon Mohamad, Ambok Bolong Abo. "Growth and Survival of Blue Swimming Crab, <i>Portunus pelagicus</i> Larvae at Different Photo Period and Light Intensity", <i>Asian Journal of Biological Sciences</i> , 2019
Publication | <1 % |
| 52 | Rawles, S.D.. "Evaluation of poultry by-product meal in commercial diets for hybrid striped bass (<i>Morone chrysops</i> @VxM. <i>saxatilis</i> @M) in recirculated tank production", <i>Aquaculture</i> , 20060908
Publication | <1 % |
| 53 | li01.tci-thaijo.org
Internet Source | <1 % |
| 54 | www.ijsh-t-journals.org
Internet Source | <1 % |
| 55 | www.rjpharmacognosy.ir
Internet Source | <1 % |
| 56 | Quy Moc Ong, Ravi Fotedar, Thy Thi Truong Ho. "Impact of different rearing systems on survival, growth and quality of mud crab (<i>Scylla paramamosain</i>) megalopae reared | <1 % |

from early zoeae", Aquaculture International, 2019

Publication

57 biodiversitas.mipa.uns.ac.id <1 %
Internet Source

58 oacis.repo.nii.ac.jp <1 %
Internet Source

59 researcherslinks.com <1 %
Internet Source

60 www.ijsrp.org <1 %
Internet Source

61 Medina, A.. "A comparative study of the ovarian development in wild and pond-reared shrimp, *Penaeus kerathurus* (Forskål, 1775)", *Aquaculture*, 19961210 <1 %
Publication

62 N Alam, Y Fujaya, Haryati, D K Sari, M Achmad, M Rusdi, N Farizah. "The effect of *Melastoma malabathricum* leaf extract on growth and spawning of blue swimming crab (*Portunus pelagicus*)", *IOP Conference Series: Earth and Environmental Science*, 2019 <1 %
Publication

63 Wade O. Watanabe, Md. Shah Alam, Andrew D. Ostrowski, Frank A. Montgomery, Jennifer E. Gabel, James A. Morris, Pamela J. Seaton. "Live prey enrichment and artificial microdiets for larviculture of Atlantic red <1 %

porgy Pagrus pagrus", Aquaculture Reports,
2016

Publication

64

Bligh, E. G., and W. J. Dyer. "A RAPID METHOD OF TOTAL LIPID EXTRACTION AND PURIFICATION", Canadian Journal of Biochemistry and Physiology, 1959.

Publication

<1 %

65

C.J. SIMON. "The effect of dietary carbohydrates on the growth response, digestive gland glycogen and digestive enzyme activities of early spiny lobster juveniles, Jasus edwardsii : Jasus edwardsii response to dietary carbohydrates", Aquaculture Nutrition, 12/2011

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off