

**Bukti Korespondensi Proses Review dan Bukti Revisi Selama Proses Review Jurnal
Syarat Khusus untuk Jabatan Guru Besar**


Nama : Dr. Ir. Gita Ciptaan, MP
NIP : 195911101986032003
NIDN : 0010115905
Unit Kerja : Fakultas Peternakan Universitas Andalas

Judul artikel	:	The Effect of Fermented Palm Kernel Cake Layer Quail Rations on Production Performance and Eggshell Thickness
Nama Jurnal	:	International Journal of Veterinary Science
Vol/No/Hal/Tahun	:	11(3): 400-403, 2022
Publisher	:	Publisher USP (Unique Scientific Publishers)
SJR	:	0,28 (Quartil Q3 bidang <i>veterinary</i> sejak 2020 hingga sekarang) https://www.scimagojr.com/journalsearch.php?q=21100820608&tip=sid

No	Perihal	Tanggal
1	Submitted	: 07 September 2021
2	Review report:1 st	: 12 Oktober 2021
3	Revised version: 1 st	: 13 Oktober 2021
4	Review report:2 nd	: 13 Oktober 2021
5	Revised version:2 nd	: 14 Oktober 2021
6	Accepted	: 21 Oktober 2021
7	proses <i>post-review</i> manuskrip	: 20 November 2021
8	Published	: 01 Juni 2022

1. Submitted (7 September 2021)

Tanggal 07 September 2021: artikel dikirim (submitted). Original paper terlampir.



Gambar : Email Author's Original Submission

MANUSKRIP VERSI PERTAMA

The effect of fermented palm kernel cake by adding humic acid in layer quail ration on production performance and eggshell thickness

Gita Ciptaan, Mirnawati, Ferawati and Malik Makmur¹

Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, Andalas University, Jl. Raya Unand, Kampus Limau Manis, Padang 25163, West Sumatera, Indonesia

gciptaan@ansci.unand.ac.id

¹ Postgraduate Student, Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, Andalas University, Jl. Raya Unand, Kampus Limau Manis, Padang 25163, West Sumatera, Indonesia

Abstract

The experiment was conducted to evaluate the effect of palm kernel cake fermented with *Sclerotium rolfsii* added with humic acid in rations on production performance and quail egg quality. The material used in this study were two hundred layer quail with 14 weeks of age (*Coturnix coturnix japonica*). Research treatment were used fermented palm kernel cake (FPKC) with 0 %, 5 %, 10 %, 15% and 20% in ration composition. Feeding trial lasts for 8 weeks. Research method used completely randomized design with four replications. The results of this research showed that feed intake, egg production, feed conversion, egg weight and eggshell thickness of layer quail were non-significant ($P>0.05$). In conclusion, palm kernel cake fermented with *S. rolfsii* can be used up to 20% in laying quail ration.

Keyword: egg quality, Japanese quail, palm kernel cake, *Sclerotium rolfsii*

Introduction

Palm kernel cake (PKC) is a by-product of palm oil processing that is potentially used as feedstuff for poultry. Seen from nutrient content of PKC as follows: crude protein 16.07%, crude fiber 21.30%, crude fat 8.23%, Ca 0.27% and P 0.94% and Cu 48.4 ppm (Mirnawati et al 2010). Although the crude protein content is quite high, but its use in poultry rations is still limited. According to Rizal (2000) get PKC can be used up to 10% or replace 40% of soybean meal in broiler ration. This is due to the high coarse fibers in β -manan (Sundu et al 2006). While poultry does not have fiber-breaking enzymes and manan in digestive tract. Therefore, it is necessary to process PKC first to improve its quality with fermentation biotechnology using cellulolytic and mananolitic molds (Meryandini et al 2008; Mirnawati et al 2018; Purwadaria and Haryati 2003) which can decrease the content of crude fiber and manan as well as the quality of palm kernel cake will be increased so that it can be replacing soybean meal in poultry rations.

Cellulolytic and mananalytic microbes which can be used for fermentation of palm kernel cake is *Sclerotium rolfsii*. Razak et al (2006) suggests that the manannase enzyme activity of *S. rolfsii* is higher than that of *Aspergillus niger*. fermentation of palm kernel cake with *S. rolfsii* has been carried out and gave the following results 26.90% crude protein, 54.86% nitrogen retention, 14.86% crude fiber, 58.41% crude fiber digestibility, 0.22% crude fat and 2557.6 kcal/kg. Although there is an increase in nutrient content and quality of palm kernel cake but its utilization in the ration is still limited 25% in broiler rations.

Mirnawati et al (2017) has been processing palm kernel cake with the addition of humic acid in the fermentation process with *S. rolfsii*. The results of this study showed an increase in nutrient content such as crude protein (27.43%), decreased crude fiber (11.53%), improving nitrogen retention (59.17%) and crude fiber digestibility (55.40%). The increasing of nutrient quality of palm kernel after fermentation is expected to be used as quail feed ingredients. Therefore, it is necessary to conduct research to determine the use of fermented palm kernel cake with *S. rolfsii* in rations on production performance and quality of laying quails.

Materials and methods

Two hundred for 14 weeks old quail laying were assigned this experiment. This laying quail was kept in individual cage (45 × 20 × 30 cm), there were 10 laying quail per unit of experiment. The experiment was performed in a completely randomized design (CRD) with five treatments (0, 5, 10, 15, and 20% FPKC) and four replicates. The diets were formulated in iso-protein 20% and iso-caloric 2700 kcal/kg ration. Diet formulation, nutrient and metabolizable energy contents of treatment diets were figured in Table 1. Diet formulation consists of yellow corn, rice bran, meat meal, CP 126 cocentrated feed (Charoen Pokphand Indonesia), top mix and FPKC. Experimental diet and drinking water were provided *ad-libitum*.

Table 1: Diet formulation (%), nutrient content (%) and metabolizable energy (kcal/kg)

	FPKC in diet (%)				
	0	5	10	15	20
Corn	45.5	45	44	43	43
Rice brand	20	18	16	13.5	11.5
Meat meal	6	6	6	6	6
CP 126 concentrate	26.5	24	22	20.5	17.5
FPKC	0	5	10	15	20
Mineral	1	1	1	1	1
Top mix	1	1	1	1	1
Total (%)	100	100	100	100	100
Crude protein	20.06	20.04	20.16	20.44	20.42
Crude fat	3.46	3.40	3.34	3.28	3.29
Crude fiber	6.13	6.17	6.23	6.27	6.50
Calcium	2.94	2.80	2.63	2.60	1.94
Phosphor	0.88	0.89	0.91	0.93	0.82
Metabolizable energy	2710	2708	2703	2706	2725

Palm kernel cake was the product of 80% PKC plus 20% rice bran that was fermented with *S. rolfesii* and addition of 200 ppm humic acid. The dose of inoculum that administered was 10% of substrate and incubated for 7 days. After harvesting the product, FPKC then dried and milled then mixed in quail diets. Feeding period of layer quail lasts for two months or eight consecutive weeks. Details of feeding or diet treatment composition showed in Table 1.

Collected data were feed intake (g/head/day), quail day egg production (%), egg weight (g/egg/head), feed conversion, egg mass production (g/head/day) and eggshell thickness (mm) of laying quail were measured following Nuraini et al (2012). All of the data were analyzed by analysis of variance based on completely randomized design according to Steel and Torrie (1991).

Result

Table 2: The effect of dietary fermented palm kernel cake (FPKC) on laying quail feed intake, egg production, egg mass, feed conversion, egg weight, and eggshell thickness

	FPKC in diet (%)					SEM
	0	5	10	15	20	
Feed intake (g/head/day)	20.37	21.61	21.71	21.47	22.30	0.43
Egg production (%)	71.80	71.61	70.80	72.05	70.93	2.01
Egg mass (g/head/day)	7.67	7.58	7.46	7.57	7.54	0.20
Feed conversion ratio	2.74	2.98	3.01	2.99	3.08	0.08
Egg weight (g)	10.69	10.61	11.71	10.48	10.65	0.50
Eggshell thickness (mm)	0.29	0.29	0.28	0.29	0.27	2.49

The feed intake of laying quail was non significant ($P>0.05$) by the levels of FPKC in the diets. The increase in level of FPKC in the diets did not reduce feed consumption of laying quail. The feed intake was ranging from 20.37-22.30 g/head/day (Table 2). The feed consumption of quail laying was non-significant ($P>0.05$) by the levels of FPKC in diets. The levels of FPKC in the diets were non-significant ($P>0.05$) on quail day egg production. The increase in levels of FPKC in quail diets maintaining egg production of laying quail. The egg production of quail laying of this experiment was ranging from 70.80 to 72.05%. Details of quail day egg production showed in Table 2.

The egg mass production of quail laying was non-significant ($P>0.05$) by the levels of FPKC in diets. The egg mass production of laying quail in this experiment was ranging from 7.46 to 7.67 g/head/day during the experiment. In accordance with the opinion of Abou El-Ghar and Debes (2013); Vercese et al (2012), stated that egg mass is linked to egg weight and egg production pattern. This result is similar to Nuraini et al (2012), which gained egg mass in the range of 6.85-7.20 g/head/day by administering a mixture of sago pulp and tofu waste fermented with *Neurospora crassa* to level of 12% in ration. Ciptaan et al (2020) findings quail eggs mass is much lower by 6.11 g/head/day with a palm oil sludge fermented with *Phanerochaete chrysosporium* and *N. crassa* to 25% level in quail ration.

The feed conversion of laying quail was non-significant ($P>0.05$) by levels of FPKC in diets was ranging from 2.74-3.08. Feed conversion is ratio between feed consumption and egg mass so that if treatment of FPKC given does not affect feed consumption and egg mass, then ration conversion is also relatively same. In Table 1 can be seen that quail got ration containing FPKC up to 20% level is equally efficient in utilize ration so it can produce egg with same ration conversion with control ration. This shows that quail is equally efficient in rations containing FPKC.

The egg weight of laying quail was non-significant ($P>0.05$) by different levels of FPKC in rations. The egg weight of laying quail in this experiment was ranging from 10.61 to 11.71 g/egg/head.

The thicknesses of eggshell from laying quail was non-significant ($P>0.05$) by the levels of FPKC in diets. The eggshell thickness of laying quail in this experiment was ranging from

0.27 to 0.29 mm. Eggshell thickness treatment of control to 20% level in ration still gives almost equal value.

Discussion

The difference between the treatments of R1 to R5 on feed consumption of laying quail rations indicates that giving FPKC with *S. rolfssii* to 20% (R5) has the same palatability. These are the rations containing FPKC which have better quality and aroma than original substrate. In accordance with opinion Mirnawati et al (2018) states that fermentation process can provide advantages physical and chemical profiles such as aroma, taste, and texture better than original substrate. It was also reported that the amount of feed consumed by poultry is influenced by feed palatability, digestibility, and diet composition (Mirnawati et al 2019; Mirnawati et al 2020; Ciptaan et al 2020). In this study, age, type and energy and protein intake for quail are relatively equal. Ciptaan et al (2020) obtaining a quail feed intake of around 22.14 g/head/day by inclusion of palm oil sludge fermented with *P. chrysosporium* and *N. crassa* to 25% in ration.

Non-significant effect of the production of quail eggs in this study due means consumption of rations associated with egg production. Daily quail production in all treatment were not affected because fermentation of palm kernel cake provides sufficient nutrient for poultry production. Fermentation can improve digestibility, which is in accordance Sukaryana et al (2010), Dairo and Fasuyi (2008), Mirnawati et al (2013) and Mirnawati et al (2019) that agriculture waste treated with fermentation will have promising nutrient profiles. Akbarillah et al (2010) and Fajrona et al (2020) that egg production is influenced by the amount of food consumed, especially the consumption of nutrients in addition to environmental factors. Consumption is what underlies the formation of eggs both quantity and quality. Most nutrient consumption will be converted into eggs, in addition to the basic needs in poultry. This result is higher than previous study conducted by Ciptaan et al (2020), which recorded daily egg production 60.21% with a palm oil sludge fermented by *N. crassa* to 12% level in ration. The study conducted by Abbas et al (2016) showed feed intake rate of Japanese quail (7 weeks of age) which had been supplemented with 15 g/kg *Cucurbita moschata* seeds oil over a period of 1-3 weeks at 135.5 g.

Ration conversion value from this study is higher than Nuraini et al (2012), which recorded from 2.82 to 2.90 with a mixture of sago pulp and tofu waste fermented with *N. crassa* to 12% in ration. While this number is lower when compared with feed conversion value from quail ration with palm kernel cake fermented with *Bacillus subtilis* in level 25% ration (Fajrona et al 2020). Another by-product is palm oil sludge which has potential to substitute conventional feed ingredients (e.g. yellow corn and soybean meal) in poultry rations. Where previous studies did not show a significant difference in terms of feed conversion ratio of broilers (Mirnawati et al 2021). However, these results were better than addition of 200-600 mg/kg L-Carnitine to Japanese quail diet in feed conversion ratio at a range of 5.8-7.7 (Mahmoud et al 2020).

A slight difference in quail egg weight caused fermentation process can breaks down complex or low digestibility substances into simpler molecule structures so that can enhance nutrient absorption as well as quality of an poultry products. On the other hand, during an incubation process. microbes produce primary and secondary metabolites that have beneficial properties. In addition, Mirnawati et al (2019) showed fermented palm kernel meal has better amino acids quality after fermentation. The egg weight obtained from this study was higher than previous results obtained by Fajrona et al (2020) were 9.57 - 9.64 g/egg/head and Nuraini et al (2012) were reported at 10.29 g/egg/head.

Combination of FPKC and humic acid addition became mineral source also increase bioavailability of calcium and phosphor, which is have important role for eggshell forming process. Enviromate (2002) stated that humic acid is a source of minerals and organic substances. Korsakov et al (2019) reported a significant effect on eggshell thickening (0.35 - 0.36 mm) of laying hens by administering humic acid (50-75 ml) via drinking water. Ciptaan et al (2020) obtained an average quail eggshell thickness of 0.26-0.28 mm. In addition Zita et al (2013), also states that average thickness of egg quail at 0.19 mm.

Conclusion

Based on result of this research, it can be concluded that palm kernel cake fermented with *Sclerotium rolfsii* can be used up to 20% level in quail rations. In terms of feed intake (22.30 g/head/day), egg production (70.93%), egg mass production (7.54 g/head/day), feed conversion (3.08), egg weight (10.65 g/egg/head), eggshell thickness (0.27 mm). It is hoped that palm kernel cake will be able to partially substitute commercial feed portions to enhance the profitability of quail layer farming.

Acknowledgement

This study was financially supported by BOPTN of Andalas University, number 42/UN.16.17/PP.RGB/LPPM/2019, dated April, 23rd 2019. I would like to thank all of those with whom I have had the pleasure to work during this project

Conflict of interest

All author declare that they have no conflict of interest.

References

Abbas R J, Alshaheen S A and Majeed T I 2017 Evaluation of the productive and physiological performance of Japanese quail (*Coturnix coturnix japonica*) fed different levels of pumpkin (*Cucurbita moschata*) seeds oil. International Journal of Veterinary Science Vol. 6: 31-35.

Abou El-Ghar R S and Debes A A 2013 Prediction of breeding values and selection for egg mass in a closed flock of white leghorn. *International Journal of Livestock Production Research* Vol. 1: 16-25.

Canellas L P and Olivares F L 2014 Physiological responses to humic substances as plant growth promoter. *Chemical and Biological Technologies in Agriculture* 1: 3. <https://doi.org/10.1186/2196-5641-1-3>

Ciptaan G, Mirnawati and Djulardi A 2020 Performance quality fed palm oil sludge fermented with *Phanerochaete cryosporium* and *Neurospora crassa*. *Livestock Research for Rural Development*, Volume 32, Article #128. Retrieved September 1, 2021, from <http://www.lrrd.org/lrrd32/8/gcpt32128.html>

Dairo F A S and Fauyi A O 2008 Evaluation of fermented palm kernel meal and fermented copra meal protein as substitute for soybean meal protein in laying hens diets. *Journal Central European Agriculture* Vol. 9: 35-44.

Enviromate 2002 Effect of humic acid on animal and humans (literature review and current research), Effects of humic acid, Enviromate Inc. 8571. Boat Club Road, Forth Worth, Texas 76719.

Fajrona K, Aini Q and Mirnawati 2020 The effect of fermented palm kernel cake with *Bacillus subtilis* on production performance and quail egg quality. *Quest Journal of Research in Agriculture and Animal Science* Vol. 7: 6-10.

Korsakov K V, Vasiliev A A, Sivokhina L A, Zabelina M V, Murtazaeva R N, Daeva T V and Kokorev V A 2019 The influence of humic acid supplement on the marketable properties of hy-line laying hen eggs. *Advances in Animal and Veterinary Sciences* Vol. 7: 66-70. <http://dx.doi.org/10.17582/journal.aavs/2019/7.s1.66.70>

Mahmoud R E, Elshopakey G E and Awadin W F 2020 Effects of feeding diets supplemented with different levels of l-carnitine on growth performance, serum metabolites, histopathological changes in growing japanese quails. *International Journal of Veterinary Science* Vol. 9: 16-23.

Meryandini A, Angreandari R and Rahmania N 2008 Isolasi bakteri mananolitik dan karakterisasi mananasenya. *Biota* Vol. 13: 82-88. <https://doi.org/10.24002/biota.v13i2.2675>

Mirnawati, Ciptaan G and Ferawati 2019 Improving the quality and nutrient content of palm kernel cake through fermentation with *Bacillus subtilis*. *Livestock Research for Rural Development*, Volume 31, Article #98. Retrieved September 2, 2021, from <http://www.lrrd.org/lrrd31/7/mirna31098.html>

Mirnawati, Ciptaan G and Ferawati 2020 Broiler performance on a diet containing palm kernel meal fermented with *Bacillus subtilis*. *Livestock Research for Rural Development*,

Volume 32, Article #21. Retrieved September 2, 2021, from <http://www.lrrd.org/lrrd32/2/mirna32021.html>

Mirnawati, Ciptaan G, Djulardi A and Makmur M 2021 Broiler response to the utilization of fermented palm oil sludge with *Phanerochaeta chrysosporium* and *Neurospora crassa*. International Journal of Veterinary Science. In press article. <https://doi.org/10.47278/journal.ijvs/2021.089>

Mirnawati, Djulardi A and Ciptaan G 2018 Utilization of fermented palm kernel cake with *Sclerotium rolfsii* in broiler ration. International Journal of Poultry Science Vol. 17 : 342-347. <https://dx.doi.org/10.3923/ijps.2018.342.347>

Mirnawati, Rizal Y and Marlida Y 2013 Effect of humic acid addition via drinking water on the performance of broiler fed diet containing fermented and non fermented palm kernel cake. Archiva Zootechnica Vol. 16: 41-53.

Nuraini, Sabrina and Latif S A 2012 Performances and quail egg quality feeding product fermented with *Neurospora crassa*. Indonesian Journal of Animal Science Vol. 14: 385-391. <https://doi.org/10.25077/jpi.14.2.385-391.2012>

Purwadaria T and Haryati T 2003 *In vitro* digestibility evaluation of coconut meal incorporated precipitate Beta D Manannase from *Eupenicillium javanicum*. Journal Microbiology Indonesia, Vol. 2003: 19 -21.

Razak N A 2006 Production and purification of mannanase degrading enzymes from palm kernel cake fermented by *Aspergillus niger* and *Sclerotium rolfsii*. Master thesis Universiti Putra Malaysia, Malaysia.

Rizal Y 2000 The response of broiler to the substitution part of soybean meal for palm kernel cake in the diet. Jurnal Peternakan Lingkungan Vo. 2: 15-20.

Steel R G D and Torrie J H 1991 Prinsip and Prosedur Statistik: Suatu pendekatan Biometrik PT. Gramedia Pustaka Utama. Jakarta, Indonesia.

Sukaryana Y, Atmomarsono U, Yuniyanto V D and Supriatna E 2010 Bioconversions of palm kernel cake and rice bran mixture by *Trichoderma viridae* toward nutritional contents. International Journal of Science and Engineering Vol. 1: 27 - 32. <http://dx.doi.org/10.12777/ijse.1.2.27-32>


Sundu B, Kumar A and Dingle J 2006 Palm kernel meal in broiler diets:effect on chicken performance and health. World's Poultry Science Journal Vol. 62 316-325. <https://doi.org/10.1079/WPS2005100>

Vercese F, Garcia E A, Sartori J R, Silva Ade P, Faitarone A B G, Berto D A, Molino A de B and Pelicia K 2012 Performance and egg quality of Japanese quails submitted to cyclic

heat stress. Brazilian Journal Poultry Science Vol. 14: 37-41. <https://doi.org/10.1590/S1516-635X2012000100007>


Zita L, Ledvinka Z and Klesalova L 2013 The effect of the age of Japanese quails on certain egg quality traits and their relationship. Veterinary Archives Vol. 83: 223-232.

1. Tanggal 09 September 2021: Pemberitahuan dan perbaikan hasil Turnitin paper



2. Review report: 1st (12 Oktober 2021)

Tanggal 12 Oktober 2021: Pemberitahuan review pertama (I) dari Reviewer



Gambar : Pemberitahuan dari editor tentang revisi pertama manuskrip

Manuskrip dari Reviewer

The Effect of Fermented Palm Kernel Cake ~~mixed with Humic Acid in~~ Layer Quail Rations on Production Performance and Eggshell Thickness

ABSTRACT

The purpose of the study was to evaluate how palm kernel cake fermented by *Sclerotium rolfsii* and ~~supplemented rationed~~ with humic acid affected production performance and ~~quail~~ egg quality of quails. The animals used in this study were 200 layer quail (*Coturnix coturnix japonica*) at 14 weeks of age. Fermented palm kernel cake (FPKC) with rationed compositions of 0%, 5%, 10%, 15%, and 20% were utilised in the research treatment, which lasted for eight weeks. Also, this study used a randomised design with four replicaties in each groupons. The results showed that feed intake, egg production, feed conversion, egg weight, and eggshell

thickness of layer quail were not statistically significant ($P>0.05$). Conclusively, palm kernel cake fermented with *S. rolfssii* ~~may can~~ be utilised in laying quail feed at a concentration of up to 20%.

Key words: Egg quality, Humic acid, Japanese quail, Palm kernel cake, *Sclerotium rolfssii*

INTRODUCTION

Palm kernel cake (PKC) is a by-product of palm oil processing, which can serve as a potential ingredient in poultry feed. Furthermore, its nutritional composition includes crude protein, crude fibre, crude fat, calcium, phosphorus at 16.07 %, 21.30 %, 8.23 %, 0.27 %, and 0.94 %, respectively, as well as copper at 48.4 ppm. (Mirnawati et al. 2010). The crude protein content of PKC is relatively high, yet its use in poultry rations is still limited. According to Rizal (2000), PKC at a concentration of up to 10 % can be used instead of 40 % soybean meal in broiler diet due to the high β -Manan content in the coarse fibres, which may be undesirable since birds do not have fibre-breaking enzymes for manan in the digestive tract (Sundu et al. 2006). Therefore, PKC must first be processed to improve its quality with the aid of fermentation biotechnology that utilises cellulolytic and mannanolytic moulds (Meryandini et al. 2008; Mirnawati et al. 2018; Purwadaria and Haryati., 2003). Furthermore, this can reduce the content of crude fibre and manan while increasing the quality of palm kernel cake such that it can replace the soybean meal in poultry rations.

Sclerotium rolfssii is a cellulolytic and mannanolytic microorganism that can be used for the fermentation of palm kernel cake. According to Razak et al. (2006), the mannanase enzyme activity of *S. rolfssii* is greater than that of *Aspergillus niger*. The fermentation of palm kernel cake with *S. rolfssii* yielded crude protein, retained nitrogen, crude fibre, and digestible crude fibre at 26.90%, 54.86%, 14.86%, and 58.41%, respectively, as well as crude fat at 0.22% and 2557.6 kcal/kg. However, the use of palm kernel cake in broiler diets is still restricted to 25% despite the rise in its nutritional content and quality.

Mirnawati et al. (2017) processed palm kernel cake mixed with humic acid through the fermentation process using *S. rolfssii*. The result of this study showed an increase in crude protein, nitrogen retention, and crude fibre digestibility at 27.43 %, 59.17 %, and 55.40 %, respectively.

Formatted: Font: Italic

Formatted: Font: Italic

respectively, as well as a decrease in crude fibre at 11.53 %. After fermentation, the increased nutritional content of palm kernels enables its use as a quail feed ingredient. Therefore, it is necessary to conduct research in order to determine the effect of fermented palm kernel cake containing *S. rolf sii* in rations on the production performance and quality of laying quails.

MATERIALS AND METHODS

The samples used in this study ~~we~~are 200 quail laying hens aged about 14-weeks old, which were confined in individual cages of size 45 × 20 × 30 cm as ten laying birds per unit. The study used a fully randomized design (CRD) with five treatments containing 0, 5, 10, 15, and 20% compositions of FPKC, as well as four ~~replicates~~ ~~duplicate~~s each. The diets used included iso-protein and iso-caloric at 20% and 2700 kcal/kg, respectively. Subsequently, Table 1 ~~showe~~d the diet formulation, nutritional and metabolizable energy levels of treatment diets. The diet formulation was made up of yellow corn, rice bran, meat meal, CP 126 concentrate feed (Charoen Pokphand Indonesia), top mix and FPKC. In addition, drinking water and experimental diet were provided *ad-libitum*.

The fermented palm kernel cake was made using a combination of PKC and rice bran at 80 % and 20 %, respectively, which were fermented with *S. rolf sii* and added to 200 ppm humic acid. The inoculum dosage was 10% of the substrate, and the incubation period was seven days. After harvesting, the product is dried and milled before being incorporated into quail diets. Meanwhile, layer quail have a feeding period of two months or eight weeks. Table 1 ~~showe~~d the composition of the feeding or diet treatments.

Data Collection

The data collected during the study included ~~d~~ feed conversion, egg mass production (g/head/day), egg weight (g/egg/head), feed intake (g/head/day), quail day egg production (%) and the eggshell thickness (mm) of laying quail, which were measured following Nuraini et al. (2012).

Data Analysis

All data were analyzed by analysis of variance based on a completely randomized design according to Steel and Torrie (1991).

Formatted: Font: Italic

RESULTS AND DISCUSSION

Separate Results and Discussion (follow the instructions for authors)

There was no significant difference ($P>0.05$) in the feed intake of the laying quails based on the levels of FPKC in the diets since an increase in the level of FPKC did not reduce the feed consumption of the laying quails. Table 2 showed the feed intake, which ranged between 20.37-22.30 g/head/day. Similarly, there was no significant difference ($P>0.05$) in the feed consumption of the laying quails based on the levels of FPKC in the diets. The difference in feed consumption of laying quail rations between treatments R1 and R5 suggests that feeding FPKC with *S. rolfssii* to 20% (R5) has the same palatability. However, these FPKC-containing meals were discovered to have a higher quality and aroma than the original substrate. According to Mirnawati et al. (2018), the fermentation process can improve the physical and chemical characteristics such as aroma, taste, and texture compared to the original substrate. Furthermore, it was also observed that feed palatability, digestibility, and diet composition all affect the amount of feed eaten by birds (Mirnawati et al. 2019; Mirnawati et al. 2020; Ciptaan et al. 2020). In this study, parameters such as age, type, as well as energy and protein consumption were all relatively equal. Ciptaan et al. (2020) obtained a quail feed intake of about 22.14 g/head/day by adding 25 % palm oil sludge fermented with *Phanerochaeta chrysosporium* and *Neurospora crassa* in rations. Palm oil sludge is another promising by-product that can replace the ingredients of standard feed, such as yellow maize and soybean meal, in poultry diets. Previous research did not reveal a significant difference in the feed conversion ratio of broilers (Mirnawati et al. 2021).

The effects of the amount of FPKC in the diet on the daily egg production of quails were not significant ($P>0.05$). However, increased amounts of FPKC in quail diets can sustain the egg production of laying quails. The egg production of laying quails in this experiment ranged between 70.80 to 72.05 %, as shown in Table 2. The daily quail production was unaffected in treatments R1, R2, R3, R4, and R5 since fermented palm kernel cake provides enough nutrients needed for poultry production. Subsequently, fermentation can increase digestibility, which is in line with the hypothesis of Sukaryana et al. (2010), Dairo and Fasuyi (2008), Mirnawati et al. (2013), and Mirnawati et al. (2019) that fermented farm waste will

have favourable nutritional profiles. Meanwhile, there was no significant difference in the egg production of quails since it was associated with the consumption of rations. This is consistent with the study by Akbarillah et al. (2010) and Fajrona et al. (2020) that egg production is controlled by the amount of food ingested, particularly nutrient consumption, as well as environmental variables. Consumption improves the quantity and quality of egg production since a majority of the nutrients consumed will be transformed into eggs, in addition to the fundamental needs of the birds. This value obtained in this result is greater than what was obtained in the previous study by Ciptaan et al. (2020), which reported daily egg production of 60.21 % using palm oil sludge fermented with *N. crassa* at a ration level of 12 %. According to a study by Abbas et al. (2016), the rate of feed intake Japanese quails at seven weeks of age, supplemented with 15 g/kg *Cucurbita moschata* seeds oil over a period of 1-3 weeks, was 135.5 g.

The quantity of FPKC in meals did not affect egg mass production of laying quails ($P>0.05$). During the trial, the egg mass output of laying quails ranged from 7.46 to 7.67 g/head/day. According to Abou El-Ghar and Debes (2013) and Vercese et al. (2012), egg mass is related to egg weight and egg production pattern. This is comparable to the results obtained by Nuraini et al. (2012), who obtained an egg mass in the range of 6.85-7.20 g/head/day by administering a mixture of sago pulp and tofu waste fermented with *N. crassa* at a 12 % ration. According to Ciptaan et al. (2020), the quail egg mass was reduced by 6.11 g/head/day when palm oil sludge fermented with *P. chrysosporium* and *N. crassa*, was incorporated to quail feed at a 25 % ration level.

Feed conversion is the ratio of feed intake to egg mass, and it was non-significant ($P>0.05$) when FPKC levels in diets ranged between 2.74-3.08. Therefore, since the FPKC treatment has no effect on feed consumption or egg mass, the ration conversion is also relatively the same. Table 1 shows that quails fed a ration containing up to 20% FPKC are similarly efficient in egg production to quails fed with the control ration (R1), which demonstrates that quails are equally efficient in FPKC-containing diets.

The result of this study is greater than that obtained by Nuraini et al. (2012), which utilised ration conversions ranging from 2.82 to 2.90 with a mixture of sago pulp and tofu pulp fermented with *N. crassa* 12% in rations. This figure is lower when compared to the feed conversion value from quail ration with palm kernel cake fermented with *Bacillus*

subtilis at a 25% ration level. (Fajrona et al. 2020). However, The results were more desirable than what was obtained from a mixture of 200-600 mg/kg L-Carnitine and Japanese quail diet with a feed conversion ratio ranging between 5.8 to 7.7 (Mahmoud et al. 2020).

Different amounts of FPKC in diets had no effect on the egg weight of laying quail ($P>0.05$), which ranged from 10.61 to 11.71 g/egg/head. A minor variation in quail egg weight produced by the fermentation process might break down complex or low digestible components into simpler molecular structures, improving nutritional absorption and the quality of poultry products. Conversely, beneficial primary and secondary metabolites are secreted by microbes throughout the incubation process. Furthermore, Mirnawati et al. (2019) showed that fermented palm kernel meal has higher amino acid quality after fermentation. The egg weight obtained from this study was higher than previous results obtained by Fajrona et al. (2020) and Nuraini et al. (2012), which were 9.57 - 9.64 g/egg/head and 10.29 g/egg/head, respectively.

The amounts of FPKC in meals had no effect on the thickness of the eggshells from laying quails ($P>0.05$), which ranged from 0.27 to 0.29 mm. The eggshell thickness treatment of R1 to R5 showed that FPKC up to 20% in the diet still provides almost similar results because the inclusion of FPKC and humic acid as mineral sources increases the bioavailability of calcium and phosphorus, both of which play significant roles in the eggshell formation process. According to Enviromate (2002), humic acid is a source of minerals and organic compounds. Also, Korsakov et al. (2019) found that about 50-75 ml of humic acid given through drinking water significantly affects the eggshell thickness, which was 0.35 - 0.36 mm in laying hens. Ciptaan et al. (2020) measured the average thickness of quail eggshells to be 0.26-0.28 mm. Additionally, Zita et al. (2013), also reported that the average thickness of quail eggs was 0.19 mm.

Conclusion

Conclusively, palm kernel cake fermented with *Sclerotium rolfsii* can be utilized up to 20% in quail diets. The results showed that the feed intake, egg production, egg mass production, feed conversion, egg weight, and eggshell thickness were found to be 22.30 g/head/day, 70.93%, 7.54 g/head/day, 3.08, 10.65 g/egg/head, and 0.27 mm, respectively. Therefore, it is

Dairo FAS and Fauyi AO, 2008. Evaluation of fermented palm kernel meal and fermented copra meal protein as substitute for soybean meal protein in laying hens diets. Journal Central European Agriculture 9: 35-44. [DOI no missing](#)

Formatted: English (United States)

Enviromate, 2002. Effect of humic acid on animal and humans (literature review and current research), Effects of humic acid, Enviromate Inc. 8571. Boat Club Road, Forth Worth, Texas 76719. [DOI no missing](#)

Formatted: English (United States)

Fajrona K, Aini Q and Mirnawati, 2020. The effect of fermented palm kernel cake with *bacillus subtilisin* rations on production performance and quail egg quality. Quest Journal of Research in Agriculture and Animal Science 7: 6-10. [DOI no missing](#)

Korsakov KV, Vasiliev AA, Sivokhina LA, Zabelina MV, Murtazaeva RN, Daeva TV and Kokorev VA, 2019. The influence of humic acid supplement on the marketable properties of hy-line laying hen eggs. Advances in Animal and Veterinary Sciences 7: 66-70. <http://dx.doi.org/10.17582/journal.aavs/2019/7.s1.66.70>

Mahmoud RE, Elshopakey GE and Awadin WF, 2020. Effects of feeding diets supplemented with different levels of l-carnitine on growth performance, serum metabolites, histopathological changes in growing japanese quails. International Journal of Veterinary Science 9: 16-23. [DOI no missing](#)

Meryandini A, Angreandari R and Rahmania N, 2008. Isolasi bakteri mananolitik dan karakterisasi mananasenya. Biota 13: 82-88. <https://doi.org/10.24002/biota.v13i2.2675>

Mirnawati, Ciptaan G and Ferawati, 2019. Improving the quality and nutrient content of palm kernel cake through fermentation with *Bacillus subtilis*. Livestock Research for Rural Development 31: 119-123. [DOI no missing](#)

Formatted: English (United States)

Mirawati, Ciptaan G and Ferawati, 2020. Broiler performance on a diet containing palm kernel meal fermented with *Bacillus subtilis*. Livestock Research for Rural Development 32: 2020. [DOI no missing](#)

Mirawati, Ciptaan G, Djulardi A and Makmur M, 2021. Broiler response to the utilization of fermented palm oil sludge with *Phanerochaeta chrysosporium* and *Neurospora crassa*. International Journal of Veterinary Science. In press article.
<https://doi.org/10.47278/journal.ijvs/2021.089>

Mirawati, Djulardi A and Ciptaan G, 2018. Utilization of fermented palm kernel cake with *Sclerotium rolfsii* in broiler ration. International Journal of Poultry Science 17 : 342-347. <https://dx.doi.org/10.3923/ijps.2018.342.347>

Mirawati, Rizal Y and Marlida Y, 2013. Effect of humic acid addition via drinking water on the performance of broiler fed diet containing fermented and non fermented palm kernel cake. Archiva Zootechnica 16: 41-53. [DOI no missing](#)

Formatted: English (United States)

Nuraini, Sabrina and Latif SA, 2012. Performances and quail egg quality feeding product fermented with *Neurospora crassa*. Indonesian Journal of Animal Science 14: 385-391. <https://doi.org/10.25077/jpi.14.2.385-391.2012>

Purwadaria T and Haryati T, 2003. *In vitro* digestibility evaluation of coconut meal incorporated precipitate Beta D Mannanase from *Eupenicillium javanicum*. Journal Microbiology Indonesia, 2003: 19 -21. [DOI no missing](#)

Razak NA, 2006. Production and purification of mannanase degrading enzymes from palm kernel cake fermented by *Aspergillus niger* and *Sclerotium rolfsii*. Master thesis Universiti Putra Malaysia, Malaysia. [DOI no missing](#)

Formatted: English (United States)

Rizal Y, 2000. The response of broiler to the substitution part of soybean meal for palm kernel cake in the diet. Jurnal Peternakan Lingkungan 2: 15-20.

Steel RGD and Torrie JH, 1991. Prinsip and Prosedur Statistik: Suatu pendekatan Biometrik PT. Gramedia Pustaka Utama. Jakarta, Indonesia. [DOI no missing](#)

Formatted: English (United States)

Sukaryana Y, Atmomarsono U, Yuniarto VD and Supriatna E, 2010. Bioconversions of palm kernel cake and rice bran mixture by *Trichoderma viridae* toward nutritional contents. International Journal of Science and Engineering 1: 27 - 32. <http://dx.doi.org/10.12777/ijse.1.2.27-32>

Sundu B, Kumar A and Dingle J, 2006 Palm kernel meal in broiler diets:effect on chicken performance and health. World's Poultry Science Journal 62 316-325. <https://doi.org/10.1079/WPS2005100>

Vercese F, Garcia EA, Sartori JR, Silva Ade P, Faitarone ABG, Berto DA, Molino A de B and Pelícia K, 2012. Performance and egg quality of Japanese quails submitted to cyclic heat stress. Brazilian Journal Poultry Science 14: 37-41. <https://doi.org/10.1590/S1516-635X2012000100007>

Zita L, Ledvinka Z and Klesalova L, 2013. The effect of the age of Japanese quails on certain egg quality traitsand their relationship. Veterinary Archives. 83: 223-232. [DOI no missing](#)

Formatted: English (United States)

Table 1: Diet formulation (%), nutrient content (%) and metabolizable energy (kcal/kg).

Item	Treatment Ration				
	R1	R2	R3	R4	R5
Corn	45.5	45	44	43	43
Rice brand	20	18	16	13.5	11.5
Meat meal	6	6	6	6	6
CP 126 concentrate	26.5	24	22	20.5	17.5

FPKC	0	5	10	15	20
Mineral B12	1	1	1	1	1
Top mix	1	1	1	1	1
Total (%)	100	100	100	100	100
<hr/>					
Crude protein	20.06	20.04	20.16	20.44	20.42
Crude fat	3.46	3.40	3.34	3.28	3.29
Crude fiber	6.13	6.17	6.23	6.27	6.50
Calcium	2.94	2.80	2.63	2.60	1.94
Phosphor	0.88	0.89	0.91	0.93	0.82
Metabolizable energy	2710	2708	2703	2706	2725
<hr/>					


Table 2: The effect of dietary fermented palm kernel cake (FPKC) on laying quail feed intake, egg production, egg mass, feed conversion, egg weight, and eggshell thickness.

Variables	Treatment (%)					SEM
	R1	R2	R3	R4	R5	
Feed intake (g/head/day)	20.37	21.61	21.71	21.47	22.30	0.43
Egg production (%)	71.80	71.61	70.80	72.05	70.93	2.01
Egg mass (g/head/day)	7.67	7.58	7.46	7.57	7.54	0.20
Feed conversion ratio	2.74	2.98	3.01	2.99	3.08	0.08
Egg weight (g)	10.69	10.61	11.71	10.48	10.65	0.50
Eggshell thickness (mm)	0.29	0.29	0.28	0.29	0.27	2.49

Note: Inclusion FPKC in R1 to R5 was R1 (0%-Control), R2 (5%), R3 (10%), R4 (15%), and R5(20%). SEM: standard error of the mean.

3. Revised version: 1st (13 Oktober 2021)

Tanggal 13 Oktober 2021: Penulis mengirim hasil perbaikan I (a revise author version)



MANUSKRIP PERBAIKAN REVISI 1

The Effect of Fermented Palm Kernel Cake Layer Quail Rations on Production Performance and Eggshell Thickness

ABSTRACT

The purpose of the study was to evaluate how palm kernel cake fermented by *Sclerotium rolfsii* and supplemented with humic acid affected production performance and egg quality of quails. The animals used in this study were 200 layer quail (*Coturnix coturnix japonica*) at 14 weeks of age. Fermented palm kernel cake (FPKC) with rationed compositions of 0%, 5%, 10%, 15%, and 20% were utilised in the research treatment, which lasted for eight weeks. Also, this study used a randomised design with four replicates in each group. The results showed that feed intake, egg production, feed conversion, egg weight, and eggshell thickness of layer quail were

not statistically significant ($P>0.05$). Conclusively, palm kernel cake fermented with *S. rolfsii* may be utilised in laying quail feed at a concentration of up to 20%.

Key words: Egg quality, Humic acid, Japanese quail, Palm kernel cake, *Sclerotium rolfsii*

INTRODUCTION

Palm kernel cake (PKC) is a by-product of palm oil processing, which can serve as a potential ingredient in poultry feed. Furthermore, its nutritional composition includes crude protein, crude fibre, crude fat, calcium, phosphorus at 16.07 %, 21.30 %, 8.23 %, 0.27 %, and 0.94 %, respectively, as well as copper at 48.4 ppm. (Mirnawati et al. 2010). The crude protein content of PKC is relatively high, yet its use in poultry rations is still limited. According to Rizal (2000), PKC at a concentration of up to 10 % can be used instead of 40 % soybean meal in broiler diet due to the high β -Manan content in the coarse fibres, which may be undesirable since birds do not have fibre-breaking enzymes for manan in the digestive tract (Sundu et al. 2006). Therefore, PKC must first be processed to improve its quality with the aid of fermentation biotechnology that utilises cellulolytic and mannanolytic moulds (Meryandini et al. 2008; Mirnawati et al. 2018; Purwadaria and Haryati., 2003).

Furthermore, this can reduce the content of crude fibre and manan while increasing the quality of palm kernel cake such that it can replace the soybean meal in poultry rations.

Sclerotium rolfsii is a cellulolytic and mannanolytic microorganism that can be used for the fermentation of palm kernel cake. According to Razak et al. (2006), the mannanase enzyme activity of *S. rolfsii* is greater than that of *Aspergillus niger*. The fermentation of palm kernel cake with *S. rolfsii* yielded crude protein, retained nitrogen, crude fibre, and digestible crude fibre at 26.90%, 54.86%, 14.86%, and 58.41%, respectively, as well as crude fat at 0.22% and 2557.6 kcal/kg. However, the use of palm kernel cake in broiler diets is still restricted to 25% despite the rise in its nutritional content and quality.

Mirnawati et al. (2017) processed palm kernel cake mixed with humic acid through the fermentation process using *S. rolfsii*. The result of this study showed an increase in crude protein, nitrogen retention, and crude fibre digestibility at 27.43 %, 59.17 %, and 55.40 %, respectively, as well as a decrease in crude fibre at 11.53 %. After fermentation, the

increased nutritional content of palm kernels enables its use as a quail feed ingredient. Therefore, it is necessary to conduct research in order to determine the effect of fermented palm kernel cake containing *S. rolfssii* in rations on the production performance and quality of laying quails.

MATERIALS AND METHODS

The samples used in this study were 200 quail laying hens aged about 14-weeks old, which were confined in individual cages of size 45 × 20 × 30 cm as ten laying birds per unit. The study used a fully randomized design (CRD) with five treatments containing 0, 5, 10, 15, and 20% compositions of FPKC, as well as four replicates each. The diets used included iso-protein and iso-caloric at 20% and 2700 kcal/kg, respectively. Subsequently, Table 1 showed the diet formulation, nutritional and metabolizable energy levels of treatment diets. The diet formulation was made up of yellow corn, rice bran, meat meal, CP 126 concentrate feed (Charoen Pokphand Indonesia), top mix and FPKC. In addition, drinking water and experimental diet were provided *ad-libitum*.

The fermented palm kernel cake was made using a combination of PKC and rice bran at 80 % and 20 %, respectively, which were fermented with *S. rolfssii* and added to 200 ppm humic acid. The inoculum dosage was 10% of the substrate, and the incubation period was seven days. After harvesting, the product is dried and milled before being incorporated into quail diets. Meanwhile, layer quail have a feeding period of two months or eight weeks. Table 1 showed the composition of the feeding or diet treatments.

Data Collection

The data collected during the study included feed conversion, egg mass production (g/head/day), egg weight (g/egg/head), feed intake (g/head/day), quail day egg production (%) and the eggshell thickness (mm) of laying quail, which were measured following Nuraini et al. (2012).

Data Analysis

All data were analyzed by analysis of variance based on a completely randomized design according to Steel and Torrie (1991).

RESULTS

There was no significant difference ($P>0.05$) in the feed intake of the laying quails based on the levels of FPKC in the diets since an increase in the level of FPKC did not reduce the feed consumption of the laying quails. Table 2 showed the feed intake, which ranged between 20.37-22.30 g/head/day. Similarly, there was no significant difference ($P>0.05$) in the feed consumption of the laying quails based on the levels of FPKC in the diets. The effects of the amount of FPKC in the diet on the daily egg production of quails were not significant ($P>0.05$). However, increased amounts of FPKC in quail diets can sustain the egg production of laying quails. The egg production of laying quails in this experiment ranged between 70.80 to 72.05 %, as shown in Table 2.

The quantity of FPKC in meals did not affect egg mass production of laying quails ($P>0.05$). During the trial, the egg mass output of laying quails ranged from 7.46 to 7.67 g/head/day. Feed conversion is the ratio of feed intake to egg mass, and it was non-significant ($P>0.05$) when FPKC levels in diets ranged between 2.74-3.08. Different amounts of FPKC in diets had no effect on the egg weight of laying quail ($P>0.05$), which ranged from 10.61 to 11.71 g/egg/head. The amounts of FPKC in meals had no effect on the thickness of the eggshells from laying quails ($P>0.05$), which ranged from 0.27 to 0.29 mm.

DISCUSSION

The difference in feed consumption of laying quail rations between treatments R1 and R5 suggests that feeding FPKC with *S. rolf sii* to 20% (R5) has the same palatability. However, these FPKC-containing meals were discovered to have a higher quality and aroma than the original substrate. According to Mirnawati et al. (2018), the fermentation process can improve the physical and chemical characteristics such as aroma, taste, and texture compared to the original substrate. Furthermore, it was also observed that feed palatability, digestibility, and diet composition all affect the amount of feed eaten by birds (Mirnawati et al. 2019; Mirnawati et al. 2020; Ciptaan et al. 2020). In this study, parameters such as age, type, as well as energy and protein consumption were all relatively equal. Ciptaan et al. (2020) obtained a quail feed intake of about 22.14 g/head/day by adding 25 % palm oil

sludge fermented with *Phanerochaeta chrysosporium* and *Neurospora crassa* in rations. Palm oil sludge is another promising by-product that can replace the ingredients of standard feed, such as yellow maize and soybean meal, in poultry diets. Previous research did not reveal a significant difference in the feed conversion ratio of broilers (Mirnawati et al. 2021).

The daily quail production was unaffected in treatments R1, R2, R3, R4, and R5 since fermented palm kernel cake provides enough nutrients needed for poultry production. Subsequently, fermentation can increase digestibility, which is in line with the hypothesis of Sukaryana et al. (2010), Dairo and Fasuyi (2008), Mirnawati et al. (2013), and Mirnawati et al. (2019) that fermented farm waste will have favourable nutritional profiles. Meanwhile, there was no significant difference in the egg production of quails since it was associated with the consumption of rations. This is consistent with the study by Akbarillah et al. (2010) and Fajrona et al. (2020) that egg production is controlled by the amount of food ingested, particularly nutrient consumption, as well as environmental variables. Consumption improves the quantity and quality of egg production since a majority of the nutrients consumed will be transformed into eggs, in addition to the fundamental needs of the birds. This value obtained in this result is greater than what was obtained in the previous study by Ciptaan et al. (2020), which reported daily egg production of 60.21 % using palm oil sludge fermented with *N. crassa* at a ration level of 12 %. According to a study by Abbas et al. (2016), the rate of feed intake Japanese quails at seven weeks of age, supplemented with 15 g/kg *Cucurbita moschata* seeds oil over a period of 1-3 weeks, was 135.5 g.

According to Abou El-Ghar and Debes (2013) and Vercese et al. (2012), egg mass is related to egg weight and egg production pattern. This is comparable to the results obtained by Nuraini et al. (2012), who obtained an egg mass in the range of 6.85-7.20 g/head/day by administering a mixture of sago pulp and tofu waste fermented with *N. crassa* at a 12 % ration. According to Ciptaan et al. (2020), the quail egg mass was reduced by 6.11 g/head/day when palm oil sludge fermented with *P. chrysosporium* and *N. crassa*, was incorporated to quail feed at a 25 % ration level.

The FPKC treatment has no effect on feed consumption or egg mass, the ration conversion is also relatively the same. Table 1 shows that quails fed a ration containing up to 20% FPKC are similarly efficient in egg production to quails fed with the control ration (R1), which demonstrates that quails are equally efficient in FPKC-containing diets.

The result of this study is greater than that obtained by Nuraini et al. (2012), which utilised ration conversions ranging from 2.82 to 2.90 with a mixture of sago pulp and tofu pulp fermented with *N. crassa* 12% in rations. This figure is lower when compared to the feed conversion value from quail ration with palm kernel cake fermented with *Bacillus subtilis* at a 25% ration level. (Fajrona et al. 2020). However, The results were more desirable than what was obtained from a mixture of 200-600 mg/kg L-Carnitine and Japanese quail diet with a feed conversion ratio ranging between 5.8 to 7.7 (Mahmoud et al. 2020).

A minor variation in quail egg weight produced by the fermentation process might break down complex or low digestible components into simpler molecular structures, improving nutritional absorption and the quality of poultry products. Conversely, beneficial primary and secondary metabolites are secreted by microbes throughout the incubation process. Furthermore, Mirnawati et al. (2019) showed that fermented palm kernel meal has higher amino acid quality after fermentation. The egg weight obtained from this study was higher than previous results obtained by Fajrona et al. (2020) and Nuraini et al. (2012), which were 9.57 - 9.64 g/egg/head and 10.29 g/egg/head, respectively.

The eggshell thickness treatment of R1 to R5 showed that FPKC up to 20% in the diet still provides almost similar results because the inclusion of FPKC and humic acid as mineral sources increases the bioavailability of calcium and phosphorus, both of which play significant roles in the eggshell formation process. According to Enviromate (2002), humic acid is a source of minerals and organic compounds. Also, Korsakov et al. (2019) found that about 50-75 ml of humic acid given through drinking water significantly affects the eggshell thickness, which was 0.35 - 0.36 mm in laying hens. Ciptaan et al. (2020) measured the average thickness of quail eggshells to be 0.26-0.28 mm. Additionally, Zita et al. (2013), also reported that the average thickness of quail eggs was 0.19 mm.

Conclusion

Conclusively, palm kernel cake fermented with *Sclerotium rolfsii* can be utilized up to 20% in quail diets. The results showed that the feed intake, egg production, egg mass production, feed conversion, egg weight, and eggshell thickness were found to be 22.30 g/head/day, 70.93%, 7.54 g/head/day, 3.08, 10.65 g/egg/head, and 0.27 mm, respectively. Therefore, it is

expected that palm kernel cake would be able to partially replace the current ingredients used in commercial feed in order to enhance the profitability of quail layer farming.

Acknowledgements

This study was funded by the BOPTN of Andalas University, number 42/UN.16.17/PP.RGB/LPPM/2019, on the 23rd of April, 2019. I am grateful to everyone with whom I had the pleasure of working on this project.

Author's Contribution

Gita Ciptaan supervised the experiment and writing original manuscript. Mirnawati and Ferawati conducted the experiment and analyzed the data. Malik Makmur finalize manuscript.

REFERENCES

- Abbas RJ, Alshaheen SA and Majeed TI, 2017. Evaluation of the productive and physiological performance of Japanese quail (*Coturnix coturnix japonica*) fed different levels of pumpkin (*Cucurbita moschata*) seeds oil. *International Journal of Veterinary Science* 6: 31-35.
- Abou El-Ghar RS and Debes AA, 2013. Prediction of breeding values and selection for egg mass in a closed flock of white leghorn. *International Journal of Livestock Production Research* 1:16-25.
- Canellas LP and Olivares, FL, 2014. Physiological responses to humic substances as plant growth promoter. *Chemical and Biological Technologies in Agriculture* 1: 3.
<https://doi.org/10.1186/2196-5641-1-3>
- Ciptaan G, Mirnawati and Djulardi A, 2020. Performance quality feed palm oil sludge fermented with *Phanerochaete cryosporium* and *Neurospora crassa*. *Livestock Reasearch for Rural Development* 32: 8.

- Dairo FAS and Fauyi AO, 2008. Evaluation of fermented palm kernel meal and fermented copra meal protein as substitute for soybean meal protein in laying hens diets. *Journal Central European Agriculture* 9: 35-44.
- Enviromate, 2002. Effect of humic acid on animal and humans (literature review and current research), Effects of humic acid, Enviromate Inc. 8571. Boat Club Road, Forth Worth, Texas 76719.
- Fajrona K, Aini Q and Mirnawati, 2020. The effect of fermented palm kernel cake with *Bacillus subtilis* on production performance and quail egg quality. *Quest Journal of Research in Agriculture and Animal Science* 7: 6-10.
- Korsakov KV, Vasiliev AA, Sivokhina LA, Zabelina MV, Murtazaeva RN, Daeva TV and Kokorev VA, 2019. The influence of humic acid supplement on the marketable properties of hy-line laying hen eggs. *Advances in Animal and Veterinary Sciences* 7: 66-70.
<http://dx.doi.org/10.17582/journal.aavs/2019/7.s1.66.70>
- Mahmoud RE, Elshopakey GE and Awadin WF, 2020. Effects of feeding diets supplemented with different levels of l-carnitine on growth performance, serum metabolites, histopathological changes in growing japanese quails. *International Journal of Veterinary Science* 9: 16-23.
- Meryandini A, Angreandari R and Rahmania N, 2008. Isolasi bakteri mananolitik dan karakterisasi mananasenya. *Biota* 13: 82-88.
<https://doi.org/10.24002/biota.v13i2.2675>
- Mirnawati, Ciptaan G and Ferawati, 2019. Improving the quality and nutrient content of palm kernel cake through fermentation with *Bacillus subtilis*. *Livestock Research for Rural Development* 31: 119-123.
- Mirnawati, Ciptaan G and Ferawati, 2020. Broiler performance on a diet containing palm kernel meal fermented with *Bacillus subtilis*. *Livestock Research for Rural Development* 32: 2020.
- Mirnawati, Ciptaan G, Djulardi A and Makmur M, 2021. Broiler response to the utilization of fermented palm oil sludge with *Phanerochaeta chrysosporium* and *Neurospora*

crassa. International Journal of Veterinary Science. In press article.

<https://doi.org/10.47278/journal.ijvs/2021.089>

Mirawati, Djulardi A and Ciptaan G, 2018. Utilization of fermented palm kernel cake with *Sclerotium rolfsii* in broiler ration. International Journal of Poultry Science 17 : 342-347. <https://dx.doi.org/10.3923/ijps.2018.342.347>

Mirawati, Rizal Y and Marlida Y, 2013. Effect of humic acid addition via drinking water on the performance of broiler fed diet containing fermented and non fermented palm kernel cake. Archiva Zootechnica 16: 41-53.

Nuraini, Sabrina and Latif SA, 2012. Performances and quail egg quality feeding product fermented with *Neurospora crassa*. Indonesian Journal of Animal Science 14: 385-391. <https://doi.org/10.25077/jpi.14.2.385-391.2012>

Purwadaria T and Haryati T, 2003. *In vitro* digestibility evaluation of coconut meal incorporated precipitate Beta D Manannase from *Eupenicillium javanicum*. Journal Microbiology Indonesia, 2003: 19 -21.

Razak NA, 2006. Production and purification of mannanase degrading enzymes from palm kernel cake fermented by *Aspergillus niger* and *Sclerotium rolfsii*. Master thesis Universiti Putra Malaysia, Malaysia.

Rizal Y, 2000. The response of broiler to the substitution part of soybean meal for palm kernel cake in the diet. Jurnal Peternakan Lingkungan 2: 15-20.

Steel RGD and Torrie JH, 1991. Prinsip and Prosedur Statistik: Suatu pendekatan Biometrik PT. Gramedia Pustaka Utama. Jakarta, Indonesia.

Sukaryana Y, Atmomarsono U, Yuniarto VD and Supriatna E, 2010. Bioconversions of palm kernel cake and rice bran mixture by *Trichoderma viridae* toward nutritional contents. International Journal of Science and Engineering 1: 27 - 32. <http://dx.doi.org/10.12777/ijse.1.2.27-32>

Sundu B, Kumar A and Dingle J, 2006 Palm kernel meal in broiler diets:effect on chicken performance and health. World's Poultry Science Journal 62 316-325. <https://doi.org/10.1079/WPS2005100>

Vercese F, Garcia EA, Sartori JR, Silva Ade P, Faltarone ABG, Berto DA, Molino A de B and Pelícia K, 2012. Performance and egg quality of Japanese quails submitted to cyclic heat stress. *Brazilian Journal Poultry Science* 14: 37-41.

<https://doi.org/10.1590/S1516-635X2012000100007>

Zita L, Ledvinka Z and Klesalova L, 2013. The effect of the age of Japanese quails on certain egg quality traits and their relationship. *Veterinary Archives*. 83: 223-232.

Table 1: Diet formulation (%), nutrient content (%) and metabolizable energy (kcal/kg).

Item	Treatment Ration				
	R1	R2	R3	R4	R5
Corn	45.5	45	44	43	43
Rice brand	20	18	16	13.5	11.5
Meat meal	6	6	6	6	6
CP 126 concentrate	26.5	24	22	20.5	17.5
FPKC	0	5	10	15	20
Mineral B12	1	1	1	1	1
Top mix	1	1	1	1	1
Total (%)	100	100	100	100	100
Crude protein	20.06	20.04	20.16	20.44	20.42
Crude fat	3.46	3.40	3.34	3.28	3.29
Crude fiber	6.13	6.17	6.23	6.27	6.50
Calcium	2.94	2.80	2.63	2.60	1.94
Phosphor	0.88	0.89	0.91	0.93	0.82
Metabolizable energy	2710	2708	2703	2706	2725

Table 2: The effect of dietary fermented palm kernel cake (FPKC) on laying quail feed intake, egg production, egg mass, feed conversion, egg weight, and eggshell thickness.

Variables	Treatment (%)					SEM
	R1	R2	R3	R4	R5	
Feed intake (g/head/day)	20.37	21.61	21.71	21.47	22.30	0.43
Egg production (%)	71.80	71.61	70.80	72.05	70.93	2.01
Egg mass (g/head/day)	7.67	7.58	7.46	7.57	7.54	0.20
Feed conversion ratio	2.74	2.98	3.01	2.99	3.08	0.08
Egg weight (g)	10.69	10.61	11.71	10.48	10.65	0.50
Eggshell thickness (mm)	0.29	0.29	0.28	0.29	0.27	2.49

Note: Inclusion FPKC in R1 to R5 was R1 (0%-Control), R2 (5%), R3 (10%), R4 (15%), and R5(20%). SEM: standard error of the mean.

Checklist Responses the Reviewer/Editor Comments

Manuscript ID: IJVS-21-395-Rev-I


Title: The Effect of Fermented Palm Kernel Cake Layer Quail Rations on Production Performance and Eggshell Thickness

Reviewer A

No.	Referee's Comments	Responses (for author)
1.	Line 1: remove "mixed with humic acid"	Thank you, we already changed
2.	Line 7: Change "rationed" to "supplemented"	Thank you, we already changed
3.	Line 7-8: Change "quail egg quality" to "egg quality of quails"	Thank you, we already changed
4.	Line 11: Change "replication" to "replicates in each group"	Thank you, we already changed
5.	Line 13-14: Change "can" to "may"	Thank you, we already changed
6.	Line 49: Erase "are" and added "were"	Thank you, we already changed the sentence
7.	Line 52: Change "duplicates" to "replicates"	Thank you, we already changed
8.	Line 53: Change "shows" to "showed"	Thank you, we already changed
9.	Line 62: Change "shows" to "showed"	Thank you, we already changed the sentence.
10.	Line 65: Change "include" to "included".	Thank you, we already changed
11.	Results and Discussion: separate results and discussion	Thank you, we already changed
12.	Added Author's Contribution.	Thank you, we already added section.
13.	Reference: DOI no missing	Several journal publishers not provide DOI link

4. Review version: 2nd (13 Oktober 2021)

Pemberitahuan Hasil review kedua (II) dari Reviewer.



Manuskrip dari Reviewer

Short Communication

The Effect of Fermented Palm Kernel Cake Layer Quail Rations on Production Performance and Eggshell Thickness

ABSTRACT

Some format issues to be solved; Follow the green font

DOI of many references missing

As Short Communication, keep list of references to 15-17. Delete old one.

The purpose of the study was to evaluate how palm kernel cake fermented by *Sclerotium rolfsii* and supplemented with humic acid affected production performance and egg quality of quails. The animals used in this study were 200 layer quail (*Coturnix coturnix japonica*) at 14 weeks of age. Fermented palm kernel cake (FPKC) with rationed compositions of 0, 5, 10, 15, and

20% were utilised in the research treatment, which lasted for eight weeks. Also, this study used a randomised design with four replicates in each group. The results showed that feed intake, egg production, feed conversion, egg weight, and eggshell thickness of layer quail were not statistically significant ($P>0.05$). Conclusively, palm kernel cake fermented with *S. rolfssii* may be utilised in laying quail feed at a concentration of up to 20%.

Key words: Egg quality, Humic acid, Japanese quail, Palm kernel cake, *Sclerotium rolfssii*

INTRODUCTION

No space between value and unit

Palm kernel cake (PKC) is a by-product of palm oil processing, which can serve as a potential ingredient in poultry feed. Furthermore, its nutritional composition includes crude protein, crude fibre, crude fat, calcium, phosphorus at 16.07 %, 21.30 %, 8.23 %, 0.27 %, and 0.94 %, respectively, as well as copper at 48.4 ppm. (Mirnawati et al. 2010). The crude protein content of PKC is relatively high, yet its use in poultry rations is still limited. According to Rizal (2000), PKC at a concentration of up to 10 % can be used instead of 40 % soybean meal in broiler diet due to the high β -Manan content in the coarse fibres, which may be undesirable since birds do not have fibre-breaking enzymes for manan in the digestive tract (Sundu et al. 2006). Therefore, PKC must first be processed to improve its quality with the aid of fermentation biotechnology that utilises cellulolytic and mannanolytic moulds (Meryandini et al. 2008; Mirnawati et al. 2018; Purwadaria and Haryati, 2003). Furthermore, this can reduce the content of crude fibre and manan while increasing the quality of palm kernel cake such that it can replace the soybean meal in poultry rations.

Arrange references yearwise.

Sclerotium rolfssii is a cellulolytic and mannanolytic microorganism that can be used for the fermentation of palm kernel cake. According to Razak et al. (2006), the mannanase enzyme activity of *S. rolfssii* is greater than that of *Aspergillus niger*. The fermentation of palm kernel cake with *S. rolfssii* yielded crude protein, retained nitrogen, crude fibre, and digestible crude fibre at 26.90, 54.86, 14.86, and 58.41%, respectively, as well as crude fat at 0.22% and 2557.6kcal/kg. However, the use of palm kernel cake in broiler diets is still restricted to 25% despite the rise in its nutritional content and quality.

Mirawati et al. (2017) processed palm kernel cake mixed with humic acid through the fermentation process using *S. rolfssii*. The result of this study showed an increase in crude protein, nitrogen retention, and crude fibre digestibility at 27.43 %, 59.17 %, and 55.40 %, respectively, as well as a decrease in crude fibre at 11.53 %. After fermentation, the increased nutritional content of palm kernels enables its use as a quail feed ingredient. Therefore, it is necessary to conduct research in order to determine the effect of fermented palm kernel cake containing *S. rolfssii* in rations on the production performance and quality of laying quails.

MATERIALS AND METHODS

The samples used in this study were 200 quail laying hens aged about 14-weeks old, which were confined in individual cages of size 45×20×30 cm as ten laying birds per unit. The study used a fully randomized design (CRD) with five treatments containing 0, 5, 10, 15, and 20% compositions of FPKC, as well as four replicates each. The diets used included iso-protein and iso-caloric at 20% and 2700 kcal/kg, respectively. Subsequently, Table 1 showed the diet formulation, nutritional and metabolizable energy levels of treatment diets. The diet formulation was made up of yellow corn, rice bran, meat meal, CP 126 concentrate feed (Charoen Pokphand Indonesia), top mix and FPKC. In addition, drinking water and experimental diet were provided *ad-libitum*.

The fermented palm kernel cake was made using a combination of PKC and rice bran at 80 % and 20 %, respectively, which were fermented with *S. rolfssii* and added to 200ppm humic acid. The inoculum dosage was 10% of the substrate, and the incubation period was seven days. After harvesting, the product is dried and milled before being incorporated into quail diets. Meanwhile, layer quail have a feeding period of two months or eight weeks. Table 1 showed the composition of the feeding or diet treatments.

Data Collection

The data collected during the study included feed conversion, egg mass production (g/head/day), egg weight (g/egg/head), feed intake (g/head/day), quail day egg production (%) and the eggshell thickness (mm) of laying quail, which were measured following Nuraini et al. (2012).

Data Analysis

All data were analyzed by analysis of variance based on a completely randomized design according to Steel and Torrie (1991).

RESULTS

There was no significant difference ($P>0.05$) in the feed intake of the laying quails based on the levels of FPKC in the diets since an increase in the level of FPKC did not reduce the feed consumption of the laying quails. Table 2 showed the feed intake, which ranged between 20.37-22.30 g/head/day. Similarly, there was no significant difference ($P>0.05$) in the feed consumption of the laying quails based on the levels of FPKC in the diets. The effects of the amount of FPKC in the diet on the daily egg production of quails were not significant ($P>0.05$). However, increased amounts of FPKC in quail diets can sustain the egg production of laying quails. The egg production of laying quails in this experiment ranged between 70.80 to 72.05 %, as shown in Table 2.

The quantity of FPKC in meals did not affect egg mass production of laying quails ($P>0.05$). During the trial, the egg mass output of laying quails ranged from 7.46 to 7.67g/head/day. Feed conversion is the ratio of feed intake to egg mass, and it was non-significant ($P>0.05$) when FPKC levels in diets ranged between 2.74-3.08. Different amounts of FPKC in diets had no effect on the egg weight of laying quail ($P>0.05$), which ranged from 10.61 to 11.71 g/egg/head. The amounts of FPKC in meals had no effect on the thickness of the eggshells from laying quails ($P>0.05$), which ranged from 0.27 to 0.29mm.

DISCUSSION

The difference in feed consumption of laying quail rations between treatments R1 and R5 suggests that feeding FPKC with *S. rolfesii* to 20% (R5) has the same palatability. However, these FPKC-containing meals were discovered to have a higher quality and aroma than the original substrate. According to Mirnawati et al. (2018), the fermentation process can improve the physical and chemical characteristics such as aroma, taste, and texture compared to the original substrate. Furthermore, it was also observed that feed palatability,

digestibility, and diet composition all affect the amount of feed eaten by birds (Mirnawati et al. 2019; Mirnawati et al. 2020; Ciptaan et al. 2020). In this study, parameters such as age, type, as well as energy and protein consumption were all relatively equal. Ciptaan et al. (2020) obtained a quail feed intake of about 22.14 g/head/day by adding 25 % palm oil sludge fermented with *Phanerochaeta chrysosporium* and *Neurospora crassa* in rations. Palm oil sludge is another promising by-product that can replace the ingredients of standard feed, such as yellow maize and soybean meal, in poultry diets. Previous research did not reveal a significant difference in the feed conversion ratio of broilers (Mirnawati et al. 2021).

The daily quail production was unaffected in treatments R1, R2, R3, R4, and R5 since fermented palm kernel cake provides enough nutrients needed for poultry production. Subsequently, fermentation can increase digestibility, which is in line with the hypothesis of Sukaryana et al. (2010), Dairo and Fasuyi (2008), Mirnawati et al. (2013), and Mirnawati et al. (2019) that fermented farm waste will have favourable nutritional profiles. Meanwhile, there was no significant difference in the egg production of quails since it was associated with the consumption of rations. This is consistent with the study by Akbarillah et al. (2010) and Fajrona et al. (2020) that egg production is controlled by the amount of food ingested, particularly nutrient consumption, as well as environmental variables. Consumption improves the quantity and quality of egg production since a majority of the nutrients consumed will be transformed into eggs, in addition to the fundamental needs of the birds. This value obtained in this result is greater than what was obtained in the previous study by Ciptaan et al. (2020), which reported daily egg production of 60.21 % using palm oil sludge fermented with *N. crassa* at a ration level of 12 %. According to a study by Abbas et al. (2016), the rate of feed intake Japanese quails at seven weeks of age, supplemented with 15 g/kg *Cucurbita moschata* seeds oil over a period of 1-3 weeks, was 135.5g.

According to Abou El-Ghar and Debes (2013) and Vercese et al. (2012), egg mass is related to egg weight and egg production pattern. This is comparable to the results obtained by Nuraini et al. (2012), who obtained an egg mass in the range of 6.85-7.20g/head/day by administering a mixture of sago pulp and tofu waste fermented with *N. crassa* at a 12 % ration. According to Ciptaan et al. (2020), the quail egg mass was reduced by 6.11 g/head/day when palm oil sludge fermented with *P. chrysosporium* and *N. crassa*, was incorporated to quail feed at a 25 % ration level.

The_FPKC treatment has no effect on feed consumption or egg mass, the ration conversion is also relatively the same. Table 1 shows that quails fed a ration containing up to 20% FPKC are similarly efficient in egg production to quails fed with the control ration (R1), which demonstrates that quails are equally efficient in FPKC-containing diets.

The result of this study is greater than that obtained by Nuraini et al. (2012), which utilised ration conversions ranging from 2.82 to 2.90 with a mixture of sago pulp and tofu pulp fermented with *N. crassa* 12% in rations. This figure is lower when compared to the feed conversion value from quail ration with palm kernel cake fermented with *Bacillus subtilis* at a 25% ration level. (Fajrona et al. 2020). However, The results were more desirable than what was obtained from a mixture of 200-600 mg/kg L-Carnitine and Japanese quail diet with a feed conversion ratio ranging between 5.8 to 7.7 (Mahmoud et al. 2020).

A minor variation in quail egg weight produced by the fermentation process might break down complex or low digestible components into simpler molecular structures, improving nutritional absorption and the quality of poultry products. Conversely, beneficial primary and secondary metabolites are secreted by microbes throughout the incubation process. Furthermore, Mirnawati et al. (2019) showed that fermented palm kernel meal has higher amino acid quality after fermentation. The egg weight obtained from this study was higher than previous results obtained by Fajrona et al. (2020) and Nuraini et al. (2012), which were **9.57 - 9.64 g/egg/head and 10.29 g/egg/head, respectively.**

The eggshell thickness treatment of R1 to R5 showed that FPKC up to 20% in the diet still provides almost similar results because the inclusion of FPKC and humic acid as mineral sources increases the bioavailability of calcium and phosphorus, both of which play significant roles in the eggshell formation process. According to Enviromate (2002), humic acid is a source of minerals and organic compounds. Also, Korsakov et al. (2019) found that **about 50-75 ml** of humic acid given through drinking water significantly affects the eggshell thickness, which was **0.35 - 0.36 mm** in laying hens. Ciptaan et al. (2020) measured the average thickness of quail eggshells to be **0.26-0.28 mm**. Additionally, Zita et al. (2013), also reported that the average thickness of quail eggs was **0.19 mm**.

Conclusion

Conclusively, palm kernel cake fermented with *Sclerotium rolfsii* can be utilized up to 20% in quail diets. The results showed that the feed intake, egg production, egg mass production, feed conversion, egg weight, and eggshell thickness were found to be **22.30 g/head/day, 70.93%, 7.54 g/head/day, 3.08, 10.65 g/egg/head, and 0.27 mm, respectively**. Therefore, it is expected that palm kernel cake would be able to partially replace the current ingredients used in commercial feed in order to enhance the profitability of quail layer farming.

Acknowledgements

This study was funded by the BOPTN of Andalas University, number 42/UN.16.17/PP.RGB/LPPM/2019, on the 23rd of April, 2019. I am grateful to everyone with whom I had the pleasure of working on this project.

Author's Contribution

Gita Ciptaan supervised the experiment and writing original manuscript. Mirnawati and Ferawati conducted the experiment and analyzed the data. Malik Makmur finalize manuscript.

REFERENCES

- Abbas RJ, Alshaheen SA and Majeed TI, 2017. Evaluation of the productive and physiological performance of Japanese quail (*Coturnix coturnix japonica*) fed different levels of pumpkin (*Cucurbita moschata*) seeds oil. International Journal of Veterinary Science 6: 31-35.
- Abou El-Ghar RS and Debes AA, 2013. Prediction of breeding values and selection for egg mass in a closed flock of white leghorn. International Journal of Livestock Production Research 1:16-25.
- Canellas LP and Olivares FL, 2014. Physiological responses to humic substances as plant growth promoter. Chemical and Biological Technologies in Agriculture 1: 3.
<https://doi.org/10.1186/2196-5641-1-3>

- Ciptaan G, Mirnawati and Djulardi A, 2020. Performance quality feed palm oil sludge fermented with *Phanerochaete cryosporium* and *Neurospora crassa*. *Livestock Research for Rural Development* 32: 8. (DOI no missing ?)
- Dairo FAS and Fauyi AO, 2008. Evaluation of fermented palm kernel meal and fermented copra meal protein as substitute for soybean meal protein in laying hens diets. *Journal Central European Agriculture* 9: 35-44. (DOI no missing ?)
- Enviromate, 2002. Effect of humic acid on animal and humans (literature review and current research), Effects of humic acid, Enviromate Inc. 8571. Boat Club Road, Forth Worth, Texas 76719.
- Fajrona K, Aini Q and Mirnawati, 2020. The effect of fermented palm kernel cake with *Bacillus subtilis* ratios on production performance and quail egg quality. *Quest Journal of Research in Agriculture and Animal Science* 7: 6-10. (DOI no missing ?)
- Korsakov KV, Vasiliev AA, Sivokhina LA, Zabelina MV, Murtazaeva RN, Daeva TV and Kokorev VA, 2019. The influence of humic acid supplement on the marketable properties of hy-line laying hen eggs. *Advances in Animal and Veterinary Sciences* 7: 66-70.
<http://dx.doi.org/10.17582/journal.aavs/2019/7.s1.66.70>
- Mahmoud RE, Elshopakey GE and Awadin WF, 2020. Effects of feeding diets supplemented with different levels of l-carnitine on growth performance, serum metabolites, histopathological changes in growing Japanese quails. *International Journal of Veterinary Science* 9: 16-23. (DOI no missing ?)
- Meryandini A, Angreandari R and Rahmania N, 2008. Isolasi bakteri mananolitik dan karakterisasi mananasenya. *Biota* 13: 82-88.
<https://doi.org/10.24002/biota.v13i2.2675>
- Mirnawati, Ciptaan G and Ferawati, 2019. Improving the quality and nutrient content of palm kernel cake through fermentation with *Bacillus subtilis*. *Livestock Research for Rural Development* 31: 119-123.
- Mirnawati, Ciptaan G and Ferawati, 2020. Broiler performance on a diet containing palm kernel meal fermented with *Bacillus subtilis*. *Livestock Research for Rural Development* 32: 2020. (DOI no missing ?)

- Mirawati, Ciptaan G, Djulardi A and Makmur M, 2021. Broiler response to the utilization of fermented palm oil sludge with *Phanerochaeta chrysosporium* and *Neurospora crassa*. International Journal of Veterinary Science. In press article.
<https://doi.org/10.47278/journal.ijvs/2021.089>
- Mirawati, Djulardi A and Ciptaan G, 2018. Utilization of fermented palm kernel cake with *Sclerotium rolfsii* in broiler ration. International Journal of Poultry Science 17 : 342-347. <https://dx.doi.org/10.3923/ijps.2018.342.347>
- Mirawati, Rizal Y and Marlida Y, 2013. Effect of humic acid addition via drinking water on the performance of broiler fed diet containing fermented and non fermented palm kernel cake. Archiva Zootechnica 16: 41-53. (DOI no missing ?)
- Nuraini, Sabrina and Latif SA, 2012. Performances and quail egg quality feeding product fermented with *Neurospora crassa*. Indonesian Journal of Animal Science 14: 385-391. <https://doi.org/10.25077/jpi.14.2.385-391.2012>
- Purwadaria T and Haryati T, 2003. *In vitro* digestibility evaluation of coconut meal incorporated precipitate Beta D Manannase from *Eupenicillium javanicum*. Journal Microbiology Indonesia, 2003: 19 -21. (DOI no missing ?)
- Razak NA, 2006. Production and purification of mannanase degrading enzymes from palm kernel cake fermented by *Aspergillus niger* and *Sclerotium rolfsii*. Master thesis Universiti Putra Malaysia, Malaysia.
- Rizal Y, 2000. The response of broiler to the substitution part of soybean meal for palm kernel cake in the diet. Jurnal Peternakan Lingkungan 2: 15-20. (DOI no missing ?)
- Steel RGD and Torrie JH, 1991. Prinsip and Prosedur Statistik: Suatu pendekatan Biometrik PT. Gramedia Pustaka Utama. Jakarta, Indonesia.
- Sukaryana Y, Atmomarsono U, Yunianto VD and Supriatna E, 2010. Bioconversions of palm kernel cake and rice bran mixture by *Trichoderma viridae* toward nutritional contents. International Journal of Science and Engineering 1: 27 - 32.
<http://dx.doi.org/10.12777/ijse.1.2.27-32>

Sundu B, Kumar A and Dingle J, 2006 Palm kernel meal in broiler diets:effect on chicken performance and health. World's Poultry Science Journal 62 316-325.

<https://doi.org/10.1079/WPS2005100>

Vercese F, Garcia EA, Sartori JR, Silva Ade P, Faltarone ABG, Berto DA, Molino A de B and Pelícia K, 2012. Performance and egg quality of Japanese quails submitted to cyclic heat stress. Brazilian Journal Poultry Science 14: 37-41.

<https://doi.org/10.1590/S1516-635X2012000100007>

Zita L, Ledvinka Z and Klesalova L, 2013. The effect of the age of Japanese quails on certain egg quality traits and their relationship. Veterinary Archives. 83: 223-232. (DOI no missing ?)

Table 1: Diet formulation (%), nutrient content (%) and metabolizable energy (kcal/kg).

Item	Treatment Ration				
	R1	R2	R3	R4	R5
Corn	45.5	45	44	43	43
Rice brand	20	18	16	13.5	11.5
Meat meal	6	6	6	6	6
CP 126 concentrate	26.5	24	22	20.5	17.5
FPKC	0	5	10	15	20
Mineral B12	1	1	1	1	1
Top mix	1	1	1	1	1
Total (%)	100	100	100	100	100
Crude protein	20.06	20.04	20.16	20.44	20.42
Crude fat	3.46	3.40	3.34	3.28	3.29
Crude fiber	6.13	6.17	6.23	6.27	6.50
Calcium	2.94	2.80	2.63	2.60	1.94

Phosphor	0.88	0.89	0.91	0.93	0.82
Metabolizable energy	2710	2708	2703	2706	2725


Table 2: The effect of dietary fermented palm kernel cake (FPKC) on laying quail feed intake, egg production, egg mass, feed conversion, egg weight, and eggshell thickness.

Variables	Treatment (%)					SEM
	R1	R2	R3	R4	R5	
Feed intake (g/head/day)	20.37	21.61	21.71	21.47	22.30	0.43
Egg production (%)	71.80	71.61	70.80	72.05	70.93	2.01
Egg mass (g/head/day)	7.67	7.58	7.46	7.57	7.54	0.20
Feed conversion ratio	2.74	2.98	3.01	2.99	3.08	0.08
Egg weight (g)	10.69	10.61	11.71	10.48	10.65	0.50
Eggshell thickness (mm)	0.29	0.29	0.28	0.29	0.27	2.49

Note: Inclusion FPKC in R1 to R5 was R1 (0%-Control), R2 (5%), R3 (10%), R4 (15%), and R5(20%). SEM: standard error of the mean.

5. Revised version: 2nd (14 Oktober 2021)

Tanggal 14 Oktober 2021: Penulis mengirim hasil perbaikan II (a revise author version)



MANUSKRIP PERBAIKAN REVISI II

Short Communication

The Effect of Fermented Palm Kernel Cake Layer Quail Rations on Production Performance and Eggshell Thickness

ABSTRACT

The purpose of the study was to evaluate how palm kernel cake fermented by *Sclerotium rolfsii* and supplemented with humic acid affected production performance and egg quality of quails. The animals used in this study were 200 layer quail (*Coturnix coturnix japonica*) at 14 weeks of age. Fermented palm kernel cake (FPKC) with rationed compositions of 0, 5, 10, 15, and 20% were utilised in the research treatment, which lasted for eight weeks. Also, this study used a randomised design with four replicates in each group. The results showed that feed intake, egg production, feed conversion, egg weight, and eggshell thickness of layer quail were not

statistically significant ($P>0.05$). Conclusively, palm kernel cake fermented with *S. rolfsii* may be utilised in laying quail feed at a concentration of up to 20%.

Key words: Egg quality, Humic acid, Japanese quail, Palm kernel cake, *Sclerotium rolfsii*

INTRODUCTION

Palm kernel cake (PKC) is a by-product of palm oil processing, which can serve as a potential ingredient in poultry feed. Furthermore, its nutritional composition includes crude protein, crude fibre, crude fat, calcium, phosphorus at 16.07%, 21.30%, 8.23%, 0.27%, and 0.94%, respectively, as well as copper at 48.4 ppm. (Mirnawati et al. 2010). The crude protein content of PKC is relatively high, yet its use in poultry rations is still limited. PKC at a concentration of up to 10% can be used instead of 40% soybean meal in broiler diet due to the high β -Manan content in the coarse fibres, which may be undesirable since birds do not have fibre-breaking enzymes for manan in the digestive tract (Sundu et al. 2006). Therefore, PKC must first be processed to improve its quality with the aid of fermentation biotechnology that utilises cellulolytic and mannanolytic moulds (Meryandini et al. 2008; Mirnawati et al. 2018). Furthermore, this can reduce the content of crude fibre and manan while increasing the quality of palm kernel cake such that it can replace the soybean meal in poultry rations.

Sclerotium rolfsii is a cellulolytic and mannanolytic microorganism that can be used for the fermentation of palm kernel cake. According to Razak (2006), the mannanase enzyme activity of *S. rolfsii* is greater than that of *Aspergillus niger*. The fermentation of palm kernel cake with *S. rolfsii* yielded crude protein, retained nitrogen, crude fibre, and digestible crude fibre at 26.90, 54.86, 14.86, and 58.41%, respectively, as well as crude fat at 0.22% and 2557.6kcal/kg. However, the use of palm kernel cake in broiler diets is still restricted to 25% despite the rise in its nutritional content and quality.

Mirnawati et al. (2017) processed palm kernel cake mixed with humic acid through the fermentation process using *S. rolfsii*. The result of this study showed an increase in crude protein, nitrogen retention, and crude fibre digestibility at 27.43%, 59.17%, and 55.40%, respectively, as well as a decrease in crude fibre at 11.53%. After fermentation, the

increased nutritional content of palm kernels enables its use as a quail feed ingredient. Therefore, it is necessary to conduct research in order to determine the effect of fermented palm kernel cake containing *S. rolfesii* in rations on the production performance and quality of laying quails.

MATERIALS AND METHODS

The samples used in this study were 200 quail laying hens aged about 14-weeks old, which were confined in individual cages of size 45×20×30 cm as ten laying birds per unit. The study used a fully randomized design (CRD) with five treatments containing 0, 5, 10, 15, and 20% compositions of FPKC, as well as four replicates each. The diets used included iso-protein and iso-caloric at 20% and 2700kcal/kg, respectively. Subsequently, Table 1 showed the diet formulation, nutritional and metabolizable energy levels of treatment diets. The diet formulation was made up of yellow corn, rice bran, meat meal, CP 126 concentrate feed (Charoen Pokphand Indonesia), top mix and FPKC. In addition, drinking water and experimental diet were provided *ad-libitum*.

The fermented palm kernel cake was made using a combination of PKC and rice bran at 80% and 20%, respectively, which were fermented with *S. rolfesii* and added to 200ppm humic acid. The inoculum dosage was 10% of the substrate, and the incubation period was seven days. After harvesting, the product is dried and milled before being incorporated into quail diets. Meanwhile, layer quail have a feeding period of two months or eight weeks. Table 1 showed the composition of the feeding or diet treatments.

Data Collection

The data collected during the study included feed conversion, egg mass production (g/head/day), egg weight (g/egg/head), feed intake (g/head/day), quail day egg production (%) and the eggshell thickness (mm) of laying quail, which were measured following Nuraini et al. (2012).

Data Analysis

All data were analyzed by analysis of variance based on a completely randomized design according to Steel and Torrie (1991).

RESULTS

There was no significant difference ($P>0.05$) in the feed intake of the laying quails based on the levels of FPKC in the diets since an increase in the level of FPKC did not reduce the feed consumption of the laying quails. Table 2 showed the feed intake, which ranged between 20.37-22.30g/head/day. Similarly, there was no significant difference ($P>0.05$) in the feed consumption of the laying quails based on the levels of FPKC in the diets. The effects of the amount of FPKC in the diet on the daily egg production of quails were not significant ($P>0.05$). However, increased amounts of FPKC in quail diets can sustain the egg production of laying quails. The egg production of laying quails in this experiment ranged between 70.80 to 72.05%, as shown in Table 2.

The quantity of FPKC in meals did not affect egg mass production of laying quails ($P>0.05$). During the trial, the egg mass output of laying quails ranged from 7.46 to 7.67g/head/day. Feed conversion is the ratio of feed intake to egg mass, and it was non-significant ($P>0.05$) when FPKC levels in diets ranged between 2.74-3.08. Different amounts of FPKC in diets had no effect on the egg weight of laying quail ($P>0.05$), which ranged from 10.61 to 11.71g/egg/head. The amounts of FPKC in meals had no effect on the thickness of the eggshells from laying quails ($P>0.05$), which ranged from 0.27 to 0.29mm.

DISCUSSION

The difference in feed consumption of laying quail rations between treatments R1 and R5 suggests that feeding FPKC with *S. rolf sii* to 20% (R5) has the same palatability. However, these FPKC-containing meals were discovered to have a higher quality and aroma than the original substrate. According to Mirnawati et al. (2018), the fermentation process can improve the physical and chemical characteristics such as aroma, taste, and texture compared to the original substrate. Furthermore, it was also observed that feed palatability, digestibility, and diet composition all affect the amount of feed eaten by birds (Mirnawati et al. 2019; Mirnawati et al. 2020; Ciptaan et al. 2020). In this study, parameters such as age, type, as well as energy and protein consumption were all relatively equal. Ciptaan et al. (2020) obtained a quail feed intake of about 22.14g/head/day by adding 25% palm oil sludge

fermented with *Phanerochaeta chrysosporium* and *Neurospora crassa* in rations. Palm oil sludge is another promising by-product that can replace the ingredients of standard feed, such as yellow maize and soybean meal, in poultry diets. Previous research did not reveal a significant difference in the feed conversion ratio of broilers (Mirnawati et al. 2021).

The daily quail production was unaffected in treatments R1, R2, R3, R4, and R5 since fermented palm kernel cake provides enough nutrients needed for poultry production. Subsequently, fermentation can increase digestibility, which is in line with the hypothesis of Sukaryana et al. (2010) and Mirnawati et al. (2019) that fermented farm waste will have favourable nutritional profiles. Meanwhile, there was no significant difference in the egg production of quails since it was associated with the consumption of rations. This is consistent with the study by Akbarillah et al. (2010) that egg production is controlled by the amount of food ingested, particularly nutrient consumption, as well as environmental variables. Consumption improves the quantity and quality of egg production since a majority of the nutrients consumed will be transformed into eggs, in addition to the fundamental needs of the birds. This value obtained in this result is greater than what was obtained in the previous study by Ciptaan et al. (2020), which reported daily egg production of 60.21% using palm oil sludge fermented with *N. crassa* at a ration level of 12%. According to a study by Abbas et al. (2017), the rate of feed intake Japanese quails at seven weeks of age, supplemented with 15g/kg *Cucurbita moschata* seeds oil over a period of 1-3 weeks, was 135.5g.

According to Abou El-Ghar and Debes (2013) and Vercese et al. (2012), egg mass is related to egg weight and egg production pattern. This is comparable to the results obtained by Nuraini et al. (2012), who obtained an egg mass in the range of 6.85-7.20g/head/day by administering a mixture of sago pulp and tofu waste fermented with *N. crassa* at a 12% ration. According to Ciptaan et al. (2020), the quail egg mass was reduced by 6.11g/head/day when palm oil sludge fermented with *P. chrysosporium* and *N. crassa*, was incorporated to quail feed at a 25% ration level.

The FPKC treatment has no effect on feed consumption or egg mass, the ration conversion is also relatively the same. Table 1 shows that quails fed a ration containing up to 20% FPKC are similarly efficient in egg production to quails fed with the control ration (R1), which demonstrates that quails are equally efficient in FPKC-containing diets.

The result of this study is greater than that obtained by Nuraini et al. (2012), which utilised ration conversions ranging from 2.82 to 2.90 with a mixture of sago pulp and tofu pulp fermented with *N. crassa* 12% in rations. However, The results were more desirable than what was obtained from a mixture of 200-600mg/kg L-Carnitine and Japanese quail diet with a feed conversion ratio ranging between 5.8 to 7.7 (Mahmoud et al. 2020).

A minor variation in quail egg weight produced by the fermentation process might break down complex or low digestible components into simpler molecular structures, improving nutritional absorption and the quality of poultry products. Conversely, beneficial primary and secondary metabolites are secreted by microbes throughout the incubation process. Furthermore, Mirnawati et al. (2019) showed that fermented palm kernel meal has higher amino acid quality after fermentation. The egg weight obtained from this study was higher than previous results obtained by Nuraini et al. (2012) which were 9.57 - 9.64g/egg/head.

The eggshell thickness treatment of R1 to R5 showed that FPKC up to 20% in the diet still provides almost similar results because the inclusion of FPKC and humic acid as mineral sources increases the bioavailability of calcium and phosphorus, both of which play significant roles in the eggshell formation process. Korsakov et al. (2019) found that about 50-75ml of humic acid given through drinking water significantly affects the eggshell thickness, which was 0.35 - 0.36mm in laying hens. Ciptaan et al. (2020) measured the average thickness of quail eggshells to be 0.26-0.28mm.

Conclusion

Conclusively, palm kernel cake fermented with *Sclerotium rolfsii* can be utilized up to 20% in quail diets. The results showed that the feed intake, egg production, egg mass production, feed conversion, egg weight, and eggshell thickness were found to be 22.30g/head/day, 70.93%, 7.54g/head/day, 3.08, 10.65g/egg/head, and 0.27mm, respectively. Therefore, it is expected that palm kernel cake would be able to partially replace the current ingredients used in commercial feed in order to enhance the profitability of quail layer farming.

Acknowledgements

This study was funded by the BOPTN of Andalas University, number 42/UN.16.17/PP.RGB/LPPM/2019, on the 23rd of April, 2019. I am grateful to everyone with whom I had the pleasure of working on this project.

Author's Contribution

Gita Ciptaan supervised the experiment and writing original manuscript. Mirnawati and Ferawati conducted the experiment and analyzed the data. Malik Makmur finalize manuscript.

REFERENCES

- Abbas RJ, Alshaheen SA and Majeed TI, 2017. Evaluation of the productive and physiological performance of Japanese quail (*Coturnix coturnix japonica*) fed different levels of pumpkin (*Cucurbita moschata*) seeds oil. *International Journal of Veterinary Science* 6: 31-35.
- Abou El-Ghar RS and Debes AA, 2013. Prediction of breeding values and selection for egg mass in a closed flock of white leghorn. *International Journal of Livestock Production Research* 1:16-25.
- Ciptaan G, Mirnawati and Djulardi A, 2020. Performance quality feed palm oil sludge fermented with *Phanerochaete cryosporium* and *Neurospora crassa*. *Livestock Reasearch for Rural Development* 32: 8.
- Korsakov KV, Vasiliev AA, Sivokhina LA, Zabelina MV, Murtazaeva RN, Daeva TV and Kokorev VA, 2019. The influence of humic acid supplement on the marketable properties of hy-line laying hen eggs. *Advances in Animal and Veterinary Sciences* 7: 66-70.
<http://dx.doi.org/10.17582/journal.aavs/2019/7.s1.66.70>
- Mahmoud RE, Elshopakey GE and Awadin WF, 2020. Effects of feeding diets supplemented with different levels of l-carnitine on growth performance, serum metabolites, histopathological changes in growing Japanese quails. *International Journal of Veterinary Science* 9: 16-23.

- Meryandini A, Angreandari R and Rahmania N, 2008. Isolasi bakteri mananolitik dan karakterisasi mananasenya. *Biota* 13: 82-88.
<https://doi.org/10.24002/biota.v13i2.2675>
- Mirawati, Ciptaan G and Ferawati, 2020. Broiler performance on a diet containing palm kernel meal fermented with *Bacillus subtilis*. *Livestock Research for Rural Development* 32: 2020.
- Mirawati, Ciptaan G, Djulardi A and Makmur M, 2021. Broiler response to the utilization of fermented palm oil sludge with *Phanerochaeta chrysosporium* and *Neurospora crassa*. *International Journal of Veterinary Science*. In press article.
<https://doi.org/10.47278/journal.ijvs/2021.089>
- Mirawati, Djulardi A and Ciptaan G, 2018. Utilization of fermented palm kernel cake with *Sclerotium rolfsii* in broiler ration. *International Journal of Poultry Science* 17 : 342-347. <https://dx.doi.org/10.3923/ijps.2018.342.347>
- Mirawati, Rizal Y, Marlida Y and Kompang IP, 2010. The role of humic acid in palm kernel cake fermented by *Aspergillus niger* for poultry ration. *Pakistan Journal of Nutrition* 9: 182-185. <https://dx.doi.org/10.3923/pjn.2010.182.185>
- Nuraini, Sabrina and Latif SA, 2012. Performances and quail egg quality feeding product fermented with *Neurospora crassa*. *Indonesian Journal of Animal Science* 14: 385-391. <https://doi.org/10.25077/jpi.14.2.385-391.2012>
- Razak NA, 2006. Production and purification of mannanase degrading enzymes from palm kernel cake fermented by *Aspergillus niger* and *Sclerotium rolfsii*. Master thesis Universiti Putra Malaysia, Malaysia.
- Steel RGD and Torrie JH, 1991. Prinsip and Prosedur Statistik: Suatu pendekatan Biometrik PT. Gramedia Pustaka Utama. Jakarta, Indonesia.
- Sukaryana Y, Atmomarsono U, Yuniarto VD and Supriatna E, 2010. Bioconversions of palm kernel cake and rice bran mixture by *Trichoderma viridae* toward nutritional contents. *International Journal of Science and Engineering* 1: 27 - 32.
<http://dx.doi.org/10.12777/ijse.1.2.27-32>

Sundu B, Kumar A and Dingle J, 2006 Palm kernel meal in broiler diets:effect on chicken performance and health. *World's Poultry Science Journal* 62 316-325.

<https://doi.org/10.1079/WPS2005100>

Vercese F, Garcia EA, Sartori JR, Silva Ade P, Faitarone ABG, Berto DA, Molino A de B and Pelícia K, 2012. Performance and egg quality of Japanese quails submitted to cyclic heat stress. *Brazilian Journal Poultry Science* 14: 37-41.

<https://doi.org/10.1590/S1516-635X2012000100007>

Table 1: Diet formulation (%), nutrient content (%) and metabolizable energy (kcal/kg).

Item	Treatment Ration				
	R1	R2	R3	R4	R5
Corn	45.5	45	44	43	43
Rice brand	20	18	16	13.5	11.5
Meat meal	6	6	6	6	6
CP 126 concentrate	26.5	24	22	20.5	17.5
FPKC	0	5	10	15	20
Mineral B12	1	1	1	1	1
Top mix	1	1	1	1	1
Total (%)	100	100	100	100	100
Crude protein	20.06	20.04	20.16	20.44	20.42
Crude fat	3.46	3.40	3.34	3.28	3.29
Crude fiber	6.13	6.17	6.23	6.27	6.50
Calcium	2.94	2.80	2.63	2.60	1.94
Phosphor	0.88	0.89	0.91	0.93	0.82
Metabolizable energy	2710	2708	2703	2706	2725

Table 2: The effect of dietary fermented palm kernel cake (FPKC) on laying quail feed intake, egg production, egg mass, feed conversion, egg weight, and eggshell thickness.

Variables	Treatment (%)					SEM
	R1	R2	R3	R4	R5	
Feed intake (g/head/day)	20.37	21.61	21.71	21.47	22.30	0.43
Egg production (%)	71.80	71.61	70.80	72.05	70.93	2.01
Egg mass (g/head/day)	7.67	7.58	7.46	7.57	7.54	0.20
Feed conversion ratio	2.74	2.98	3.01	2.99	3.08	0.08
Egg weight (g)	10.69	10.61	11.71	10.48	10.65	0.50
Eggshell thickness (mm)	0.29	0.29	0.28	0.29	0.27	2.49

Note: Inclusion FPKC in R1 to R5 was R1 (0%-Control), R2 (5%), R3 (10%), R4 (15%), and R5(20%). SEM: standard error of the mean.

Checklist Responses the Reviewer/Editor Comments

Manuscript ID: IJVS-21-395-Rev-II

Title: The Effect of Fermented Palm Kernel Cake Layer Quail Rations on Production Performance and Eggshell Thickness

Reviewer A

No.	Referee's Comments	Responses (for author)
1.	As short communication, keep list references to 15-17. Delete old one	Thank you, we already reduced
2.	No space value and unit	Thank you, we already changed in entire manuscript
3.	DOI of many references missing	Thank you, we already added but some journal publisher doesn't provide DOI, such as Livestock Research for Rural Development: https://www.lrrd.org/

6. Accepted (21 Oktober 2021)

Pemberitahuan Artikel di *acceptance*.

The screenshot shows a Gmail interface on a Windows desktop. The email is from Editor IJVS to Malik Makmur. The subject is 'Short Communication-Manuscripts-IJVETS'. The email content states that the revised manuscript (IJVS-21-395) has been found acceptable after processing on FAST TRACK. It details Article Publishing Charges (APCs) for Fast Track processed paper: US\$ 250 + Over and above 06 printed pages: US\$ 25/Printed page. The paper is composed of 4 pages. The author is requested to pay US\$250 as APCs through the Western Union using recipient's first name: RAO, Middle Name: ZAHID; last name: ABBAS, city: FAISALABAD, country: PAKISTAN as early as possible. The author is also asked to send a receipt issued by the Western Union office bearing MTCN (Money Transfer Control Number) and first and last name of the sender along with the country name from where money was sent.

Dear Author,

Your revised manuscript (IJVS-21-395) has been found acceptable after processing on **FAST TRACK**. Article Publishing Charges (APCs) for Fast Track processed paper are APCs up to 6 Printed Pages: US\$ 250 + Over and above 06 printed pages: US\$ 25/Printed page. Your paper is composed of 4 pages; therefore, you are requested to pay US\$250 as APCs through the Western Union using recipient's first name: RAO, Middle Name: ZAHID; last name: ABBAS, city: FAISALABAD, country: PAKISTAN as early as possible so that your paper be sent for publishing and other processes. You will be sending us the receipt issued by the Western Union office bearing MTCN (Money Transfer Control Number) and first and last name of the sender along with the country name from where money was sent.

[Quoted text hidden]

23°C Berawan 23:43 11/04/2023

7. proses *post-review* manuskrip (20 November 2021)

The screenshot shows a Gmail interface on a Windows desktop. The email is from Editor IJVS to Malik Makmur. The subject is 'Short Communication-Manuscripts-IJVETS'. The email content thanks the author for choosing the International Journal of Veterinary Science (IJVS) for their valuable publication. It states that the paper has been composed and asks the author to perform proofreading within 1-2 days and indicate mistakes in a Table form.

Dear Author,

Thank you very much for choosing the International Journal of Veterinary Science (<https://www.ijvets.com/>) for your valuable publication. Now your paper has been composed, please perform proofreading within 1-2 days and indicate mistakes in a Table form:

Page #	Line #	Column 1 or 2	In Proofs	What change required

If you don't find any mistakes, even then inform us.

Red font references in the text or list are missing if any. You have to provide the missing references, and if present in the list, then cite these references in the text or delete.

Thank you very much again for your valuable contribution. We look forward to receiving your new research/review article soon.

[Quoted text hidden]

IJVS 21-395-4-SC-Indonesia.pdf
218K

Status baterai tersisa 64%
5h 34min

23°C Berawan 23:49 11/04/2023

Lampiran Perbaikan (Revised) Post Review/ Proofread

No.	Page	Line	Column	In Proofs	What change required
1.	1	9-10	1	Doctoral Student, Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, Andalas University, 10 Jl. Raya Unand, Kampus Limau Manis, Padang 25163, West Sumatera, Indonesia	Post-doctoral researcher, IPB University, Bogor, West Java, Indonesia
2.	1	46	2	Mirawati et al. (2017)	Mirawati, Ciptaan G and Ferawati, 2017. The effect of mannanolytic fungi and humic acid dosage to improve the nutrient content and quality of fermented palm kernel cake. <i>International Journal ChemTech Research</i> 10: 56-61.
3.	2, 3	155, 172, 222	2, 1	Mirawati et al. (2019)	Mirawati, Ciptaan G and Ferawati, 2019. Improving the quality and nutrient content of palm kernel cake through fermentation with <i>Bacillus subtilis</i> . <i>Livestock Research for Rural Development</i> 31: 2019.
4.	3	177	1	Akbarillah et al. (2010)	Akbarillah T, Kusuyah and Hidayat, 2010. The effect of fresh indigofera leaves utilization as feed supplementation on egg production and its yolk color of ducks. <i>Jurnal Sain Peternakan Indonesia</i> 5: 27-33. https://doi.org/10.31186/jspi.id.5.1.27-33
5.	3	183	1	needs of the birds These values	needs of the birds. These values
6.	3	241	2	<i>Sclerotium rolfsii</i>	<i>Sclerotium rolfsii</i>

8. Published

Artikel Publish, tersedia online di International Journal of Veterinary Science:

<http://www.ijvets.com/pdf-files/Volume-11-no-3-2022/400-403.pdf>

The screenshot displays the website interface for the International Journal of Veterinary Science. The main content area lists several research articles with their titles, authors, and publication details. On the right side, there is a SCOPUS CiteScore Trend chart showing an upward trend from 2017 to 2022. Below the chart, there are sections for 'Key Features' including 'doi' and 'Crossref' logos, 'OPEN ACCESS' badge, and a Creative Commons Attribution-NonCommercial (CC BY-NC) license logo.

Characterization of Lactobacillus Species Recovered from Raw Dromedary Milk in Relation to its Antimicrobial Activity
Hani M. Khyria, Sherif Marouf, Mai H Hanafy, Alif Hussein, Aalia SA Saad and Heidy Abo-El Yazeed
Int J Vet Sci. 2022, 11(3): 373-377.
Abstract Full text pdf

Prevalence and Types of Fungi in Kintamani Bali Dogs
Pulu Henrywaesa Sudipa and Ketul Tono Pasek Geigel
Int J Vet Sci. 2022, 11(3): 378-383.
Abstract Full text pdf

Nanofomulations of Pentacyclic Triterpenoids: Chemoprevention and Anticancer
Khalifa El-Dawy, Dosa Mohamed and Zenab Abbou
Int J Vet Sci. 2022, 11(3): 384-391.
Abstract Full text pdf

Effect of Acetaminophen on the Estrous Cycle and Reproductive Hormones of Female Mice
Nofei AN, Mutembei HM, Kaingu CK, Muthee JK, Birech Z and Ondeki AM
Int J Vet Sci. 2022, 11(3): 392-395.
Abstract Full text pdf

Prevalence of Lung Lesions in Imported Cattle Slaughtered at Abu Simble Abattoir, Egypt
Ameer Mohamed Abdel-Az Abdel-Rassid, Ali Mewarad Ahmed, Hassan Mohamed Sobhy, Sherin S. Abdelgayed and Sahar Hussein Abdalla Hakiel
Int J Vet Sci. 2022, 11(3): 396-399.
Abstract Full text pdf

The Effect of Fermented Palm Kernel Cake Layer Quail Rations on Production Performance and Eggshell Thickness
Gita Ciptaan, Mirrawati, Farwati and Malik Malmur
Int J Vet Sci. 2022, 11(3): 400-403.
Abstract Full text pdf

The response of Vitamin D to the Motility of Isolated Intestines in Male Rats
Nabil A. Soliman, Sherif Waigh Mansour, Safaa E. Nassar, Aya Shawky Mabrouly and Reem Adly Ahmad
Int J Vet Sci. 2022, 11(3): 404-408.
Abstract Full text pdf

SCOPUS CiteScore Trend
2017 2018 2019 2020 2021 2022
CiteScore Value
Research in category

Key Features
doi Crossref
ISSN: 1842-2129/ijvets@ijv
OPEN ACCESS
CC BY-NC