

# Jurnal Ajava\_2016\_Zain et al

*by* Mardiati Zain

---

**Submission date:** 06-Jun-2023 11:40AM (UTC+0800)

**Submission ID:** 2109998824


**File name:** Artikel\_Supplementation\_of\_Saccharomyces\_Cerevisiae\_2.pdf (674.87K)

**Word count:** 3705

**Character count:** 19813

# aJava

Asian Journal of Animal and Veterinary Advances



Academic  
Journals Inc.

[www.academicjournals.com](http://www.academicjournals.com)



## Research Article

# Supplementation of *Saccharomyces cerevisiae* and *Sapindus rarak* in Diet Based of Oil Palm Frond (OPF) on Nutrient Digestibility and Daily Weight Gain of Goat

Mardiati Zain, Jurnida Rahman, Khasrad and Erpomen

Department of Animal Nutrition, Faculty of Animal Science, Andalas University, Kampus Limau Manis, 25163 Padang, Indonesia

## Abstract

This experiment was to study the effect of adding *Saccharomyces cerevisiae* and *Sapindus rarak* in diet Oil Palm Frond (OPF) based of nutrients digestibility and body weight gain of goats. Experimental design used randomized blok design, with 4 treatments and 4 groups as replications. This experiment used sixteen ettawa goats ( $40 \pm 8.5$  kg liveweight). The four treatments were (A) native grass+concentrate (B), OPF (previously treated with 6% urea)+concentrate, (C) Diet B+1% *Saccharomyces cerevisiae* and (D) Diet C+4% *Sapindus rarak*. The animals were adjusted to their treatments over a 2 week preliminary period which was followed by an 8 week experimental period. Feed intake was determined daily and live weight was measured every two week. Digestibility was measured by total faecal collection over 6 consecutive days on last week of experiment period. The results showed that the dry matter digestibility of treatment B (50.74%) were significantly lower than treatments A, C and D respectively (67.37, 62.38 and 65.71%) and supplementation of *Saccharomyces cerevisiae* was able to improve nutrient digestibility and body weight gain of goat, but still low compared to control (A). Addition of 4% *Sapindus rarak* in treatment D ( $123 \text{ g day}^{-1}$ ) was able to provide digestibility and body weight gain similar to the control ration ( $132 \text{ g day}^{-1}$ ). It can be concluded that the use of OPD as a substitute for grass in goat diets would give the same results with the grass when added 1% *Saccharomyces cerevisiae* and 4% *Sapindus rarak*.

**Key words:** Oil palm frond, *Sapindus rarak*, *Saccharomyces cerevisiae*, goat, digestibility, weight gain, ammoniated, defaunation

**Received:** March 11, 2016

**Accepted:** March 29, 2016

**Published:** April 15, 2016

**Editor:** Dr. Kuldeep Dhama, Principal Scientist, Division of Pathology, Indian Veterinary Research Institute (IVRI), Izatnagar, Uttar Pradesh, India

**Citation:** Mardiati Zain, Jurnida Rahman, Khasrad and Erpomen, 2016. Supplementation of *Saccharomyces cerevisiae* and *Sapindus rarak* in diet based of oil palm frond (OPF) on nutrient digestibility and daily weight gain of goat. Asian J. Anim. Vet. Adv., 11: 314-318.

**Corresponding Author:** Mardiati Zain, Department of Animal Nutrition, Faculty of Animal Science, Andalas University, Kampus Limau Manis, 25163 Padang, Indonesia

**Copyright:** © 2016 Mardiati Zain *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

**Competing Interest:** The authors have declared that no competing interest exists.

**Data Availability:** All relevant data are within the paper and its supporting information files.

## INTRODUCTION

Indonesia has a extensive oil palm plantation till now. According to Pertanian<sup>1</sup> Indonesia has 8,908,000 ha of oil palm plantations in 2011 and in 2014 has reached 10,210,892 ha. The plantation and industry of palm oil produced oil palm byproduct such as Oil Palm Fronds (OPF), Palm Kernel Cake (PKC), Palm Oil Sludge (POS)<sup>2</sup>. Oil palm fronds could be used as replace grasses in ration of ruminant in Indonesia. Based on study of Simanuhuruk *et al.*<sup>3</sup> production of palm frond oil can reach 40-50 fresh frond/tree/year with weights frond dry weight of 4.5 kg per frond. In one hectare of oil palm is expected to produce 6400-7500 fresh frond per year.

Oil Palm Fronds could be used as a source of fiber feed or as a component of ration for ruminants but it is contain low protein and high fiber content<sup>4,5</sup>. To maximize the utilization of available this feed resources as ruminant feed is needed the strategy to improve it's quality through processing by treated with 3% urea (ammoniation technique) as suggested of Zain *et al.*<sup>5</sup>. In addition to improving its use in rations also necessary effort to improved the microbial population in the rumen by adding rumen microbial growth factor.

Rumen microbial growth factor that can be used to increased rumen microbial population like as a defaunating agent that can reduced population of rumen protozoa as discussed by Zain *et al.*<sup>6</sup> and direct feed microbial to got an optimum condition for rumen bacterial growth as discussed elsewhere<sup>7,8</sup>. Number of protozoa in rumen must be reduced when using low quality feed such as oil palm fronds because protozoa would predation and digestion of rumen bacteria. Rumen protozoa can be reduced by used the *Sapindus rarak* fruit pericarp extract<sup>9</sup>. Direct fed microbial such as *Saccharomyces cerevisiae* supplementation on ruminant productive performance have been widely studied, but results are inconsistent. Cellulolytic activity of fibrolytic bacteria in the rumen fibrolytic bacteria in the rumen could stimulated with supplemented with yeast based study<sup>10-13</sup> improve feed intake and production of animal<sup>14,8</sup>. Therefore, the objective of our experiment was to determine whether supplementation of direct fed microbial like as *Saccharomyces cerevisiae* and defaunating agent (*Sapindus rarak*) could increase the nutrient digestibility and goat performance that feeding oil palm fronds ammoniated.

## MATERIALS AND METHODS

Sixteen male Ettawa goats (40+4.5 kg SEM) live weight equipped were randomly distributed in a 4×4 Completely

Table 1: Composition of feedstuffs to formulate complete feed in each treatment

Item	Diet (DM %)			
	A	B	C	D
Oil palm frond ammoniated	-	50.0	50.0	50.0
Grass	50.0	-	-	-
Rice brand	28.0	28.0	28.0	28.0
Corn	10.0	10.0	10.0	10.0
Coconute cake	11.0	11.0	11.0	11.0
Salt	0.5	0.5	0.05	0.05
Mineral	0.5	0.5	0.05	0.5
Total	100.0	100.0	100.0	100.0
Supplementation				
<i>Saccharomyces cerevisiae</i>	-	-	1	1
<i>Sapindus rarak</i>	-	-	-	4
<b>Nutrition (%)</b>				
Protein	12.66	12.01	12.01	12.01
TDN	64.73	63.83	63.83	63.83
Lemak	3.60	3.95	3.95	3.95
BETN	49.91	48.66	48.66	48.66
NDF	43.14	46.75	46.75	46.75
ADF	35.24	37.55	37.55	37.55

TDN: Total digestible nutrients, NDF: Neutral Detergent Fiber and ADF: Acid Detergent Fiber

Randomized Design (CRD). The four treatments were (A) native grass+concentrate (B), OPF (previously treated with 6% urea)+concentrate, (C) Diet B+1% *S. cerevisiae* and (D) Diet C+4% *Sapindus rarak*. Basal diet contained (dry basis) 50% OPF and 50% concentrate. The composition of feeds including concentrate is given in Table 1. The study was divided into two phases i.e., preliminary period and experiment period (data collection phase). In preliminary period, preparation of ration complete feed and male goat as material of research had done. Adaptation on farm environment and feed was done in two weeks and was continued by observation and data collection in 8 weeks.

Digestibility trials conducted using four animals for each treatment which were separated in individual pens. Goats were fed *ad-libitum* during the preliminary phase (15 days) and then restricted during the collection phase (last 6 days) at 90% of the intake feed that was offered at 7:00 and 16:00 h last 15 days of collection period. Animals were equipped with bags fitted to the animals with harness for total collection. During the collection period accurate records were kept for individual feed intake. Live weight was measured every two week. Total fecal excretion was collected once daily and 10% representative samples were dried at 60°C over night and kept in sealed bags until analysis. Feed and fecal were ground to pass through a 1 mm screen and composited. Dry matter, organic matter and nitrogen were analyzed by standard methods<sup>15</sup>. Neutral Detergent Fiber (NDF), Acids Detergent Fiber (ADF), cellulose were determined by procedures outlined by Goering and Van Soest<sup>16</sup>.

**Preparation of *Sapindus rarak* and *Saccharomyces cerevisiae*:**

Fruit of *Sapindus rarak* were dried in an oven at 60°C until they had of 90% dry matter constant. The whole fruits (including seed) were ground immediately after drying. *Saccharomyces cerevisiae* was obtained in Biotechnology Laboratory of Gajah Mada University, Yogyakarta Indonesia collection, strain Meyen ex Hansen) and it is contained  $4 \times 10^8$  live organisms/g, in medium on which it was grown. The composition of *Saccharomyces cerevisiae* medium were 4 g glucose, 2 g Yeast Extract Broth (YEB) and 3 g jelly powder. Mixed medium with 200 mL aquadest and inclusion on erlenmeyer glass. Autoclave on 121°C temperature, 1 atm pressure during 15 min. After 15 min remove the medium from autoclave and included in petri disk. Inoculation *Saccharomyces cerevisiae* product 1 mL in 99 mL aquadest. Prepared eight test tube for inoculated. Given test tube with code, 1 till 8. Filled test tube with 9 mL aquadest. After that, included 1 mL *Saccharomyces cerevisiae* on test tube 1, then took 1 mL from test tube 1 and removed in test tube 2, conducted it same treatment till test tube 8. Removed 1 mL from test tube 8 in petri disk that filled medium and let stand during 24 h. After 24 h, the total colonies of *Saccharomyces cerevisiae* counted in counting chamber.

**Statistical analysis:** Data were analyzed using the General Linear Model procedure in Statistical Analysis Software (SAS, version).

**RESULTS AND DISCUSSION**

**Digestibility of nutrient:** The result of effects addition of *S. cerevisiae* and *Sapindus rarak* on nutrient digestibility of ration based on oil palm frond ammoniated of goat are presented in Table 2. Effects of treatments were significant ( $p < 0.05$ ) increased dry matter, organic matter, NDF, ADF, cellulose and hemicellulose digestibility. Dry matter and organic matter digestibility in A, B, C and D treatments were 67.37, 50.74, 62.38 and 65.71% and organic matter digestibility were 70.20, 55.90, 66.33 and 67.64%, respectively. These result indicated that combination and *Saccharomyces cerevisiae* and *Sapindus rarak* in treatment D had higher dry matter and organic matter digestibility compared with treatment B and C.

This increased is likely due to the role of *Saccharomyces cerevisiae* can increased rumen microbial growth. More over increased dry matter digestibility and organic matter this may have indicated increased activity of rumen bacteria. The increasing activity of rumen microbial can level up degrading

Table 2: Digestibility nutrient of goat with experimental diet

Items	Diet			
	A	B	C	D
Dry matter digestibility (%)	67.37 <sup>a</sup>	50.74 <sup>d</sup>	62.38 <sup>c</sup>	65.71 <sup>b</sup>
Organic matter digestibility (%)	70.20 <sup>a</sup>	55.90 <sup>f</sup>	66.33 <sup>b</sup>	67.64 <sup>b</sup>
NDF digestibility (%)	60.22 <sup>a</sup>	47.90 <sup>f</sup>	54.34 <sup>b</sup>	60.21 <sup>a</sup>
ADF digestibility (%)	53.12 <sup>a</sup>	35.89 <sup>f</sup>	42.74 <sup>c</sup>	49.52 <sup>b</sup>
Cellulosa digestibility	64.43 <sup>a</sup>	43.84 <sup>f</sup>	53.97 <sup>b</sup>	59.31 <sup>ab</sup>
Hemicellulosa digestibility	64.79 <sup>a</sup>	50.66 <sup>f</sup>	57.70 <sup>b</sup>	65.55 <sup>ab</sup>

Means in the same row with different in their superscript differ ( $p < 0.05$ )

activity of organic matter into simple soluble compounds and it will lead to the increasing of organic substance absorption. This is in line with the study conducted by Paryad and Rashidi<sup>17</sup> who stated that the nutrient digestibility of goat and sheep rations supplemented with yeast was significantly increased compared with controls. *Saccharomyces cerevisiae* also had the capability to reduced oxygen so, the rumen environment become conducive for growth rumen bacteria especially cellulolytic bacteria as reported by Wallace<sup>18</sup>. Supplementation of *Sapindus rarak* would suppression of protozoa and may enhanced the flow of microbial protein from the rumen, increased the efficiency of feed utilization and improved the nutrition of the animal, provided that the loss of protozoa does not impair the fiber breakdown as reported by Newbold *et al.*<sup>19</sup>.

In this study, NDF, ADF and cellulose digestibility obtained was also significantly affected by *Saccharomyces cerevisiae* and *Sapindus rarak* supplementation. This is presumably because *Saccharomyces cerevisiae* that containing  $4 \times 10^8$  live microorganism/g have ability to degraded the fiber. NDF, ADF and cellulose that have potential as a source of energy for ruminants because they were fiber from the carbohydrate fraction. Supplementation of *Saccharomyces cerevisiae* and *Sapindus rarak* can increased activity of rumen microbial growth, then cases increased digestibility of NDF, ADF and cellulose. According to Fadel Elseed and Abusamra<sup>20</sup>, *Saccharomyces cerevisiae* could increased OM plus NDF digestibility compared good quality forages with control diet. Furthermore, supplementation single Direct Fed Microbial (DFM) of *Saccharomyces cerevisiae* was suitable to be used single or in combination with *Aspergillus oryzae* or *Bacillus amilolyquifaciens* could increased fermentability and degradability of ammoniated palm frond<sup>21</sup>. Some researchers<sup>22,23</sup>, have been reported that treatment with some yeast cultures increased the number of total and cellulolytic bacteria in the rumen and in some cases increased cellulose degradation. The result of Miller-Webster *et al.*<sup>22</sup>, showed that *Aspergillus oryzae* fermentation extract and *Saccharomyces cerevisiae* culture stimulated fiber digestion by ruminal



Table 3: Feed intake, average daily gain of goat with experimental diet

Items	Diet			
	A	B	C	D
Dry matter intake (g day <sup>-1</sup> )	993 <sup>a</sup>	734 <sup>b</sup>	828 <sup>ab</sup>	906 <sup>a</sup>
Organic matter intake (g day <sup>-1</sup> )	735 <sup>a</sup>	556 <sup>b</sup>	679 <sup>ab</sup>	720 <sup>a</sup>
Average daily gain (g day <sup>-1</sup> )	132 <sup>a</sup>	80 <sup>b</sup>	108 <sup>c</sup>	123 <sup>a</sup>

Means in the same row with different in their superscript differ (p<0.05)

microorganisms. Increased digestibility of NDF, ADF and cellulose with supplementation *Saccharomyces cerevisiae* indicated that supplementation could promote rumen cellulolytic bacteria<sup>12-24</sup>. Increasing the number of rumen cellulolytic caused *Saccharomyces cerevisiae* provided essential metabolite for its growth. The relatively stable rumen pH for growth of rumen microbes, especially cellulolytic bacteria causing growth to be better so the fiber fraction digestibility also increased. It is also reported by several researchers<sup>22,20,7</sup>.

**Feed intake and daily gain:** Feed intake and daily weight gain of this treatments are shown in Table 3. Dry matter, organic matter intake and average daily gain significantly affected of the treatments (p<0.05). This Table showed that supplementation of *Saccharomyces cerevisiae* can increased of nutrient intake and daily weight gain of goat when compared without *Saccharomyces cerevisiae* supplementation but still lower compared control diet. Supplementation of *Sapindus rarak* in treatment D could increased the nutrient intake, the same as with control diet and could improved the animal performance.

The positive effect of *Saccharomyces cerevisiae* in increasing nutrient intake and average daily gain of goat may be attributed to the increase of numbers of rumen total viable bacteria and cellulolytic bacteria<sup>25</sup>. Moreover, the stabilization of rumen environment could to be the reason for increased of numbers of rumen bacteria also may be related to pH modulation via reductions in lactic acid concentration<sup>26</sup>.

Supplementation *Sapindus rarak* in D treatment gave the best result compared with other treatment. This is likely due to the role of *Sapindus rarak* can release protozoa, thus release of protozoa can increase activity of rumen bacteria because existence of protozoa undesirable when goat received that high fiber feed. Existence of protozoa tend to prey rumen bacteria to viability of protozoa<sup>9</sup>. The daily DMI were ranged at 734-993 g/head/day. The lowest OMI was in the B. The high feed consumption was correlated with the high animal productivity and was influenced of digestibility. High nutrient digestibility will increased the nutrient intake.

Average Daily Gain (ADG) of goat was lower in this study than the study in indigenous rams fed complete feed by Purbowati *et al.*<sup>27</sup> who reported the average daily gain were 145.22-164.98 g. This might be caused by the different kinds and feedstuffs composition compose the complete feed. The main factor affected animal ADG is total nutrients consumed and biological value of feed. Feed consumption showed total energy and nutrient intake to promote the growth and animal production, so that if feed consumption is high, the ADG will also high.

## CONCLUSION

Oil palm fronds can be used to replace the field grass in formulation of complete feed for goat if supplemented with 1% *Saccharomyces cerevisiae* and 4% *Sapindus rarak*.

## ACKNOWLEDGMENTS

The authors are grateful for the support of National Strategic Research Grant by Directorate General Higher Education, Department of National Education Republic of Indonesia contract no 06/UN.16/PL-SN/2014. The study would not have been possible without the cooperation of Laboratorium Ruminant Nutrition of Faculty of Animal Science, Special thanks for my graduated student and technical assistance of Laboratory Ruminant Nutrition of Animal Science, Faculty of Andalas University.

## REFERENCES

- Pertanian, K., 2014. Agricultural Statistics 2014. Ministry of Agriculture Republic of Indonesia, Jakarta, Indonesia.
- Elisabeth, J. and S.P. Ginting, 2003. [Utilization of oil palm industry by-products as feedstuffs for beef cattle]. Proceedings of the National Workshop on Systems Integration Palm Oil-Ox, September 9-10, 2003, Bengkulu, Indonesia, pp: 110-119.
- Simanuhuruk, K. Junjungan and A. Tarigan, 2007. Utilization of oil palm fronds as basal feed for Kacang goats on growing phase. Proceedings of the Seminar Nasional Teknologi Peternakan dan Veteriner, August 21-22, 2007, Sungei Putih, Galang.
- Oshio, S., O. Abu Hassan, A. Takigawa, D. Mohd. Jaafar, A. Abe, I. Dahlan and N. Nakanishi, 1990. Processing and utilization of oil palm by-products for ruminants. MARDI/TARC @ JIRCAS Collaborative Study Report, pp: 110.
- Zain, M., J. Rahman and Khasrad, 2014. Effect of palm oil by products on *in vitro* fermentation and nutrient digestibility. Anim. Nutr. Feed Technol., 14: 175-181.

6. Zain, M., T. Sutardi, Suryahadi and N. Ramli, 2008. Effect of defaunation and supplementation methionine hydroxy analogue and branched chain amino acid in growing sheep diet based on palm press fiber ammoniated. Pak. J. Nutr., 7: 813-816.
7. Zain, M., J. Rahman, Khasrad and Erpomen, 2015. *In vitro* fermentation characteristics of palm oil byproducts which is supplemented with growth factor rumen microbes. Pak. J. Nutr., 14: 625-628.
8. Herawaty, R., N. Jamarun, M. Zain, Anim and R.W.S. Ningrat, 2013. Effect of supplementation *Sacharomyces cerevisiae* and *Leucaena leucocephala* on low quality roughage feed in beef cattle diet. Pak. J. Nutr., 12: 182-184.
9. Benchaar, C., T.A. McAllister and P.Y. Chouinard, 2008. Digestion, ruminal fermentation, ciliate protozoal populations and milk production from dairy cows fed cinnamaldehyde, Quebracho condensed tannin, or *Yucca schidigera* Saponin extracts. J. Dairy Sci., 91: 4765-4777.
10. Thrune, M., A. Bach, M. Ruiz-Moreno, M.D. Stern and J.G. Linn, 2009. Effects of *Saccharomyces cerevisiae* on ruminal pH and microbial fermentation in dairy cows: Yeasts supplementation on rumen fermentation. Livestock Sci., 124: 261-265.
11. Fereli, F., A.F. Branco, C.C. Jobim, S.M. Coneglian, F. Granzotto and J.C. Barreto, 2010. Sodium monensin and *Saccharomyces cerevisiae* in cattle diets: Ruminal fermentation, nutrient digestibility and microbial synthesis efficiency. Revista Brasileira de Zootecnia, 39: 183-190.
12. Zain, M., N. Jamarun, A. Anim, R.W.S. Ningrat and R. Herawati, 2011. Effect of yeast (*Saccharomyces cerevisiae*) on fermentability, microbial population and digestibility of low quality roughage *in vitro*. Archiva Zootechnica, 14: 51-58.
13. Jurkovich, V., E. Brydl, J. Kutasi, A. Harnos and P. Kovacs *et al.*, 2014. The effects of *Saccharomyces cerevisiae* strains on the rumen fermentation in sheep fed with diets of different forage to concentrate ratios. J. Applied Anim. Res., 42: 481-486.
14. Tripathi, M.K. and S.A. Karim, 2011. Effect of yeast cultures supplementation on live weight change, rumen fermentation, ciliate protozoa population, microbial hydrolytic enzymes status and slaughtering performance of growing lamb. Livestock Sci., 135: 17-25.
15. AOAC., 2007. Official Method of Analysis. 18th Edn., Association of Official Analytical Chemist, Gaithersburg, MD, USA.
16. Goering, H.K. and P.J. Van Soest, 1970. Forage Fiber Analysis (Apparatus, Reagents, Procedure and Some Applications). In: Agricultural Handbook, 379, US Department of Agricultural (Ed.). US Government Printing Office, Washington, DC., pp: 1-20.
17. Paryad, A. and M. Rashidi, 2009. Effect of yeast (*Saccharomyces cerevisiae*) on apparent digestibility and nitrogen retention of tomato pomace in sheep. Pak. J. Nutr., 8: 273-278.
18. Wallace, R.J., 1994. Ruminal microbiology, biotechnology and ruminant nutrition: Progress and problems. J. Anim. Sci., 72: 2992-3003.
19. Newbold, C.J., S.M. El Hassan, J. Wang, M.E. Ortega and R.J. Wallace, 1997. Influence of foliage from African multipurpose trees on activity of rumen protozoa and bacteria. Br. J. Nutr., 78: 237-249.
20. Fadel Elseed, A.M.A. and M.R. Abusamra, 2007. Effects of supplemental yeast (*Saccharomyces cerevisiae*) culture on NDF digestibility and rumen fermentation of forage sorghum hay in Nubian goat's kids. Res. J. Agric. Biol. Sci., 3: 133-137.
21. Suryani, H., M. Zain, R.W.S. Ningrat and N. Jamarun, 2016. Supplementation of Direct Fed Microbial (DFM) on *in vitro* Fermentability and degradability of ammoniated palm frond. Pak. J. Nutr., 15: 90-95.
22. Miller-Wester, T., W.H. Hoover, M. Holt and J.E. Nocek, 2002. Influence of yeast culture on ruminal microbial metabolism in continuous culture. J. Dairy Sci., 85: 2009-2014.
23. Newbold, C.J., P.E.V. Williams, N. Mckain, A. Walker and R.J. Wallace, 1990. The effects of yeast culture on yeast numbers and fermentation in the rumen of sheep. Proc. Nutr. Soc., 49: 47-47.
24. Callaway, E.S. and S.A. Martin, 1997. Effects of a *Saccharomyces cerevisiae* culture on ruminal bacteria that utilize lactate and digest cellulose. J. Dairy Sci., 80: 2035-2044.
25. Marghany, M., M.A. Sarhan, A. Abd. El-Hey and A.A.H. El-Tahan, 2005. Performance of lactating buffaloes fed rations supplemented with different levels of baker's yeast (*Saccharomyces cerevisiae*). Egypt. J. Nutr. Feed, 8: 21-34.
26. El-Waziry, A.M., H.E.M. Kamel and M.H.M. Yacout, 2000. Effect of bakery's yeast (*Saccharomyces cerevisiae*) supplementation to berseem (*Trifolium alexandrinum*) hay diet on protein digestion and rumen fermentation of sheep. Egypt. J. Nutr. Feeds, 3: 71-82.
27. Purbowati, E., C.I. Sutrisno, E. Baliarti, S.P.S. Budhi and W. Lestariana, 2008. [Energy utilization of Complete feed with different protein-energy levels in male local lamb on feedlots system]. J. Pengembangan Peternakan Tropis, 33: 59-65.

# Jurnal Ajava\_2016\_Zain et al

---

## ORIGINALITY REPORT

---

**13%**

SIMILARITY INDEX

**20%**

INTERNET SOURCES

**12%**

PUBLICATIONS

**0%**

STUDENT PAPERS

---

## MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

6%

★ **studylib.net**

Internet Source

---

Exclude quotes  On

Exclude bibliography  Off

Exclude matches  < 3%